

VU Learning HUB

BIO301 Quiz File

Chapter-wise Solved MCQs

vulearninghub.xyz

Lesson 1: What is Genetics?

1. Genetics is defined as the study of which of the following?

- A) Cells, tissues, and organs
- B) Genes, heredity, and variation
- C) Proteins, enzymes, and lipids
- D) Bacteria, viruses, and fungi

2. Genetics is considered a sub-field of which broader discipline?

- A) Chemistry
- B) Physics
- C) Biology
- D) Geology

3. Who explained the fundamental principles of heredity in 1866?

- A) Strausberger
- B) Gregor Mendel
- C) Waldeyer
- D) T. H. Morgan

4. Which plant did Gregor Mendel use as an experimental plant to formulate the laws of heredity?

- A) Sweet pea
- B) Garden pea
- C) Sunflower
- D) Snapdragon

Answer Key

1. B — Genetics is specifically defined as the study of genes, heredity, and variation
2. C — Genetics is categorized as a field of Biology
3. B — Gregor Mendel explained the core principles of heredity in the year 1866
4. B — Mendel utilized the garden pea as his experimental plant for formulating inheritance laws

Lesson 2: Sub Types of Genetics

1. Which sub-discipline of genetics deals with the movement of genes and genetic traits from parents to offspring, Mendel's Laws, and genetic recombinations?

- A) Population Genetics
- B) Transmission (Classical) Genetics
- C) Quantitative Genetics
- D) Molecular Genetics

2. Which branch of genetics studies traits in a group of population when those traits are determined by one or a few genes?

- A) Molecular Genetics
- B) Quantitative Genetics
- C) Population Genetics
- D) Transmission Genetics

3. Quantitative genetics specifically studies group heredity for traits that are determined by:

- A) A single gene only
- B) Molecular structure of DNA
- C) Many genes simultaneously
- D) Extracellular enzymes

4. What is the primary focus of Molecular Genetics?

- A) Mendel's Laws of inheritance
- B) Traits determined by one or a few genes in a population
- C) The molecular structure and function of genes
- D) Physical traits like skin color and height

Answer Key

1. B — Transmission (Classical) Genetics deals with the movement of genes from parents to offspring, Mendel's Laws, and recombinations.
2. C — Population genetics studies traits in a group of population when determined by one or a few genes.
3. C — Quantitative genetics studies group heredity for traits determined by many genes simultaneously (e.g., skin color, height).
4. C — Molecular genetics deals specifically with the molecular structure and function of genes.

Lesson 3: Common Genetics Terminologies**1. How is a "character" defined in genetics?**

- A) A variant for a specific feature
- B) A heritable feature such as skin color or height
- C) The physical appearance of a disease
- D) An offspring resulting from a hybrid cross

2. What is a "trait"?

- A) A heritable feature
- B) The total genetic makeup of an organism
- C) A variant for a character (e.g., brown, black, white)
- D) The cross between two different species

3. A cross between a true-breed plant or animal with another true-breed plant or animal is called a:

- A) Monohybrid test cross
- B) Pure cross
- C) Dihybrid reciprocal cross
- D) Back cross

4. The first filial generation resulting from a parental cross is denoted as:

- A) P generation
- B) F1 generation
- C) F2 generation
- D) F3 generation

Answer Key

1. B — A character is defined as a heritable feature such as skin color or height

2. C — A trait represents a specific variant for a character, such as brown or white color
3. B — A pure cross refers to crossing a true-breed organism with another true-breed organism
4. B — The first filial generation derived from the parent (P) generation is termed the F1 generation

Lesson 4: Mendel's Law of Dominance

1. According to Mendel's law of dominance, what happens in a cross of parents pure for contrasting traits?

- A) Both traits blend equally in the next generation
- B) Only one form of the trait appears in the next generation
- C) Neither form of the trait appears
- D) A completely novel phenotype is created

2. In a monohybrid cross of true-breeding plants, what happens to one version of the trait in the F1 generation?

- A) It duplicates
- B) It disappears
- C) It mutates
- D) It becomes co-dominant

3. When true-breeding purple-flowered pea plants are crossed with white-flowered pea plants, all F1 plants have purple flowers because:

- A) Purple is recessive to white
- B) Purple is dominant to white
- C) White flower genes are completely destroyed
- D) Environment altered flower pigments

4. Mendel's law of dominance specifically describes the behavior of contrasting traits during which type of cross?

- A) Monohybrid cross
- B) Trihybrid cross
- C) Test cross
- D) Pedigree cross

Answer Key

1. B — Mendel's law of dominance states that in a cross of pure parents with contrasting traits, only one trait form appears in the next generation
2. B — In a monohybrid cross, one version of the contrasting trait disappears in the F1 generation
3. B — Purple color is dominant to white color, causing all F1 offspring to exhibit purple flowers
4. A — The law of dominance illustrates trait expression in a monohybrid cross

Lesson 5: Monohybrid cross and Dihybrid cross

1. Cross-fertilization of true-breeding plants that differ in just one character is defined as a:

- A) Dihybrid cross
- B) Monohybrid cross
- C) Trihybrid cross
- D) Testcross

2. What did Gregor Mendel determine using a monohybrid cross?

- A) The segregation of alleles at a single gene locus
- B) The physical length of chromosomes in micrometers
- C) The G-C base ratio across species
- D) The rate of histone acetylation

3. What is the primary purpose of conducting a dihybrid cross?

- A) To count the number of centrioles in dividing cells
- B) To determine whether alleles at two different gene loci segregate dependently or independently
- C) To measure the water content of active nuclei
- D) To observe polytene chromosome puffs

4. How are two characters transmitted from parents to offspring when their genes are located on different chromosomes?

- A) As a linked package
- B) Independently
- C) As a single fused allele
- D) Exclusively through maternal mitochondria

Answer Key

- 1. B — A monohybrid cross involves cross-fertilization of true-breeding plants differing in a single character.
- 2. A — Mendel used monohybrid crosses to demonstrate the segregation of alleles at a single gene locus.
- 3. B — Dihybrid crosses evaluate whether alleles at two distinct gene loci segregate dependently or independently.
- 4. B — Characters controlled by genes on separate chromosomes transmit independently to offspring.

Lesson 6: Law of Segregation**1. According to the Law of Segregation, what happens to the two alleles responsible for a trait during gamete formation?**

- A) They mutate into new alleles
- B) They separate from each other
- C) They fuse permanently
- D) They remain bound to histones

2. What process recombines or reunites the separated alleles to form a new genotype in offspring?

- A) Transcription
- B) Fertilization
- C) Translation
- D) Cytokinesis

3. According to the Modern Concept of Segregation, when do the two alleles of a gene separate?

- A) During fertilization
- B) During gametogenesis
- C) During DNA replication in S phase
- D) During organogenesis

4. In a cross, what determines the resulting phenotype of an offspring after gametes reunite?

- A) The environmental temperature alone
- B) The resulting genotype formed during fertilization
- C) The size of the ribosome
- D) Random chance without genetic influence

Answer Key

1. B — The Law of Segregation specifies that the two alleles for a trait separate from each other during gamete formation.
2. B — Separated alleles are recombined during the process of fertilization to form new genotypes.
3. B — Under the modern concept, alleles separate specifically during gametogenesis.
4. B — Gametes combine during fertilization to yield a new genotype, which determines the resulting phenotype.

Lesson 7: Mendel's Law of Independent Assortment**1. What does Mendel's Law of Independent Assortment state regarding alleles for different traits?**

- A) They are always inherited as a single linked package
- B) They are distributed to sex cells independently of one another
- C) They destroy one another during meiotic prophase
- D) They blend together to produce a single uniform phenotype

2. In Mendel's dihybrid cross example with pod color and seed color, which traits were dominant?

- A) Yellow pod color and green seed color
- B) Green pod color and yellow seed color
- C) Yellow pod color and yellow seed color
- D) Green pod color and green seed color

3. What phenotypic ratio did Mendel observe in the F₂ generation after self-pollinating F₁ dihybrid plants?

- A) 3:1
- B) 1:2:1
- C) 9:3:3:1
- D) 1:1:1:1

4. The 9:3:3:1 phenotypic ratio in the F₂ generation of a dihybrid cross supports which hypothesis?

- A) Dependent segregation unit hypothesis
- B) Independent assortment of alleles
- C) Acquired environmental variation
- D) Maternal mitochondrial inheritance

Answer Key

1. B — The Law of Independent Assortment states that alleles for different traits distribute to sex cells independently.
2. B — In the source example, green pod color (GG) and yellow seed color (YY) are dominant traits.

3. C — Mendel observed a classic 9:3:3:1 phenotypic ratio among the F2 offspring of a dihybrid cross.
4. B — The 9:3:3:1 ratio supports the hypothesis that alleles for different traits sort into gametes independently.

Lesson 8: Modern Genetics Concepts

1. How is genetic material organized in prokaryotic cells compared to eukaryotic cells?

- A) Prokaryotes have multiple linear chromosomes, while eukaryotes have one circular chromosome
- B) Prokaryotes generally have one circular chromosome, while eukaryotes have linear chromosomes in their nuclei
- C) Both prokaryotes and eukaryotes possess linear chromosomes only
- D) Prokaryotes lack chromosomes entirely and rely solely on RNA

2. In modern genetics, a gene is defined as which of the following?

- A) A complete cell membrane structure
- B) A heredity unit located on chromosomes
- C) A protein enzyme in the cytoplasm
- D) A carbohydrate energy polymer

3. Which of the following is listed as an example of a Non-Mendelian inheritance pattern?

- A) Monohybrid 3:1 dominance
- B) Mitochondrial inheritance
- C) Pure-breed monohybrid segregation
- D) Dihybrid 9:3:3:1 independent assortment

4. In genetics, how is a mutation broadly defined, and what impacts can it have?

- A) It strictly indicates a lethal disease in all cases
- B) It indicates "a change" and can be harmful, beneficial, or have no effect
- C) It refers only to changes caused by cold temperature
- D) It is an acquired physical training adaptation

Answer Key

1. B — Prokaryotes typically have a single circular chromosome, whereas eukaryotes have linear chromosomes in their nucleus.
2. B — A gene is defined in modern genetics as a unit of heredity located on chromosomes.
3. B — Non-Mendelian inheritance includes patterns such as mitochondrial inheritance, linkage, maternal effects, and epigenetics.
4. B — In genetics, mutation indicates a change that may be harmful, beneficial, or neutral.

Lesson 9: Importance of Genetics

1. Which of the following is listed as one of the three core characteristics of human genetic data?

- A) Temporary
- B) Personal
- C) Changeable
- D) Anonymous

2. Genetic information is described as Personal, Predictive, and:

- A) Permanent

- B) Reversible
- C) Environmental
- D) Diagnostic

3. Genetics assists medical science in which of the following areas?

- A) Newborn and population screening
- B) Pharmacogenetics and individualized medicine
- C) Predictive planning and diagnostic treatments
- D) All of the above

4. How does genetics contribute to our understanding of biological history?

- A) It proves traits do not change
- B) It helps us understand the process of evolution
- C) It eliminates environmental influences
- D) It prevents random mutations

Answer Key

- 1. B — Genetic data is characterized as Personal, Permanent, and Predictive
- 2. A — Genetic data possesses three key characteristics: Personal, Permanent, and Predictive
- 3. D — Genetics aids diagnostics, treatments, screening, and individualized pharmacogenetics
- 4. B — Genetics helps scientists understand the process and mechanisms of evolution

Lesson 10: What is Heredity?

1. What is the fundamental definition of heredity?

- A) The change in species over evolutionary time
- B) The transmission of traits from parents to offspring
- C) The environmental modification of physical organs
- D) The breakdown of proteins during digestion

2. Gregor Mendel demonstrated that inheritance occurs through which model rather than blending?

- A) Blending inheritance
- B) Particulate inheritance
- C) Acquired inheritance
- D) Environmental inheritance

3. According to the particulate theory of inheritance, how are characters (alleles) passed from parents to offspring?

- A) They are blended and permanently diluted
- B) They are inherited unchanged
- C) They are destroyed during fertilization
- D) They are altered by physical exercise

4. What did the observed pattern of inheritance of unchanged characters provide in early genetics?

- A) The first definition of a gene
- B) The discovery of electron microscopy
- C) The proof that DNA lacks nitrogenous bases

D) The complete classification of all animal species

Answer Key

1. B — Heredity is defined as the transmission of traits from parents to offspring.
2. B — Mendel showed that inheritance follows a particulate model rather than a blending model.
3. B — In particulate inheritance, hereditary units (alleles) are transmitted unchanged across generations.
4. A — The inheritance pattern of unchanged particulate characters provided the first biological definition of a gene.

Lesson 11: Concept of variation

1. How is biological variation defined among living organisms?

- A) Differences seen among individuals of the same species
- B) Structural differences between completely different kingdoms
- C) The exact duplication of parental physical traits
- D) Chemical changes in non-living minerals

2. What two primary factors influence variations seen in individuals of the same species?

- A) Atmospheric pressure and soil pH only
- B) Genotype and environment
- C) Ribosome count and tRNA length
- D) Gravitational pull and magnetic fields

3. In the handout's example of snail shells, which variations are observed among individuals?

- A) Length of antennae and eye size
- B) Shell color, presence/absence of bands, and number of bands
- C) Number of legs and internal lung capacity
- D) Thickness of the foot muscle alone

4. Which of the following is listed in the handout as an example of variation in humans (Homo sapiens)?

- A) Presence of circular prokaryotic plasmids
- B) Variations in hair curliness, eye color, and skin color
- C) Presence of polytene giant chromosomes
- D) Ability to perform photosynthesis in warm sunlight

Answer Key

1. A — Variation refers to the observed differences among individuals belonging to the same species.
2. B — Variation among individuals arises due to their genetic makeup (genotype), environment, or a combination of both.
3. B — Snail shell variations include shell color, presence or absence of bands, and band count.
4. B — Human variations highlighted in the text include skin color, hair color, eye color, hair curliness, and sex.

Lesson 12: Genetic and Acquired Variation

1. Inherited variations result directly from the activity of:

- A) Environmental factors
- B) Genes

- C) Nutritional habits
- D) Physical training

2. Which of the following is an example of an inherited variation?

- A) Language spoken
- B) Athletic skills
- C) Blood groups
- D) Obesity

3. Acquired variations result from an individual's activities, nutrition, or:

- A) Gene activity
- B) Environmental conditions during a lifetime
- C) Meiotic recombination
- D) Chromosomal nondisjunction

4. Can acquired characteristics be inherited by offspring?

- A) Yes, always
- B) Yes, if beneficial
- C) No, they cannot be inherited
- D) Only under cold conditions

Answer Key

- 1. B — Inherited variations are genetically controlled and result from gene activity
- 2. C — Blood groups, fingerprints, and sex are genetically controlled inherited variations
- 3. B — Acquired variations arise from an individual's activities, diet, or lifetime environmental conditions
- 4. C — Acquired characteristics cannot be inherited by subsequent generations

Lesson 13: Types of variation: Two types of Variation

1. Which type of variation is entirely genetically controlled and cannot be altered by external environmental conditions?

- A) Continuous variation
- B) Discontinuous variation
- C) Acquired variation
- D) Environmental variation

2. Which of the following human traits is an example of discontinuous variation?

- A) Height
- B) Weight
- C) Blood group
- D) Skin color gradient

3. How is continuous variation characterized regarding gene control and environmental influence?

- A) Controlled by a single gene and unaffected by the environment
- B) Controlled by a lot of genes and strongly influenced by the environment
- C) Caused exclusively by physical exercise
- D) Unaffected by both genes and environment

4. When plotted on a graph, measurements of continuous variation (such as human height) typically yield what type of curve?

- A) Disjointed bar chart with separated bars
- B) Smooth bell-shaped curve
- C) Flat horizontal line
- D) Exponential upward curve

Answer Key

1. B — Discontinuous variation is entirely genetically controlled and unaffected by external conditions.
2. C — Human blood groups represent discontinuous variation with distinct qualitative categories.
3. B — Continuous variation is polygenic (controlled by many genes) and strongly influenced by environmental factors.
4. B — Continuous variation measurements, such as height, produce a smooth bell-shaped curve when graphed.

Lesson 14: Source of Variation

1. Which of the following is categorized as a genetic cause of variation?

- A) Nutrients
- B) Meiosis
- C) Temperature
- D) Physical training

2. Sexual reproduction contributes to genetic variation primarily through:

- A) Environmental adaptation
- B) Recombination of genetic material
- C) Acquired physical training
- D) Temperature shifts

3. Which of the following is an environmental cause of variation?

- A) Meiosis
- B) Mutations
- C) Temperature
- D) Sexual reproduction

4. Which type of variation acts as the raw material for natural selection?

- A) Acquired variation
- B) Environmental variation
- C) Inherited variation
- D) Nutritional variation

Answer Key

1. B — Meiosis, sexual reproduction, and mutations are genetic causes of variation
2. B — Sexual reproduction enables the recombination of genetic material in new ways
3. C — Temperature, nutrients, drugs, and training are environmental causes of variation
4. C — Inherited variation serves as the fundamental raw material for natural selection

Lesson 15: Heredity and variation in Mendel's work

1. In Mendel's work, what was demonstrated when pure-breeding tall plants were self-pollinated?

- A) They produced a mix of tall and dwarf plants in equal numbers
- B) They produced only tall offspring
- C) They produced only purple flowers
- D) They failed to produce any offspring

2. What key evidence of heredity was shown by Mendel's pure-breeding purple-flowered plants?

- A) They consistently produced white-flowered plants
- B) They produced only purple-flowered offspring
- C) They produced plants with variable yellow seeds
- D) They changed color depending on soil temperature

3. According to the conclusions in Lesson 15, how is biological variation inherited in nature?

- A) Solely by random chance
- B) According to genetic laws
- C) Exclusively through environmental trauma
- D) By direct physical training of parents

4. How does Mendel's work address the occurrence of variation in natural populations?

- A) Variation is extremely rare and artificial
- B) Variation is widespread in nature and follows genetic rules
- C) Variation only occurs in bacterial plasmids
- D) Variation is impossible in true-breeding lines

Answer Key

- 1. B — In Mendel's experiments, pure-breeding tall plants consistently produced only tall offspring.
- 2. B — Pure-breeding purple-flowered plants demonstrated heredity by producing only purple-flowered offspring.
- 3. B — Variation is inherited according to established genetic laws rather than solely by chance.
- 4. B — Mendel's work showed that variation is widespread in nature and operates under defined genetic principles.

Lesson 16: Mendelian and Non-Mendelian inheritance

1. In Gregor Mendel's classic model of inheritance, how many alleles control a single trait?

- A) Multiple alleles exceeding ten
- B) Exactly two alleles with a clear dominant-recessive relationship
- C) One allele per chromosome pair only
- D) Zero alleles, as traits are non-genetic

2. Mendelian inheritance specifically applies to genes that obey which principles?

- A) The laws of dominance, segregation, and independent assortment
- B) The rules of acquired inheritance and physical training
- C) The principles of cytoplasmic organelle duplication
- D) The laws of thermodynamics and fluid dynamics

3. How is Non-Mendelian inheritance defined?

- A) Patterns of inheritance that strictly follow the 3:1 monohybrid ratio
- B) Patterns of inheritance that deviate from Mendelian patterns
- C) Traits that are expressed only in bacterial cells
- D) Traits that cannot be passed to offspring under any condition

4. Which of the following is listed in the handout as an example of Non-Mendelian inheritance?

- A) Autosomal dominant monohybrid inheritance
- B) Mitochondrial inheritance and genomic imprinting
- C) Pure-breeding purple flower inheritance
- D) Complete dominance in garden peas

Answer Key

1. B — Mendel's classic model assumes each trait is controlled by a single gene with two alleles showing a dominant-recessive relationship.
2. A — Mendelian inheritance involves genes obeying Mendel's laws of dominance, segregation, and independent assortment.
3. B — Non-Mendelian inheritance refers to inheritance patterns that deviate from standard Mendelian rules.
4. B — Examples of Non-Mendelian inheritance include mitochondrial inheritance, genomic imprinting, incomplete dominance, co-dominance, and epigenetics.

Lesson 17: Conclusions of Mendelian Inheritance**1. Mendel's conclusions regarding inheritance were summarized into how many laws?**

- A) Two
- B) Three
- C) Four
- D) Five

2. In Mendel's experiments, smooth seeds (S) showed what relationship to wrinkled seeds (s)?

- A) Incompletely dominant
- B) Completely dominant
- C) Co-dominant
- D) Epistatic

3. The Law of Segregation states that two members of a gene pair separate during:

- A) Fertilization
- B) Formation of gametes
- C) Mitotic anaphase
- D) DNA packaging

4. The Law of Independent Assortment asserts that alleles for different traits:

- A) Always remain together on the same strand
- B) Assort independently of one another
- C) Mutate simultaneously
- D) Blend in equal ratios

Answer Key

1. B — Mendel's conclusions were formulated into three laws: Dominance, Segregation, and Independent Assortment
2. B — Smooth seeds (S) are completely dominant over wrinkled seeds (s)
3. B — Gene pairs segregate from each other during gamete formation
4. B — Alleles for different traits assort independently during meiosis

Lesson 18: Exceptions to Mendelian inheritance

1. Which exception to Mendelian inheritance occurs when one gene masks or hides the expression of another gene?

- A) Pleiotropy
- B) Epistasis
- C) Incomplete dominance
- D) Monohybrid segregation

2. What is defined as a phenomenon where a single gene produces multiple distinct phenotypic effects?

- A) Polygenic inheritance
- B) Pleiotropy
- C) Co-dominance
- D) Complete dominance

3. When a single phenotypic trait is influenced and determined by the combined action of multiple genes, it is called:

- A) Monohybrid cross
- B) Polygenic inheritance
- C) Incomplete dominance
- D) Sex linkage

4. Which of the following is included in the handout's list of exceptions to Mendelian inheritance?

- A) Environmental effects on gene expression and linkage
- B) Law of Segregation
- C) Pure-breeding monohybrid crosses
- D) Law of Dominance

Answer Key

1. B — Epistasis is an exception to Mendelian inheritance where one gene masks the expression of another gene.
2. B — Pleiotropy occurs when a single gene exerts multiple distinct phenotypic effects.
3. B — Polygenic inheritance refers to a single phenotypic trait being controlled by multiple genes.
4. A — Linkage, sex linkage, and environmental effects on gene expression are listed as exceptions to Mendelian inheritance.

Lesson 19: Incomplete dominance

1. In incomplete dominance, what type of phenotype is produced in heterozygous individuals?

- A) The phenotype of the dominant parent only

- B) The phenotype of the recessive parent only
- C) An intermediate phenotype that is a blend of both alleles
- D) A completely unexpressed, lethal phenotype

2. In Japanese 4 o'clock flowers, crossing a red-flowered plant (RR) with a white-flowered plant (WW) produces offspring with which phenotype?

- A) All red flowers
- B) All white flowers
- C) All pink flowers
- D) A 3:1 ratio of red to white flowers

3. What is the genotype of the pink-flowered Japanese 4 o'clock plant?

- A) RR
- B) WW
- C) RW
- D) rr

4. Why does a "new" blended phenotype appear in incomplete dominance?

- A) Because one allele completely destroys the other allele
- B) Because neither allele is completely dominant over the other
- C) Because environmental temperature alters DNA bases
- D) Because chromosomes fail to segregate during meiosis

Answer Key

- 1. C — Incomplete dominance results in an intermediate phenotype that blends the traits of both alleles.
- 2. C — Crossing red (RR) and white (WW) Japanese 4 o'clock plants yields heterozygous offspring (RW) with pink flowers.
- 3. C — The pink-flowered Japanese 4 o'clock plant has the heterozygous genotype RW.
- 4. B — A blended phenotype appears because neither allele is dominant over the other.

Lesson 20: Co-dominance

1. How is co-dominance defined in genetics?

- A) One allele completely masks the expression of the other allele
- B) Both alleles are simultaneously expressed in the heterozygous individual
- C) The dominant allele blends with the recessive allele to form an intermediate phenotype
- D) Both alleles are completely suppressed in heterozygotes

2. In roan cattle inheritance, what coat phenotype is produced by the heterozygous genotype (Rr)?

- A) Pure red hairs
- B) Pure white hairs
- C) Roan coat (a mix of red and white hairs)
- D) Golden yellow fur

3. In the Appaloosa horse example, what is the phenotype of the heterozygous individual?

- A) A pure white coat
- B) A pure dark coat
- C) A white horse with gray spots

D) A black horse with white mane

4. Which human trait is cited in Lesson 20 as an example of co-dominance?

- A) Eye color
- B) Blood group inheritance (ABO)
- C) Skin pigmentation
- D) Hair texture

Answer Key

1. B — Co-dominance occurs when both alleles appear and are expressed in heterozygous individuals.
2. C — In roan cattle, the heterozygous genotype (Rr) produces a roan coat, which is a mix of red and white hairs.
3. C — In Appaloosa horses, the heterozygous phenotype is a white horse with gray spots.
4. B — Human blood group inheritance is cited as an example of co-dominance.

Lesson 21: Polygenic traits

1. How are polygenic traits defined in genetics?

- A) Traits controlled by a single gene locus with two alleles
- B) Traits controlled by the combined interaction of many gene loci
- C) Traits inherited exclusively through sex chromosomes
- D) Traits produced by point mutations in intronic sequences

2. Which human feature is explicitly listed in Lesson 21 as an example of a polygenic trait?

- A) ABO blood group
- B) Human height
- C) Color blindness
- D) Albinism

3. In contrast to single-gene Mendelian traits, polygenic traits typically exhibit:

- A) Only two discrete non-overlapping phenotypes
- B) Continuous variation across a range of phenotypic values
- C) Complete embryonic lethality
- D) Strict sex-linked inheritance

4. Qualitative and quantitative traits that show continuous variation are usually described as being:

- A) Polygenic in nature
- B) Mono-allelic
- C) Acquired variations
- D) Epigenetically silenced

Answer Key

1. B — Polygenic traits are determined by the combined interaction of multiple gene loci.
2. B — Human height is cited in Lesson 21 as a primary example of a polygenic trait.
3. B — Combined interactions across multiple loci yield continuous variation in phenotypes.
4. A — Qualitative and quantitative traits exhibiting continuous variation are polygenic in nature.

Lesson 22: Epistasis

1. What is the definition of epistasis as presented in Lesson 22?

- A) A single gene controlling multiple distinct physical traits
- B) Alleles at one gene locus hiding or preventing the expression of alleles at a second gene locus
- C) The failure of homologous chromosomes to separate during meiosis
- D) A point substitution resulting in a silent amino acid change

2. In Labrador retrievers, what coat color is specified by the dominant allele B when pigment deposition is active?

- A) Brown (Chocolate)
- B) Black
- C) Yellow
- D) White

3. What coat color phenotype results when a Labrador retriever is homozygous recessive (ee) at the pigment deposition locus?

- A) Black fur
- B) Brown fur
- C) Yellow fur, regardless of the alleles present at the first locus
- D) Spotted fur

4. What is the genotype of a brown (chocolate) Labrador retriever according to the lesson examples?

- A) BBEE or BbEe
- B) bbEE or bbEe
- C) BBee or Bbee
- D) bbee

Answer Key

1. B — Epistasis occurs when alleles at one gene locus prevent or mask the expression of alleles at another locus.
2. B — The dominant allele B specifies black coat pigment in Labrador retrievers.
3. C — Homozygous recessive `ee` blocks pigment deposition in fur, producing yellow fur regardless of the B locus.
4. B — Genotypes `bbEE` and `bbEe` combine brown pigment (`bb`) with functional deposition (`E\_`) to yield brown fur.

Lesson 23: Pleiotropy

1. How is pleiotropy defined in genetics?

- A) The combined interaction of many genes controlling a single trait
- B) The ability of a single gene locus/allele to affect an organism in many ways by producing more than one trait
- C) The exchange of genetic material between non-sister chromatids
- D) The duplication of an entire chromosome set during mitosis

2. Which group of human genetic conditions is cited in Lesson 23 as examples of pleiotropy?

- A) Sickle cell anemia, Cystic fibrosis, and Huntington disease

- B) Down syndrome, Turner syndrome, and Klinefelter syndrome
- C) Hemophilia A and Red-green color blindness
- D) Scurvy, Rickets, and Beriberi

3. In the Huntington disease example of pleiotropy, what two distinct phenotypic effects stem from the single gene mutation?

- A) Short stature and webbed neck
- B) Progressive dementia (at age 40–50) and death ~5 years after disease onset
- C) Extra fingers and fused toes
- D) Blindness and deafness

4. How does pleiotropy differ fundamentally from polygenic inheritance?

- A) Pleiotropy involves multiple genes affecting one trait, whereas polygenic inheritance involves one gene affecting multiple traits
- B) Pleiotropy involves one gene affecting multiple traits, whereas polygenic inheritance involves multiple genes affecting one trait
- C) Pleiotropy applies only to prokaryotes, whereas polygenic traits apply only to eukaryotes
- D) Neither pleiotropy nor polygenic inheritance involves nuclear genes

Answer Key

- 1. B — Pleiotropy occurs when a single gene locus produces multiple phenotypic effects.
- 2. A — Sickle cell anemia, Cystic fibrosis, and Huntington disease are human examples of pleiotropic gene action.
- 3. B — A single mutant gene in Huntington disease causes both progressive dementia and early death.
- 4. B — Pleiotropy involves one gene influencing multiple traits, whereas polygenic inheritance involves many genes influencing a single trait.

Lesson 24: Environmental effects

1. According to Lesson 24, an organism's phenotype depends on an interaction between which two factors?

- A) Genotype and environment
- B) Diet and exercise only
- C) Temperature and humidity only
- D) Age and gender only

2. In Himalayan rabbits and Siamese cats, what effect occurs when the temperature exceeds 33 °C?

- A) Fur becomes darker due to increased melanin production
- B) The tyrosinase enzyme inactivates, resulting in reduced melanin and lighter fur
- C) The allele mutates permanently into a dominant allele
- D) Melanin converts into blue anthocyanin

3. What specific environmental factor alters the color of Hydrangea flowers?

- A) Atmospheric pressure
- B) Soil pH
- C) Light intensity
- D) Wind speed

4. Which of the following animals is explicitly listed as an example where coat color is affected by temperature?

- A) Polar bear
- B) Arctic fox
- C) Desert lizard
- D) Tropical parrot

Answer Key

1. A — The phenotype of an organism depends on its genotype as well as the environment under which genes act.
2. B — At temperatures above 33 °C, the tyrosinase enzyme inactivates, resulting in reduced melanin production.
3. B — Soil pH impacts phenotype by altering the color of Hydrangea flowers.
4. B — Arctic foxes are listed alongside Himalayan rabbits and Siamese cats as examples of temperature-sensitive coat color.

Lesson 25: What Are Chromosomes?

1. Chromosomes are described as rod-shaped, filamentous bodies located in the:

- A) Cytoplasm
- B) Nucleus
- C) Cell wall
- D) Vacuole

2. Who first described chromosomes in 1875?

- A) Waldeyer
- B) Strausberger
- C) Gregor Mendel
- D) T. H. Morgan

3. The term "chromosome" (meaning colored body) was first used in 1888 by:

- A) Strausberger
- B) Waldeyer
- C) Watson
- D) Crick

4. During which cell division stage can chromosome numbers be counted most easily?

- A) Interphase
- B) Prophase
- C) Mitotic metaphase
- D) Telophase

Answer Key

1. B — Chromosomes are rod-shaped bodies present within the cell nucleus
2. B — Strausberger first described chromosomes in 1875
3. B — Waldeyer coined the term "chromosome" in 1888
4. C — Chromosome numbers can be counted easily during mitotic metaphase

Lesson 26: Chromosome Structure

1. Chromosomes are composed of which two primary constituents?

- A) Carbohydrates and lipids
- B) DNA and associated proteins
- C) RNA and cell wall polysaccharides
- D) Centrosomes and centrioles

2. What is the constricted region of a chromosome where sister chromatids are attached?

- A) Nucleolus
- B) Centromere
- C) Telomere
- D) Kinetochore corona

3. Where are chromosomes or DNA located within a cell?

- A) In the cell membrane
- B) In the nucleus or nucleoid region
- C) Floating freely in the cell wall
- D) Inside cytoplasmic vesicles

4. Which type of protein is specifically mentioned as part of a chromosome's structure alongside DNA?

- A) Tubulin
- B) Histones
- C) Securin
- D) Separase

Answer Key

1. B — Chromosomes are made of DNA and associated proteins.
2. B — The centromere is defined as the constricted region where chromatids are attached.
3. B — DNA or chromosomes are present within the cell's nucleus or nucleoid region.
4. B — A chromosome constitutes a DNA molecule, Histone proteins, and supercoiled DNA.

Lesson 27: Variation in number of chromosomes

1. What is the total chromosome count in human somatic cells?

- A) 23 chromosomes
- B) 46 chromosomes
- C) 48 chromosomes
- D) 78 chromosomes

2. Which organism is noted as having only two chromosomes?

- A) Mosquito
- B) Fruit fly
- C) Round worm
- D) Fungi

3. Which pair of organisms both possess a total chromosome count of 78?

- A) Chimpanzee and Horse

- B) Dog and Chicken
- C) Goldfish and Fruit fly
- D) Mosquito and Round worm

4. Most organisms possess chromosome numbers falling within which general range?

- A) 2 to 6
- B) 12 to 50
- C) 100 to 200
- D) 300 or more

Answer Key

1. B — Humans have 46 total chromosomes (23 pairs: 22 pairs of autosomes and one pair of sex chromosomes).
2. C — Round worms have a total chromosome count of only two.
3. B — Both the Dog and the Chicken have 78 chromosomes.
4. B — Most organisms have chromosome numbers ranging between 12 and 50.

Lesson 28: Centromere-function

1. What are the two essential features of all eukaryotic chromosomes, each absolutely necessary for chromosome stability?

- A) Centrosomes and centrioles
- B) Centromere and telomeres
- C) Chromatin and nucleoplasm
- D) Kinetochore and spindle fibers

2. For which process are centromeres required?

- A) DNA replication during interphase
- B) Segregation of chromosomes during meiosis and mitosis
- C) Protein synthesis at the ribosome
- D) Formation of the nuclear envelope

3. When chromosomes are stained, how does the centromere appear, and what does it join?

- A) A light-stained region where chromatids repel each other
- B) A dark-stained region where the two sister chromatids of a chromosome are joined during mitotic metaphase
- C) A dark-stained region at the terminal ends of the chromosome
- D) A light-stained region that dissolves before anaphase

4. What must happen to the centromere shared by the sister chromatids during mitosis, and what is observed in anaphase?

- A) It must fuse with the telomeres so chromatids stay together
- B) It must divide so chromatids can migrate to opposite poles; centromeres are the first parts of chromosomes seen moving towards opposite poles during anaphase
- C) It remains unchanged while the telomeres lead the movement
- D) It dissolves so the chromatids can separate randomly

Answer Key

1. B — Centromere and telomeres are the two essential features of all eukaryotic chromosomes, each absolutely necessary for chromosome stability.
2. B — Centromeres are required for segregation of chromosomes during meiosis and mitosis; telomeres provide terminal stability and ensure survival.
3. B — When chromosomes are stained, the centromere shows as a dark-stained region where the two sister chromatids of a chromosome are joined during mitotic metaphase.
4. B — During mitosis the centromere shared by the sister chromatids must divide so chromatids can migrate to opposite poles; centromeres are the first parts seen moving toward opposite poles during anaphase, with the remaining regions lagging behind as if pulled by the centromere.

Lesson 28: Chromosomes in Prokaryotes and Eukaryotes

1. Prokaryotic cells typically possess:

- A) Many linear chromosomes made of DNA and histones
- B) A single circular chromosome plus plasmids
- C) Multiple membrane-bound nuclear chromosomes
- D) No DNA molecules

2. Prokaryotic chromosomes are composed of:

- A) DNA and histones
- B) Made only of DNA
- C) RNA and carbohydrates
- D) Lipids and histones

3. Eukaryotic chromosomes are situated in the:

- A) Cytoplasm
- B) Nucleus
- C) Cell membrane
- D) Nucleoid region

4. Unlike prokaryotes, eukaryotic chromosomes are composed of DNA bound to:

- A) Plasmids
- B) Histones
- C) Ribosomes
- D) Peptidoglycan

Answer Key

1. B — Prokaryotes contain a single circular chromosome along with plasmids
2. B — Prokaryotic chromosomes consist only of DNA without histone proteins
3. B — Eukaryotic chromosomes are enclosed within a true membrane-bound nucleus
4. B — Eukaryotic DNA is packaged together with histone proteins

Lesson 29: Salient features of Chromosomes

1. Which of the following statements accurately describes the shape of chromosomes as detailed in Lesson 29?

- A) Chromosomes are always spherical in shape
- B) Chromosomes can be either circular or linear in shape

- C) Chromosomes are always rod-shaped and never circular
- D) Chromosomes lack any defined shape

2. According to Lesson 29, what is the estimated gene count in humans?

- A) 5,800
- B) 13,700
- C) 25,000 - 27,000
- D) 45,000

3. Which organism listed in Lesson 29 possesses the highest total number of genes among the given examples?

- A) Human
- B) Drosophila
- C) E. coli
- D) Rice

4. How does the gene density per Mb of DNA in prokaryotes compare to that in eukaryotes?

- A) Prokaryotes have lower gene density per Mb of DNA than eukaryotes
- B) Prokaryotes exhibit higher gene density per Mb of DNA than eukaryotes
- C) Both have identical gene density per Mb of DNA
- D) Gene density cannot be compared between the two groups

Answer Key

1. B — Lesson 29 establishes that chromosome shapes can be either circular or linear.
2. C — The source handout specifies that humans have an estimated 25,000 to 27,000 genes.
3. D — Rice is listed with 45,000 genes, which exceeds the counts for humans, Drosophila, and E. coli.
4. B — Prokaryotic genomes exhibit higher gene density per Mb of DNA than eukaryotic genomes.

Lesson 30: Size of Chromosomes

1. The length of chromosomes generally varies between:

- A) 0.01 μm to 0.1 μm
- B) 0.5 μm to 32 μm
- C) 50 μm to 100 μm
- D) 1 mm to 5 mm

2. Which plant species possesses the longest known normal chromosomes (32 μm)?

- A) Maize
- B) Trillium
- C) Onion
- D) Pea

3. Giant chromosomes found in Diptera can reach lengths up to:

- A) 32 μm
- B) 100 μm
- C) 300 μm
- D) 1000 μm

4. In general, how do plant chromosome sizes compare to animal chromosome sizes?

- A) Plants have shorter chromosomes than animals
- B) Plants have longer chromosomes than animals
- C) Plant and animal chromosomes are identical in size
- D) Plants lack chromosomes entirely

Answer Key

1. B — Chromosome length ranges from 0.5 μm to 32 μm
2. B — The longest normal chromosomes are found in Trillium (32 μm)
3. C — Giant polytene chromosomes in Diptera can measure up to 300 μm in length
4. B — Generally, plant species possess longer chromosomes than animal species

Lesson 31: Types of Chromosomes

1. Which two primary types of eukaryotic chromosomes are described in Lesson 31?

- A) Homologous chromosomes and heterologous chromosomes
- B) Autosomes and sex chromosomes
- C) Linear chromosomes and circular plasmids
- D) Metacentric chromosomes and telocentric chromosomes

2. How are autosomes defined in genetics?

- A) Chromosomes that differ between males and females
- B) Paired chromosomes with the same length, shape, centromere location, and genes (any chromosomes other than sex chromosomes)
- C) Unpaired chromosomes found only in gametes
- D) Chromosomes composed entirely of heterochromatin

3. Sex chromosomes are defined as members of a pair of chromosomes that:

- A) Are identical in length and shape in both sexes
- B) Differ between males and females
- C) Carry no genes at all
- D) Exist only in prokaryotes

4. Which of the following statements is true regarding autosomes?

- A) They are non-paired single chromosomes
- B) They include all chromosomes other than the sex chromosomes
- C) They determine the sex of the offspring
- D) They are found only in plants

Answer Key

1. B — Eukaryotic chromosomes are classified into two main types: autosomes and sex chromosomes.
2. B — Autosomes are paired chromosomes with matching length, shape, centromere location, and genes (all chromosomes except sex chromosomes).
3. B — Sex chromosomes are defined as members of a chromosome pair that differ between males and females.
4. B — Any chromosome that is not a sex chromosome is categorized as an autosome.

Lesson 32: What Are Telomeres?

1. What are the two physical ends of a chromosome called?

- A) Centromeres
- B) Kinetochores
- C) Telomeres
- D) Chromatids

2. Telomeres are required for chromosomal:

- A) Cleavage and degradation
- B) Replication and stability
- C) Nondisjunction
- D) Spindle attachment

3. The handout compares telomeres to which everyday object?

- A) Zipper teeth
- B) Ends of shoestrings
- C) Buttons
- D) Paper clips

4. How do normal chromosome ends compare to broken chromosome ends regarding stickiness?

- A) Normal ends are sticky, broken ends are not
- B) Broken ends are sticky, whereas normal ends are not sticky
- C) Both normal and broken ends are highly sticky
- D) Neither normal nor broken ends display stickiness

Answer Key

- 1. C — The terminal structural ends of chromosomes are called telomeres
- 2. B — Telomeres are essential for preserving chromosome stability and enabling replication
- 3. B — Telomeres function analogously to the protective plastic tips at the ends of shoestrings
- 4. B — Broken chromosome ends readily fuse because they are sticky, whereas normal intact telomeres are non-sticky

Lesson 33: Euchromatin and Heterochromatin

1. What term is used for the darker-staining chromosomal regions generally found near the centromere?

- A) Euchromatin
- B) Heterochromatin
- C) Constitutive chromatin
- D) Facultative chromatin

2. What are the light-staining chromosomal regions called?

- A) Heterochromatin
- B) Euchromatin
- C) Constitutive heterochromatin
- D) Facultative heterochromatin

3. Heterochromatin is categorized into which two distinct groups?

- A) Primary and secondary heterochromatin

- B) Active and inactive heterochromatin
- C) Constitutive heterochromatin and facultative heterochromatin
- D) Dense and loose heterochromatin

4. Which type of heterochromatin remains permanently in the heterochromatic state and does not revert?

- A) Facultative heterochromatin
- B) Constitutive heterochromatin
- C) Euchromatin
- D) Centromeric chromatin

Answer Key

1. B — Dark-staining chromosomal regions near centromeres or telomeres represent heterochromatin.
2. B — Light-staining chromosomal regions are designated as euchromatin.
3. C — Heterochromatin is classified into constitutive heterochromatin and facultative heterochromatin.
4. B — Constitutive heterochromatin remains permanently in the heterochromatic stage without reverting to euchromatin.

Lesson 34: Staining and banding chromosomes

1. In Feulgen staining, what treatment produces a free aldehyde group in deoxyribose molecules?

- A) Mild hydrolysis in 1N HCl at 60 C for 10 minutes
- B) Staining with quinacrine mustard under UV light
- C) Digestion with restriction endonucleases
- D) Heating in 5N NaOH at 100 C for 1 hour

2. What chemical reagent is used in Q-banding to reveal fluorescent bands under UV light?

- A) Giemsa stain
- B) Quinacrine mustard
- C) Feulgen reagent
- D) Ethidium bromide

3. What chromosomal regions do the bands produced by C-banding correspond to?

- A) Euchromatin regions
- B) Constitutive heterochromatin
- C) Single nucleotide polymorphisms
- D) Histone H1 protein tails

4. How are R-bands (reverse bands) characterized relative to Q-bands?

- A) They stain distal ends exclusively
- B) They are located in zones between Q-bands that do not fluoresce with quinacrine mustard
- C) They only stain centromeric AT-rich regions
- D) They destroy the sugar-phosphate backbone

Answer Key

1. A — Feulgen staining subjects cells to mild hydrolysis in 1N HCl at 60 C for 10 minutes, freeing aldehyde groups in deoxyribose.
2. B — Q-banding utilizes quinacrine mustard to produce fluorescent bands when observed under UV light.

3. B — C-banding specifically stains constitutive heterochromatin, primarily located around centromeres.
4. B — R-bands represent reverse banding situated in non-fluorescing zones between Q-bands.

Lesson 35: Nucleosomes

1. What length of DNA wraps around the octamer core of a single nucleosome?
 - A) 20 bp
 - B) 60 bp
 - C) 147 bp
 - D) 300 bp
2. What is the length range of linker DNA located between adjacent nucleosomes?
 - A) 5–10 bp
 - B) 20–60 bp
 - C) 147–200 bp
 - D) 500–1000 bp
3. Which five abundant histone types exist in eukaryotic chromatin?
 - A) H1, H2A, H2B, H3, and H4
 - B) CDE-I, CDE-II, CDE-III, CEN, and CBF
 - C) A, B, C, D, and E
 - D) DNA, RNA, ATP, GTP, and CTP
4. When are nucleosomes assembled during the cell cycle?
 - A) Exclusively during mitotic metaphase
 - B) Immediately after DNA replication
 - C) During cytokinesis in plant cells
 - D) Prior to fertilization in mature gametes

Answer Key

1. C — A nucleosome consists of a core of eight histone proteins with 147 bp of DNA wrapped around them.
2. B — Linker DNA situated between nucleosomes spans 20 to 60 base pairs in length.
3. A — The five abundant histone classes in chromatin are H1, H2A, H2B, H3, and H4.
4. B — Nucleosome assembly takes place immediately following DNA replication.

Lesson 36: Chromatin and Chromosomes

1. The complexes formed between eukaryotic DNA and proteins are termed:
 - A) Plasmids
 - B) Chromatin
 - C) Kinetochores
 - D) Centrioles
2. What are the major proteins associated with eukaryotic chromatin?
 - A) Polymerases
 - B) Histones
 - C) Ligases
 - D) Endonucleases

3. Histone proteins contain a high proportion of which basic amino acids?

- A) Alanine and Glycine
- B) Arginine and Lysine
- C) Tryptophan and Tyrosine
- D) Glutamate and Aspartate

4. How many abundant histone types exist in eukaryotic chromatin (H1, H2A, H2B, H3, H4)?

- A) Three
- B) Five
- C) Ten
- D) Twelve

Answer Key

1. B — Chromatin is the eukaryotic complex composed of DNA and associated proteins
2. B — Histones are the primary protein components bound to eukaryotic DNA
3. B — Histones are basic proteins rich in lysine and arginine residues
4. B — Eukaryotic chromatin contains five main histone classes: H1, H2A, H2B, H3, and H4

Lesson 38: Higher order chromatin folding**1. What is the structural diameter of the basic "beads-on-a-string" nucleosome form of chromatin?**

- A) 2 nm
- B) 11 nm
- C) 30 nm
- D) 300 nm

2. When nucleosomes pack closely together, they form a coiling chromatin fiber with what diameter?

- A) 11 nm
- B) 20 nm
- C) 30 nm
- D) 700 nm

3. In extended chromosome structure, loops of the 30-nm chromatin fiber attach to a central chromosome scaffold, producing a fiber of what diameter?

- A) 11 nm
- B) 30 nm
- C) 300 nm
- D) 1400 nm

4. What is the total diameter of an entire mitotic metaphase chromosome in its fully condensed state?

- A) 300 nm
- B) 300 nm (scaffold-associated fiber)
- C) 700 nm
- D) 1400 nm

Answer Key

1. B — The "beads-on-a-string" nucleosome arrangement measures 11 nm in diameter.
2. C — Packed nucleosomes assemble into a 30-nm chromatin fiber.
3. C — Loops of 30-nm chromatin fiber associate with a chromosome scaffold to form a 300-nm fiber.
4. D — A fully condensed metaphase chromosome spans a total diameter of 1400 nm across its two chromatids.

Lesson 39: Kinetochore

1. What is the actual site within the centromere region where spindle fibers attach during cell division?

- A) The telomere
- B) The kinetochore
- C) The nucleolus organizer
- D) The secondary constriction

2. How long is the CEN DNA sequence found within kinetochore regions?

- A) 50 base pairs
- B) 120 base pairs
- C) 300 base pairs
- D) 1000 base pairs

3. The CEN DNA sequence consists of which three sub-domains?

- A) CDE-I, CDE-II, and CDE-III
- B) H1, H2A, and H2B
- C) CEN-A, CEN-B, and CEN-C
- D) TATA, CAAT, and GC boxes

4. A point mutation in which CEN DNA sub-domain completely eliminates the centromere's ability to function?

- A) CDE-I
- B) CDE-II
- C) CDE-III
- D) CDE-IV

Answer Key

1. B — The actual attachment location for spindle fibers within the centromere is called the kinetochore.
2. B — CEN DNA located within kinetochore regions is 120 base pairs long.
3. A — CEN DNA comprises three sub-domains designated CDE-I, CDE-II, and CDE-III.
4. C — A point mutation specifically in CDE-III completely abolishes centromere function.

Lesson 40: Karyotyping

1. Karyotype is defined as the general morphology of which type of chromosomes?

- A) Gametic chromosomes
- B) Somatic chromosomes
- C) Bacterial chromosomes
- D) Viral chromosomes

2. What is a diagrammatic representation of a species' karyotype showing all its morphological features called?

- A) An idiotype (idiogram)
- B) A phenotype
- C) A genotype
- D) A haplotype

3. Which stain is explicitly mentioned in Lesson 40 for producing banding patterns to identify chromosomes?

- A) Feulgen stain
- B) Quinacrine mustard
- C) Giemsa stain
- D) Ethidium bromide

4. In what order are chromosomes arranged when constructing a standard karyotype display?

- A) Ascending order of size
- B) Descending order of size with centromeres aligned
- C) Alphabetical order of gene names
- D) Random order

Answer Key

1. B — Karyotype refers to the general morphological arrangement of somatic chromosomes.
2. A — An idiotype is a diagrammatic representation of a species' karyotype detailing morphological features.
3. C — Giemsa staining is specified in Lesson 40 for generating characteristic chromosomal banding patterns.
4. B — Karyotypes arrange chromosomes in descending order of size with their centromeres aligned.

Lesson 41: Prokaryotic Chromosomes

1. According to Lesson 41, which group has the smallest genome length?

- A) Eukaryotic yeast
- B) RNA viruses
- C) Bacteria
- D) Drosophila

2. Larger bacteriophage genomes may contain up to how many genes?

- A) 10 genes
- B) 150 genes
- C) 3,000 genes
- D) 45,000 genes

3. Approximately how many genes are organized into the single circular chromosome of E. coli?

- A) 100 genes
- B) 1,500 genes
- C) 3,000 to 4,000 genes
- D) 25,000 genes

4. How many primary chromosomes do prokaryotic organisms usually possess?

- A) Only one circular chromosome (plus plasmids)

- B) Two linear chromosomes
- C) 46 chromosomes
- D) No chromosomes at all

Answer Key

1. B — Genome length is smallest among RNA viruses.
2. B — Larger bacteriophage genomes can contain up to 150 genes.
3. C — E. coli contains roughly 3,000 to 4,000 genes organized on its single circular chromosome.
4. A — Prokaryotes typically possess a single circular chromosome plus plasmids.

Lesson 42: Sex chromosomes and sex determination

1. In human gender determination, which sex is heterogametic?

- A) Female (XX)
- B) Male (XY)
- C) Female (ZW)
- D) Male (XO)

2. In birds, amphibians, reptiles, and butterflies, which combination represents the female heterogametic sex?

- A) XX female
- B) XY female
- C) ZW female
- D) XO female

3. In the XO sex determination system found in grasshoppers, what is the male sex chromosome constitution?

- A) XX
- B) XO (a single X chromosome, no Y)
- C) XY
- D) ZW

4. What is the probability of an offspring being male or female in human gender inheritance?

- A) 25% male, 75% female
- B) 50% male, 50% female
- C) 75% male, 25% female
- D) 100% determined by temperature

Answer Key

1. B — Human males are heterogametic (XY), producing two types of sex cells (X or Y).
2. C — In birds, reptiles, and butterflies, females are heterogametic with ZW sex chromosomes.
3. B — Male grasshoppers possess a single X chromosome without a Y, represented as XO.
4. B — Human fertilization results in a 50% chance of a girl (XX) and a 50% chance of a boy (XY).

Lesson 43: Salient features of DNA

1. According to the base pairing rules in Lesson 43, Adenine specifically forms complementary pairs with:

- A) Guanine
- B) Cytosine
- C) Thymine
- D) Uracil

2. What does Chargaff's G-C rule state regarding the (G+C) percentage in DNA?

- A) It is always 50% across all species
- B) It remains constant within a given species but varies between different species
- C) It equals the (A+T) percentage in all organisms
- D) It is always higher than (A+T)

3. What does Chargaff's "Cluster rule" specify regarding pyrimidines?

- A) Pyrimidines never occur near purines
- B) Pyrimidines cluster together on one strand, dictating purine clustering on the complementary strand
- C) Pyrimidines pair only with other pyrimidines
- D) Pyrimidines are absent from coding strands

4. What chemical components form the structural outer backbones of the DNA double helix?

- A) Amino acids and peptide bonds
- B) Nitrogenous bases and hydrogen bonds
- C) Alternating sugar and phosphate groups
- D) Fatty acids and glycerol

Answer Key

1. C — Purines pair with pyrimidines, specifically Adenine (A) with Thymine (T).
2. B — The (G+C)% remains constant within a given species but varies between different species.
3. B — The cluster rule notes pyrimidines cluster together on one strand, dictating purine clustering on the opposite strand.
4. C — Alternating sugar and phosphate groups form the double-stranded helical backbone.

Lesson 44: Nucleotide

1. A complete nucleotide monomer is composed of how many structural components?

- A) One
- B) Two
- C) Three
- D) Four

2. What are the three components of a nucleotide?

- A) Pentose sugar, nitrogenous base, and phosphate group
- B) Hexose sugar, amino acid, and lipid
- C) Glycerol, fatty acid, and nitrogenous base
- D) Histone, DNA, and centromere

3. A nucleic acid polymer containing fewer than 10 monomer units is termed an:

- A) Polynucleotide
- B) Oligonucleotide
- C) Nucleoside
- D) Octamer

4. A nucleoside consists of:

- A) Sugar + base + phosphate
- B) Monomer of sugar + base
- C) Base + phosphate group only
- D) Histone protein + DNA

Answer Key

1. C — Every nucleotide is constructed from three distinct sub-units
2. A — Nucleotides comprise a 5-carbon pentose sugar, a nitrogenous base, and a phosphate group
3. B — Short nucleic acid polymers under 10 units long are classified as oligonucleotides
4. B — A nucleoside is a two-part molecule formed solely by a sugar and a nitrogenous base

Lesson 45: Conformations of DNA**1. Which DNA conformation is a left-handed helix?**

- A) A-DNA
- B) B-DNA
- C) Z-DNA
- D) C-DNA

2. Which DNA conformation is the widest helix and features a 6Å central hole along its axis?

- A) A-DNA
- B) B-DNA
- C) Z-DNA
- D) Triple-stranded DNA

3. Standard B-DNA is characterized as a:

- A) Left-handed narrowest helix
- B) Right-handed helix with intermediate width
- C) Single-stranded left-handed helix
- D) Helix lacking a minor groove

4. Which DNA conformation completely lacks a major groove?

- A) A-DNA
- B) B-DNA
- C) Z-DNA
- D) All helical forms possess a major groove

Answer Key

1. C — Z-DNA is unique among common DNA conformations for having a left-handed helical structure
2. A — A-DNA represents the widest right-handed DNA conformation with an internal central hole
3. B — B-DNA exhibits a right-handed double-helical structure of intermediate width
4. C — Z-DNA is the narrowest form and lacks a major groove

Lesson 46: Chargaff Rule of Base Ratios**1. Chargaff's rules state that in double-stranded DNA, purines pair with:**

- A) Other purines

- B) Pyrimidines
- C) Amino acids
- D) Phosphate backbones

2. According to Chargaff's parity rules, the percentage of Adenine (A) approximately equals:

- A) Percentage of Guanine (G)
- B) Percentage of Cytosine (C)
- C) Percentage of Thymine (T)
- D) Percentage of Uracil (U)

3. Chargaff's G-C rule states that the (G+C)% is:

- A) Variable within individual tissue cells of a single organism
- B) Constant within a species, but often differs between species
- C) Exactly 50% in all living organisms
- D) Zero in prokaryotic DNA

4. The cluster rule states that pyrimidines often occur in clusters, meaning the complementary strand contains clusters of:

- A) Pyrimidines
- B) Purines
- C) Histones
- D) Ribose sugars

Answer Key

1. B — Chargaff established that purines specifically pair with pyrimidines
2. C — Parity rules dictate that % A equals % T and % G equals % C
3. B — The overall (G+C) percentage remains constant within a single species but varies across different species
4. B — Pyrimidine clusters on one DNA strand dictate complementary purine clusters on the opposing strand

Lesson 47: Giant or Polytene chromosome

1. Giant chromosomes were first discovered in 1881 in dipteran salivary glands by which scientist?

- A) Flemming
- B) Balbiani
- C) Mendel
- D) Morgan

2. What is the total length of giant chromosomes in *Drosophila melanogaster*?

- A) 32 μ
- B) 300 μ
- C) 2,000 μ
- D) 1400 nm

3. Approximately how many dark-staining bands can be recognized in the giant chromosomes of *Drosophila*?

- A) 500
- B) 1,000

- C) 5,000
- D) 45,000

4. How many individual strand copies are estimated to form the polytene giant chromosomes of Chironomus?

- A) 64
- B) 512
- C) 4,096
- D) 10,000

Answer Key

1. B — Balbiani first discovered giant chromosomes in 1881 within dipteran salivary glands.
2. C — The total combined length of giant chromosomes in *D. melanogaster* is approximately 2,000 μ .
3. C — *Drosophila* giant chromosomes display about 5,000 distinct dark-staining bands under the microscope.
4. C — Repeated DNA replication without cell division yields about 4,096 strands in *Chironomus* polytene chromosomes.

Lesson 48: Lampbrush Chromosome

1. Lampbrush chromosomes were first observed in 1882 by:

- A) Balbiani
- B) Flemming
- C) Mendel
- D) Morgan

2. In vertebrates, lampbrush chromosomes are predominantly found in which cell type?

- A) Somatic liver cells
- B) Oocytic nuclei
- C) Red blood cells
- D) Muscle fibers

3. What does a single lateral loop in a lampbrush chromosome structurally represent?

- A) One centromere
- B) One chromatid (a single DNA molecule)
- C) One ribosome
- D) One nucleolus

4. How are the two ends of each lateral loop in a lampbrush chromosome structurally distinguished?

- A) One thin end and one thick end (asymmetry)
- B) Both ends are identical
- C) Both ends have centromeres
- D) One end is always broken

Answer Key

1. B — Flemming first observed lampbrush chromosomes in the year 1882.
2. B — Lampbrush chromosomes occur prominently in the oocytic nuclei of vertebrates.
3. B — Each extended loop corresponds to one chromatid, representing a single DNA molecule.

4. A — Each lampbrush loop displays asymmetry, having one thin end and one thick end.

Lesson 49: Cell division and reproduction

1. According to Lesson 49, what essential event must occur before a cell's nucleus divides?

- A) Cytoplasm is completely pinched off
- B) Genetic material replicates
- C) Cell wall dissolves entirely
- D) Spindle fibers break down and dissolve

2. Cell division and reproduction are described as two fundamental processes occurring in:

- A) Eukaryotes only
- B) Prokaryotes only
- C) Both prokaryotes and eukaryotes
- D) Viruses only

3. Why is cell division required in living organisms according to Lesson 49?

- A) For cell growth, development, and reproduction
- B) To eliminate all histone proteins
- C) To convert DNA into single-stranded RNA
- D) To prevent DNA replication during interphase

4. What happens to a cell's genetic material prior to nuclear division?

- A) It is degraded by endonucleases
- B) It replicates so genetic material can be distributed
- C) It converts permanently into heterochromatin
- D) It translocates to non-homologous chromosomes

Answer Key

- 1. B — Before the nucleus divides, the cell's genetic material replicates.
- 2. C — Cell division and reproduction represent fundamental processes in both prokaryotes and eukaryotes.
- 3. A — Cell division is a fundamental process required for organismal growth, development, and reproduction.
- 4. B — Genetic material replicates prior to nuclear division to enable proper distribution to daughter cells.

Lesson 50: Cell Cycle

1. The eukaryotic cell cycle is divided into two primary phases:

- A) Prophase and Anaphase
- B) Interphase and Mitotic phase
- C) G1 phase and G2 phase
- D) Meiosis I and Meiosis II

2. The G0 phase of the cell cycle is also referred to as the:

- A) Synthesis phase
- B) Post-mitotic phase
- C) Pre-mitotic growth phase
- D) Cytokinesis phase

3. Eukaryotic cells generally enter the non-dividing G0 state from which cell cycle stage?

- A) G1 phase
- B) S phase
- C) G2 phase
- D) Metaphase

4. Interphase proceeds in three sequential stages in what order?

- A) S → G1 → G2
- B) G1 → S → G2
- C) G2 → S → G1
- D) G1 → G2 → S

Answer Key

1. B — The cell cycle consists of Interphase and the Mitotic (M) phase
2. B — G0 is a quiescent post-mitotic state
3. A — Cells exit the G1 phase to enter the G0 post-mitotic state
4. B — Interphase progresses through G1, S (synthesis), and G2 stages

Lesson 51: Mitosis

1. What is the result of mitotic cell division from a single parent cell?

- A) Four genetically distinct haploid gametes
- B) Two genetically identical daughter cells
- C) Eight non-identical daughter nuclei
- D) A single tetraploid zygote

2. Which cell types can undergo mitosis according to Lesson 51?

- A) Diploid cells only
- B) Haploid cells only
- C) Both haploid and diploid cells
- D) Bacterial plasmids only

3. What essential components must each newly formed daughter cell receive following mitotic division?

- A) Its own cytoplasm and DNA
- B) Plasmids and a cell wall
- C) Four centriole pairs and a cleavage furrow
- D) A haploid set of 23 chromosomes only

4. Before sister chromatids break apart during mitosis, what process must occur to produce identical copies of the DNA?

- A) Recombination during synapsis
- B) DNA replication prior to mitosis
- C) Histone acetylation in G0
- D) Cytokinesis via cell plate

Answer Key

1. B — Mitosis divides a single cell into two new daughter cells that are genetically identical to the original cell.
2. C — Mitotic cell division occurs in both haploid and diploid cells.

3. A — Each daughter cell produced by mitosis must receive its own cytoplasm and DNA.
4. B — Chromosomes replicate before mitosis to produce identical copies of DNA before sister chromatids separate.

Lesson 52: Un-replicated and replicated chromosome

1. Duplicated chromosomes that remain attached at their centromere are known as:

- A) Homologous pairs
- B) Sister chromatids
- C) Tetrads
- D) Polynucleotides

2. When does chromosome duplication (replication) take place during the cell cycle?

- A) During metaphase alignment
- B) Before cell division takes place
- C) During cytokinesis
- D) After telophase is complete

3. As long as duplicated chromosomes remain attached at their centromere, they are designated as:

- A) Two independent chromosomes
- B) Sister chromatids
- C) Non-homologous translocations
- D) Acquired variations

4. What causes chromosomes to transition from an un-replicated state to a replicated state?

- A) Splitting of centromeres in anaphase
- B) Replication of the entire DNA of each chromosome
- C) Dissolution of the nuclear membrane
- D) Attachment of asters to centrioles

Answer Key

1. B — Duplicated chromosomes attached at their centromere are referred to as sister chromatids as long as they remain attached.
2. B — Chromosomes (DNA) replicate before cell division takes place.
3. B — Attached duplicated chromosome copies are defined as sister chromatids.
4. B — Duplicated chromosomes arise through DNA replication, duplicating the entire DNA of each chromosome.

Lesson 53: Interphase

1. What proportion of total cell cycle time is spent in Interphase?

- A) ~10%
- B) ~50%
- C) ~90%
- D) 100%

2. What key molecular process occurs specifically during the S phase of Interphase?

- A) Absorption of nutrients and initial growth

- B) Synthesis of new DNA for daughter cells
- C) Splitting of centromeres
- D) Pinching of the cleavage furrow

3. What takes place during the G1 phase of Interphase?

- A) The new cell absorbs nutrients and grows larger
- B) Homologous chromosomes form tetrads
- C) Nuclear membranes break down
- D) Chromosomes line up at the equator

4. Why does a cell in the G2 phase eventually need to enter cell division?

- A) Because it loses all its DNA
- B) Because it continues to grow and gets too large, needing to divide
- C) Because centrioles fail to replicate
- D) Because histone proteins are degraded

Answer Key

1. C — Interphase accounts for approximately 90% of total cell cycle duration.
2. B — Genomic DNA synthesis for daughter cells occurs exclusively during the S phase of Interphase.
3. A — In G1 phase, the newly formed cell absorbs nutrients and expands in size.
4. B — During G2, continued growth causes the cell to get too large, triggering the need for cell division.

Lesson 54: Prophase of mitosis

1. Prophase is defined as which stage of mitotic cell division?

- A) The final stage
- B) The middle alignment stage
- C) The initial stage
- D) The resting stage

2. What happens to the nuclear membrane during prophase of mitosis?

- A) It doubles in size
- B) It disappears
- C) It becomes rigid
- D) It divides into four

3. Where do centrioles position themselves during prophase?

- A) At the metaphase plate
- B) On opposite poles of the cell
- C) Inside the nucleus
- D) At the cell membrane

4. At the start of mitosis, DNA wraps around which proteins to condense into visible chromosomes?

- A) Tubulin
- B) Actin
- C) Histone proteins
- D) Keratin

Answer Key

1. C — Prophase is the initial stage of mitotic cell division.
2. B — Disappearance of the nuclear membrane is one of four major events characterizing prophase.
3. B — Centrioles separate and move to opposite poles of the cell during prophase.
4. C — Chromatin condenses at mitosis onset as DNA wraps tightly around histone proteins.

Lesson 55: Metaphase

1. What key positioning event occurs to chromosomes during metaphase?

- A) They split into single chromatids
- B) They line up along the middle alignment (metaphase) plate
- C) They dissolve into chromatin
- D) They migrate to the poles

2. What structures attach the centromere of each sister chromatid to the centrioles at the poles?

- A) Cleavage furrows
- B) Spindle fibers
- C) Nuclear pores
- D) Ribosomes

3. How many major events characterize metaphase according to Lesson 55?

- A) One event
- B) Two events
- C) Three events
- D) Four events

4. Where are centrioles positioned during metaphase?

- A) At the cell center alongside chromosomes
- B) At the cell poles anchoring the spindle apparatus
- C) Inside the nucleus
- D) At the cleavage furrow

Answer Key

1. B — Metaphase is defined by chromosomes lining up along the middle alignment plate of the cell.
2. B — Spindle fibers connect centromeres to centrioles stationed at opposite cell poles.
3. B — Lesson 55 outlines two key metaphase events: alignment in the middle and spindle attachment.
4. B — Centrioles reside at the cell poles to anchor the spindle apparatus.

Lesson 56: Anaphase of mitosis

1. What event happens to centromeres during anaphase of mitosis?

- A) They duplicate
- B) They split, separating sister chromatids
- C) They dissolve
- D) They fuse

2. Once sister chromatids separate during anaphase, they are designated as:

- A) Homologous tetrads
- B) Individual daughter chromosomes
- C) Bivalents
- D) Chromatins

3. Destruction of which inhibitory protein permits separase protease to split centromeres during anaphase?

- A) Cyclin
- B) Securin
- C) Tubulin
- D) Actin

4. Which spindle fibers push against each other to stretch the dividing cell into an oval shape?

- A) Kinetochore fibers
- B) Non-kinetochore (polar) spindle fibers
- C) Astral fibers only
- D) Intermediate filaments

Answer Key

1. B — Anaphase begins with centromere splitting, which separates sister chromatids.
2. B — Upon separation, former sister chromatids become individual daughter chromosomes.
3. B — Securin inhibits separase; its destruction allows separase to cleave centromeres.
4. B — Non-kinetochore spindle fibers elongate and push against one another to stretch the cell.

Lesson 57: Telophase

1. What occurs to spindle fibers during telophase?

- A) They condense into chromatin
- B) They break down and disappear
- C) They duplicate
- D) They form the cell plate

2. Telophase effectively reverses the cellular changes brought about by which phase(s)?

- A) S phase and G2 phase
- B) Prophase (and prometaphase)
- C) Metaphase only
- D) Cytokinesis

3. What nuclear structures re-emerge inside each newly forming daughter nucleus during telophase?

- A) Centrioles
- B) Nucleoli (with nuclear membrane reformation)
- C) Spindle fibers
- D) Chromosomes condense further

4. How do chromosomes change structurally during telophase?

- A) They hyper-condense into 1,400 nm fibers
- B) They uncoil and return to a relaxed chromatin state
- C) They split into chromatids

D) They attach to the cell membrane

Answer Key

1. B — Spindle fibers dissolve and disappear during telophase.
2. B — Telophase reverses the nuclear breakdown and condensation events of prophase.
3. B — Reformation of nuclear membranes is accompanied by the reappearance of nucleoli.
4. B — Condensed chromosomes uncoil and return to a relaxed chromatin state.

Lesson 58: Cytokinesis

1. What is the fundamental definition of cytokinesis?

- A) Division of the cell nucleus
- B) Division of the cytoplasm into two daughter cells
- C) Duplication of chromosomes
- D) Formation of the spindle apparatus

2. In animal cells, cytokinesis occurs through the formation of what structure that pinches the cell?

- A) Cell plate
- B) Contractile cleavage furrow
- C) Phragmoplast
- D) Nuclear envelope

3. In plant cells, cytokinesis is accomplished via the formation of what midway structure?

- A) Cleavage furrow
- B) Cell plate (developing into a cell wall)
- C) Contractile ring
- D) Spindle equator

4. Into how many parts does cytokinesis divide a single parent cell?

- A) Two equal parts
- B) Four haploid parts
- C) Three unequal parts
- D) Eight parts

Answer Key

1. B — Cytokinesis is specifically the division of the cytoplasm into two daughter cells.
2. B — Animal cells divide by forming a contractile cleavage furrow that pinches the cell in two.
3. B — Plant cells construct a cell plate midway between nuclei, which develops into a cell wall.
4. A — Cytokinesis splits the parent cell cytoplasm into two equal daughter cells.

Lesson 59: Mitotic apparatus/ spindles

1. What three structural components comprise the mitotic apparatus/spindles?

- A) Centromeres, telomeres, and kinetochores
- B) Centrosomes, spindle microtubules, and asters
- C) Histones, nucleosomes, and chromatin
- D) Exons, introns, and promoters

2. What are asters in the mitotic spindle apparatus?

- A) Radial arrays of short microtubules extending from each centrosome
- B) Protein structures within the CDE-III domain of CEN DNA
- C) Contractile rings that form cleavage furrows in animal cells
- D) Double-stranded RNA molecules that silence genes

3. Where does the assembly of spindle microtubules begin within the cell?

- A) Inside the nucleolus
- B) At the centrosome
- C) On the cell plate
- D) Within the telomere cap

4. During mitosis, spindle microtubules attach to which specific chromosomal structure to move chromosomes to the metaphase plate?

- A) Kinetochores
- B) Histone tails
- C) Nuclear envelope pores
- D) Poly-A tails

Answer Key

- 1. B — The mitotic apparatus includes centrosomes, spindle microtubules, and asters.
- 2. A — Asters are radial arrays of short microtubules extending outward from each centrosome.
- 3. B — Spindle microtubule assembly initiates within the centrosome.
- 4. A — Spindle microtubules attach specifically to kinetochores to align chromosomes along the metaphase plate.

Lesson 60: Diploid cells and haploid cells**1. What symbol is used to represent haploid sex cells (gametes)?**

- A) $2n$
- B) n
- C) $3n$
- D) $4n$

2. Diploid cells ($2n$) are also known as:

- A) Gametes
- B) Somatic cells (body cells)
- C) Polar bodies
- D) Plasmids

3. Fertilization is defined as the fusion of spermatozoa and ova to produce a diploid cell called a:

- A) Gamete
- B) Zygote
- C) Tetrad
- D) Barr body

4. What is the primary function of meiosis regarding chromosome number?

- A) It doubles the diploid chromosome number to tetraploid

- B) It reduces the diploid chromosome number into a haploid number
- C) It preserves identical diploid chromosome sets in body cells
- D) It converts linear chromosomes into circular plasmids

Answer Key

1. B — Haploid sex cells (gametes) are represented by the symbol n .
2. B — Diploid cells carrying two chromosomes at each location ($2n$) are somatic/body cells.
3. B — Fertilization is the fusion of haploid spermatozoa and ova to form a diploid zygote ($2n=46$).
4. B — Meiosis is the cellular process that reduces diploid chromosome numbers to haploid sets.

Lesson 61: Homologous chromosomes

1. Homologous chromosomes are defined as chromosome pairs that share similar:

- A) Sequence length only
- B) Shape and overall genetic content
- C) Centromere number only
- D) Telomere caps only

2. Where are gene alleles that control the same inherited traits situated on homologous chromosomes?

- A) At the same locus
- B) At random loci
- C) Only on the p arm
- D) Outside the centromere

3. In a diploid organism, where do the two homologous chromosomes originate?

- A) Both from the female parent
- B) Both from the male parent
- C) One from each parent
- D) From sister chromatids

4. How many autosome pairs are present in human somatic cells?

- A) 22 pairs
- B) 23 pairs
- C) 46 pairs
- D) 1 pair

Answer Key

1. B — Homologous chromosomes are paired chromosomes that share similar shape and overall genetic content.
2. A — Corresponding gene alleles reside at the exact same locus on homologous chromosomes.
3. C — Diploid individuals inherit one homologous chromosome from the maternal parent and one from the paternal parent.
4. A — Human somatic cells contain 22 pairs of autosomes and 1 pair of sex chromosomes.

Lesson 62: Meiosis

1. Meiosis reduces the chromosome number of a diploid cell to form:

- A) Tetraploid somatic cells
- B) Haploid gametes
- C) Diploid zygotes directly
- D) Triploid endosperm

2. Why is Meiosis I designated as a reductional division?

- A) It reduces cytoplasm volume by half
- B) Homologous pairs separate, reducing chromosome number from diploid to haploid
- C) It eliminates all sex chromosomes
- D) It doubles the DNA content

3. Human gametes (sperm and egg) formed through meiotic division contain how many total chromosomes?

- A) 46
- B) 23
- C) 92
- D) 12

4. Fusion of haploid spermatozoa and ova during fertilization yields a diploid cell called a:

- A) Gamete
- B) Zygote
- C) Spore
- D) Tetrad

Answer Key

- 1. B — Meiosis reduces diploid chromosome sets to yield haploid gametes.
- 2. B — Homologous pairs separate during Meiosis I, reducing chromosome numbers from diploid to haploid.
- 3. B — Meiotically produced human gametes carry a haploid set of 23 chromosomes.
- 4. B — Sperm and egg fusion restores diploidy, forming a single-celled zygote.

Lesson 63: Stages of meiosis

1. What major event occurs during Meiosis I?

- A) Sister chromatids separate
- B) Homologous chromosomes separate
- C) DNA replication
- D) Cytokinesis only

2. Why is Meiosis I categorized as a reductional division?

- A) It reduces the cytoplasm volume by 90%
- B) It reduces the diploid chromosome number to haploid
- C) It reduces protein synthesis
- D) It reduces the nuclear membrane

3. What structures separate from each other during Meiosis II?

- A) Homologous chromosome pairs
- B) Sister chromatids
- C) Tetrads
- D) Bivalents

4. Why is Meiosis II referred to as an equational division?

- A) It produces four haploid cells with no reduction in chromosome number (sister chromatids separate)
- B) It doubles the chromosome number
- C) It equates cytoplasm between cells
- D) It replicates DNA again

Answer Key

1. B — Homologous chromosomes separate from one another during Meiosis I.
2. B — Meiosis I is termed reductional division because it reduces the diploid chromosome number to haploid.
3. B — Sister chromatids separate during Meiosis II.
4. A — Meiosis II is an equational division because sister chromatids separate without reducing chromosome number.

Lesson 64: Prophase I of meiosis**1. What percentage of the total time required for meiosis is occupied by Prophase I?**

- A) ~10%
- B) ~50%
- C) Over 90%
- D) ~25%

2. How many sub-stages comprise Prophase I of meiosis?

- A) Two sub-stages
- B) Three sub-stages
- C) Five sub-stages (Leptotene, Zygotene, Pachytene, Diplotene, Diakinesis)
- D) Seven sub-stages

3. During which sub-stage of Prophase I do homologous chromosomes pair up (synapsis) to form tetrads?

- A) Leptotene
- B) Zygotene
- C) Diplotene
- D) Diakinesis

4. The X-shaped sites where non-sister chromatids exchange genetic material during Prophase I are called:

- A) Centromeres
- B) Chiasmata
- C) Kinetochores
- D) Telomeres

Answer Key

1. C — Prophase I is the longest meiotic phase, occupying over 90% of total meiotic duration.
2. C — Prophase I is divided into five distinct sub-stages: Leptotene, Zygotene, Pachytene, Diplotene, and Diakinesis.
3. B — Homologous pairing (synapsis) begins during the Zygotene sub-stage of Prophase I.
4. B — Chiasmata represent the physical crossover sites where non-sister chromatids exchange segments.

Lesson 65: Metaphase I of meiosis

1. What structures line up along the metaphase plate during Metaphase I of meiosis?

- A) Single unattached chromatids
- B) Tetrads (paired homologous chromosomes)
- C) Individual chromosomes like in mitosis
- D) Nuclear membranes

2. How do spindle microtubules attach to kinetochores in Metaphase I?

- A) Microtubules from one pole attach to both kinetochores of a tetrad
- B) Microtubules from opposite poles attach to kinetochores of opposing homologs
- C) Microtubules do not attach in Metaphase I
- D) Microtubules attach only to one homolog

3. In Metaphase I, how are the two chromosomes of each tetrad oriented?

- A) Both facing the same pole
- B) One chromosome facing one pole, the other facing the opposite pole
- C) Randomly without orientation
- D) Facing the cell center

4. How does Metaphase I of meiosis differ fundamentally from Metaphase of mitosis?

- A) Homologous pairs line up as tetrads in Metaphase I, unlike individual chromosomes in mitotic metaphase
- B) There is no difference
- C) Mitosis has tetrads
- D) Meiosis I lacks spindle fibers

Answer Key

- 1. B — Tetrads (paired homologous chromosomes) align at the metaphase plate during Metaphase I.
- 2. B — Microtubules extending from opposite poles attach to the kinetochores of opposing homologs.
- 3. B — One chromosome of each tetrad faces one pole, while the other chromosome faces the opposite pole.
- 4. A — Homologous pairs line up as tetrads in Metaphase I, unlike mitotic metaphase where individual chromosomes align.

Lesson 66: Anaphase I of meiosis

1. What structures separate and move toward opposite poles during Anaphase I of meiosis?

- A) Sister chromatids
- B) Homologous chromosome pairs
- C) Nuclear envelopes
- D) Spindle poles

2. What happens to sister chromatids during Anaphase I of meiosis?

- A) They split at their centromeres
- B) They remain joined at their centromeres and move together to the poles
- C) They dissolve completely
- D) They duplicate again

3. What guides the movement of separated chromosomes toward opposite poles during Anaphase I?

- A) Spindle apparatus
- B) Cytoplasmic streaming
- C) Nuclear pores
- D) Cell plate formation

4. How does Anaphase I of meiosis differ fundamentally from Anaphase of mitosis?

- A) Sister chromatids separate in Anaphase I
- B) Homologous chromosomes separate in Anaphase I, while sister chromatids stay joined
- C) No chromosomes move in Anaphase I
- D) Centromeres divide in Anaphase I

Answer Key

1. B — Homologous chromosome pairs separate from each other during Anaphase I.
2. B — Sister chromatids remain joined at their centromeres and move together to poles during Anaphase I.
3. A — Chromosome movement toward poles in Anaphase I is directed by the spindle apparatus.
4. B — Meiosis I keeps sister chromatids joined at centromeres, unlike mitosis where sister chromatids separate.

Lesson 67: Telophase I of meiosis

1. What is the structural state of chromosomes when they arrive at each pole during Telophase I?

- A) Single chromatids, haploid
- B) Haploid set, each chromosome still composed of two sister chromatids
- C) Diploid set of single chromatids
- D) Dissolved chromatin

2. How many daughter cells are formed at the completion of Telophase I and cytokinesis?

- A) Two haploid daughter cells
- B) Four haploid cells
- C) Two diploid cells
- D) One cell

3. Does DNA replication occur between the end of Meiosis I and the beginning of Meiosis II?

- A) Yes, a full S phase occurs
- B) No, chromosomes do not replicate between Meiosis I and Meiosis II
- C) Yes, but only partially
- D) Only in plant cells

4. How does cytokinesis proceed during Telophase I in animal cells compared to plant cells?

- A) Both use cell plates
- B) Animal cells divide via a contractile cleavage furrow, whereas plant cells construct a cell plate
- C) Both use cleavage furrows
- D) Cytokinesis does not occur

Answer Key

1. B — Each pole receives a haploid set of chromosomes, each still composed of two sister chromatids.

2. A — Telophase I and simultaneous cytokinesis divide a diploid cell into two haploid daughter cells.
3. B — No chromosome replication takes place between Meiosis I and Meiosis II.
4. B — Animal cells divide via a contractile cleavage furrow, whereas plant cells construct a cell plate.

Lesson 68: Prophase II of meiosis

1. What positioning event occurs to centrioles during Prophase II of meiosis?

- A) They dissolve permanently
- B) They separate and take up positions at opposite poles
- C) They remain at the center
- D) They duplicate

2. Is there a round of DNA replication immediately preceding Prophase II?

- A) Yes, a brief S phase occurs
- B) No, chromosomes are already replicated and do not replicate again
- C) Yes, in animal cells only
- D) Only the Y chromosome replicates

3. What happens to the nuclear membrane during Prophase II?

- A) It thickens into a cell wall
- B) It disappears
- C) It doubles
- D) It remains intact

4. Which structures form during Prophase II and radiate toward the center of the cell?

- A) Cleavage furrows
- B) Spindle fibers
- C) Cell plates
- D) Nuclear pores

Answer Key

1. B — Centrioles separate and take up positions at opposite poles during Prophase II.
2. B — Chromosomes do not replicate prior to Prophase II because they are already replicated.
3. B — The nuclear membrane disappears during Prophase II.
4. B — Spindle fibers form and radiate toward the center of the cell during Prophase II.

Lesson 69: Metaphase II of meiosis

1. What structures line up along the metaphase plate during Metaphase II of meiosis?

- A) Tetrads of homologous pairs
- B) Sister chromatids arranged across the middle of the cell
- C) Unreplicated single chromatids
- D) Nuclear envelopes

2. Microtubules extending from opposite poles attach to which specific structure on sister chromatids during Metaphase II?

- A) Telomeres
- B) Kinetochores

- C) Nucleoli
- D) Histone octamers

3. Why are the two sister chromatids of each chromosome no longer genetically identical during Metaphase II?

- A) Because of DNA replication errors
- B) Due to crossing over that occurred during Meiosis I
- C) Because of independent assortment
- D) Due to chromosome loss

4. Where are chromosomes positioned within the cell during Metaphase II?

- A) At the cell poles
- B) Lined up along the central metaphase plate in the middle of the cell
- C) Inside the nucleolus
- D) Attached to the cell membrane

Answer Key

1. B — Sister chromatids line up across the middle metaphase plate during Metaphase II.
2. B — Microtubules from opposing spindle poles attach to kinetochores on individual sister chromatids.
3. B — Crossing over during Prophase I recombines non-sister chromatids, making sister chromatids non-identical by Metaphase II.
4. B — Chromosomes align along the central metaphase plate in the middle of the cell during Metaphase II.

Lesson 70: Anaphase II of meiosis

1. What critical structural event permits sister chromatids to separate during Anaphase II?

- A) Splitting of centromeres
- B) Dissolution of the nuclear membrane
- C) DNA replication
- D) Formation of tetrads

2. What designation do separated sister chromatids take once they move apart in Anaphase II?

- A) Homologous pairs
- B) Individual daughter chromosomes
- C) Bivalents
- D) Tetrads

3. Where do separated chromatids migrate during Anaphase II?

- A) To the metaphase plate
- B) To opposite poles of the cell
- C) To the cell center
- D) Out of the cell

4. How does Anaphase II differ fundamentally from Anaphase I?

- A) Sister chromatids separate in Anaphase II, whereas homologous chromosomes separate in Anaphase I
- B) They are identical
- C) Anaphase I separates sister chromatids
- D) Anaphase II involves tetrads

Answer Key

1. A — Splitting of centromeres enables sister chromatids to separate during Anaphase II.
2. B — Once separated in Anaphase II, former sister chromatids become individual daughter chromosomes.
3. B — Separated chromatids are pulled toward opposite poles of the cell.
4. A — Anaphase II involves centromere splitting and sister chromatid separation, whereas Anaphase I separates homologous chromosomes.

Lesson 71: Telophase II of meiosis

1. How many total daughter cells are produced at the conclusion of Telophase II and cytokinesis?

- A) Two
- B) Four haploid daughter cells
- C) Eight
- D) One

2. How is the genetic composition of the four daughter cells described following Telophase II?

- A) All genetically identical
- B) Genetically distinct due to crossing over and independent assortment
- C) Diploid
- D) Triploid

3. What nuclear event takes place during Telophase II as chromosomes arrive at the poles?

- A) Nuclear membranes reassemble around haploid sets as chromosomes de-condense
- B) Nuclear membranes dissolve
- C) Chromosomes condense further
- D) Nucleoli duplicate

4. What chromosome set number is present in each resulting daughter cell after Telophase II?

- A) Diploid ($2n$)
- B) Haploid (n) set of unreplicated chromosomes
- C) Tetraploid
- D) Aneuploid

Answer Key

1. B — Meiosis II concludes with Telophase II and cytokinesis, producing four haploid daughter cells.
2. B — Crossing over and independent assortment ensure each of the four daughter cells is genetically unique.
3. A — Nuclear membranes reassemble around the haploid sets as chromosomes de-condense during Telophase II.
4. B — Each resulting daughter cell contains a single haploid (n) set of unreplicated chromosomes.

Lesson 72: Mitosis and meiosis

1. How does mitosis affect the number of chromosome sets in daughter cells relative to the parent cell?

- A) It halves the chromosome sets
- B) It conserves the number of chromosome sets, producing genetically identical cells

- C) It doubles the chromosome sets
- D) It eliminates sex chromosomes

2. What change in chromosome sets occurs as a result of meiotic cell division?

- A) Chromosome sets double
- B) Chromosome sets are reduced from diploid ($2n$) to haploid (n)
- C) Chromosome sets remain unchanged
- D) Chromosome sets become triploid

3. How do the daughter cells produced by meiosis compare genetically to one another and to the parent?

- A) They are genetically identical to the parent
- B) They are genetically distinct due to recombination and independent assortment
- C) They contain no genetic material
- D) They are twice the size of the parent cell

4. Which division process generates cells dedicated to growth, tissue repair, and asexual reproduction?

- A) Meiosis
- B) Mitosis
- C) Binary fission only
- D) Budding only

Answer Key

- 1. B — Mitosis maintains chromosome number and set counts, creating genetically identical daughter cells.
- 2. B — Meiosis reduces chromosome sets from diploid ($2n$) to haploid (n).
- 3. B — Genetic recombination and independent assortment make meiotic daughter cells genetically distinct.
- 4. B — Mitosis provides identical somatic cells needed for growth, repair, and tissue regeneration.

Lesson 73: Summary - Mitosis and meiosis

1. How many cell divisions occur in mitosis compared to meiosis?

- A) Mitosis has one division, whereas meiosis has two sequential divisions
- B) Both have one division
- C) Mitosis has two, meiosis has one
- D) Both have four divisions

2. During which phase does synapsis of homologous chromosomes and crossing over occur?

- A) Mitotic prophase
- B) Prophase I of meiosis
- C) Metaphase of mitosis
- D) Anaphase II

3. What is the primary role of meiosis in the animal body?

- A) To repair damaged tissue cells
- B) To produce haploid gametes while introducing genetic variability
- C) To grow organs
- D) To replace skin cells

4. How do the daughter cells of mitosis compare to those of meiosis in number and genetic composition?

- A) Mitosis yields two genetically identical diploid ($2n$) cells, whereas meiosis yields four genetically distinct haploid (n) cells
- B) Both yield identical cells
- C) Mitosis yields four haploid cells
- D) Meiosis yields two diploid cells

Answer Key

1. A — Mitosis involves a single nuclear division, whereas meiosis consists of two sequential divisions.
2. B — Synapsis and crossing over take place exclusively during Prophase I of meiosis.
3. B — Meiosis generates haploid gametes while introducing genetic variability among offspring.
4. A — Mitosis yields two genetically identical diploid ($2n$) cells, whereas meiosis yields four genetically distinct haploid cells.

Lesson 74: Chromosomal theory of inheritance**1. According to the Chromosomal Theory of Inheritance, where are genes physically located?**

- A) In the cytoplasm
- B) On chromosomes at specific loci
- C) In the mitochondria only
- D) In the cell membrane

2. The movement and behavior of chromosomes during cell division aligns with which laws?

- A) Lamarck's laws
- B) Mendel's laws of segregation and independent assortment
- C) Hardy-Weinberg principles
- D) Central dogma

3. Around what year did scientists begin to recognize parallels between Mendelian hereditary factors and chromosome behavior?

- A) 1865
- B) Around 1900
- C) 1953
- D) 1970

4. Which meiotic processes physically account for Mendel's laws of segregation and independent assortment?

- A) DNA replication and cytokinesis
- B) Segregation and independent sorting of chromosomes during meiosis
- C) Crossing over only
- D) Random fertilization only

Answer Key

1. B — The Chromosomal Theory of Inheritance establishes that genes occupy specific physical positions (loci) on chromosomes.
2. B — Chromosome movements during meiotic divisions directly follow Mendel's laws of segregation and independent assortment.

3. B — Parallels between Mendelian factors and cytogenetic chromosome behavior were first recognized around 1900.
4. B — Segregation and independent sorting of chromosomes during meiosis provide the physical basis for Mendel's laws.

Lesson 75: Experiment of T. H. Morgan

1. In T. H. Morgan's experiment with *Drosophila melanogaster*, which eye color was the wild type and which was mutant?

- A) Red wild-type, white mutant
- B) White wild-type, red mutant
- C) Both wild-type
- D) Brown wild-type

2. In Morgan's cross between red-eyed females and white-eyed males, what phenotype was observed in the F1 generation?

- A) All white-eyed
- B) All red-eyed
- C) Half red, half white
- D) All brown-eyed

3. In the F2 generation of Morgan's experiment, in which sex did the white-eyed mutant phenotype appear?

- A) Females only
- B) Males only (exclusively in male offspring)
- C) Both sexes equally
- D) Neither sex

4. What fundamental genetic conclusion did T. H. Morgan derive from his eye color experiment?

- A) Eye color is autosomal
- B) Eye color inheritance correlates with X chromosome transmission (sex linkage)
- C) Eye color is random
- D) Eye color follows blending inheritance

Answer Key

1. A — Red eye color represented the wild-type phenotype, while white eye color was the mutant.
2. B — All F1 offspring resulting from wild-type females crossed with mutant males displayed red eyes.
3. B — In the F2 generation, the white-eyed mutant trait appeared exclusively in male fruit flies.
4. B — Morgan demonstrated that eye color inheritance correlates with X chromosome transmission.

Lesson 76: Effect of linkage on inheritance

1. Which two traits did T. H. Morgan test in *Drosophila* to study the effect of gene linkage on inheritance?

- A) Body color (gray vs. black) and wing size (normal vs. vestigial)
- B) Eye color and bristle number
- C) Body color and eye color

D) Wing size and leg length

2. What testcross phenotypic ratio was expected if the genes for body color and wing size assorted independently?

- A) 3:1
- B) 9:3:3:1
- C) 1:1:1:1
- D) 1:2:1

3. In Morgan's testcross experiment, what were the two predominant phenotypic classes observed among progeny?

- A) Recombinant classes
- B) Parental phenotypes: wild-type (965) and double mutant
- C) All white-eyed
- D) All vestigial

4. What biological mechanism did Morgan propose to explain the presence of a smaller number of recombinant offspring?

- A) Independent assortment
- B) Crossing over breaking the physical linkage between genes
- C) Mutation
- D) Random drift

Answer Key

1. A — Morgan examined body color (gray vs. black) and wing size (normal vs. vestigial).
2. C — Independent assortment predicts a 1:1:1:1 phenotypic ratio among testcross offspring.
3. B — The vast majority of progeny exhibited parental phenotypes: wild-type (965) and double mutant.
4. B — Recombinant phenotypes arise when crossing over breaks the physical linkage between genes.

Lesson 77: Linkage map

1. What is a linkage map?

- A) A microscopic photograph of condensed metaphase chromosomes
- B) An ordered map of genetic loci along a chromosome based on recombination frequencies
- C) A pedigree chart
- D) A karyotype

2. What parameters are used to determine gene positions on a linkage map rather than physical base-pairs?

- A) Chromosome size
- B) Recombination frequencies (relative positions in terms of recombination)
- C) Gene expression levels
- D) Protein sizes

3. What is the relationship between the frequency of recombination between two genetic markers and their physical distance?

- A) A higher recombination frequency indicates a greater physical distance between two markers
- B) Higher recombination means shorter distance
- C) No relationship

D) Recombination only occurs in mitosis

4. Which molecular markers are explicitly mentioned in Lesson 77 as tools used in constructing linkage maps?

- A) Microsatellites and Restriction Fragment Length Polymorphisms (RFLPs)
- B) SNPs only
- C) Karyotypes
- D) Pedigrees

Answer Key

1. B — A linkage map orders genetic loci along a chromosome based on recombination frequencies.
2. B — Linkage map positions reflect the relative position of genes in terms of recombination frequency.
3. A — A higher recombination frequency indicates a greater physical distance between two markers.
4. A — Microsatellites and Restriction Fragment Length Polymorphisms (RFLPs) serve as key markers.

Lesson 78: Linkage map of drosophila chromosome

1. Which scientist first utilized recombination frequencies from fruit fly crosses to construct a chromosome map?

- A) Morgan
- B) Alfred Sturtevant
- C) Mendel
- D) Watson

2. How is one map unit (1 cM / centiMorgan) defined on a genetic linkage map?

- A) Equivalent to a 1% recombination frequency
- B) Equivalent to 1 base pair
- C) Equivalent to one chromosome
- D) Equivalent to 100 genes

3. Sturtevant mapped three *Drosophila* genes with recombination frequencies: $y-w = 1\%$, $w-m = 9.5\%$, $y-m = 17\%$. What is the correct relative order of these genes?

- A) $y - m - w$
- B) $y - w - m$ (w between y and m, since $1\% + 9.5\%$ is approximately the y-m distance)
- C) $m - y - w$
- D) $w - y - m$

4. Why does a greater distance between two genes increase their recombination frequency?

- A) Because greater distance provides more opportunity and crossover points between the genes
- B) Because genes attract each other
- C) Because of mutation
- D) Because of independent assortment

Answer Key

1. B — Alfred Sturtevant used recombination frequencies from fruit fly crosses to construct the first chromosome map.
2. A — One map unit (1 centiMorgan) is defined as equal to a 1% recombination frequency.

3. B — Because the smaller distances add up to the largest distance, the relative gene order places the middle gene between the other two.
4. A — A greater distance between genes provides more opportunity and crossover points along the chromosome.

Lesson 79: Sex determination and chromosomes role

- 1. Which specific gene located on the human Y chromosome triggers the development of male gonads (testes)?**
- A) SRY gene
 - B) BRCA1 gene
 - C) CFTR gene
 - D) HTT gene
- 2. What happens to generic embryonic gonads in an embryo that lacks a functional SRY gene?**
- A) They form testes
 - B) They develop into ovaries by default
 - C) They disappear completely
 - D) They become kidneys
- 3. Besides sex determination via SRY, what other essential role do genes on the Y chromosome serve?**
- A) Production of functional sperms
 - B) Determination of eye color
 - C) Regulation of blood clotting
 - D) Control of height
- 4. In the human X-Y sex determination system, what are the sex chromosome compositions of females and males?**
- A) Females XY, males XX
 - B) Females XX, males XY
 - C) Females XO, males XY
 - D) Females ZW, males ZZ

Answer Key

1. A — The SRY gene on the Y chromosome serves as the master switch initiating male testes development.
2. B — Without SRY gene activity, embryonic gonadal tissue defaults to developing into ovaries.
3. A — Y chromosome genes carry vital information required for spermatogenesis and functional sperm production.
4. B — Human females possess two X chromosomes (XX), while males possess one X and one Y chromosome (XY).

Lesson 80: Sex determination systems

- 1. In the X-O sex determination system observed in insects, what are the sex chromosome compositions of females and males?**
- A) Females XY and males XX
 - B) Females are XX and males are XO
 - C) Females are ZW and males are ZZ

D) Both females and males are XX

2. In birds, some fishes, and certain insects, which sex chromosome combination represents the female?

- A) XX
- B) XO
- C) ZW
- D) ZZ

3. In the haploid-diploid sex determination system found in bees and ants, how do male offspring develop?

- A) From fertilized diploid eggs
- B) From unfertilized ova, becoming haploid
- C) From diploid zygotes with ZW chromosomes
- D) Through mitotic fragmentation of sperm cells

4. In honeybees (*Apis mellifera*), what is the chromosome count of a diploid female worker or queen bee compared to a haploid male drone?

- A) Females have 46 chromosomes and males have 23 chromosomes
- B) Females have 32 chromosomes and males have 16 chromosomes
- C) Both females and males have 46 chromosomes
- D) Females have 23 chromosomes and males have 22 chromosomes

Answer Key

1. B — Under the X-O system in insects, females carry XX sex chromosomes while males carry a single X (XO).
2. C — In the Z-W sex determination system, females represent the heterogametic sex with ZW chromosomes.
3. B — Males in haploid-diploid systems (such as bees and ants) develop parthenogenetically from unfertilized haploid eggs.
4. B — Female bees develop from fertilized diploid eggs (32 chromosomes), whereas male drones develop from unfertilized haploid eggs (16 chromosomes).

Lesson 81: Sex linked genes

1. How is a sex-linked gene defined in genetics?

- A) A gene located on an autosomal chromosome pair
- B) A gene located on either sex chromosome
- C) A gene that codes exclusively for primary sex organs
- D) A gene present only in mitochondrial DNA

2. When a father with an X-linked recessive genetic disorder has children with an unmutated female, the mutant X chromosome is passed:

- A) To all of his sons, but none of his daughters
- B) To all of his daughters, but none of his sons
- C) To 50% of his sons and 50% of his daughters
- D) To none of his children

3. If a carrier female for an X-linked recessive trait mates with a normal male, what is the probability that her sons will be affected?

- A) 0%
- B) 25%
- C) 50%
- D) 100%

4. Do sex chromosomes carry genes for traits unrelated to sex determination?

- A) No, sex chromosomes carry only sex-determining genes
- B) Yes, sex chromosomes contain genes for many traits unrelated to sex
- C) Only Y chromosomes carry non-sex traits
- D) Sex chromosomes carry no functional genes

Answer Key

1. B — A sex-linked gene is defined as any gene located on either sex chromosome (X or Y).
2. B — An affected father passes his single mutant X chromosome to all daughters (making them carriers), but to none of his sons.
3. B — A carrier female ($X^A X^a$) has a 50% chance of passing the mutant X^a allele to each son ($X^a Y$).
4. B — Sex chromosomes carry numerous genes responsible for general biological functions unrelated to sex determination.

Lesson 82: Genetic mapping in humans

1. What is the primary purpose of genetic mapping in human medicine?

- A) To synthesize custom histones
- B) To provide crucial clues about which chromosome carries a disease gene
- C) To count chromosomes
- D) To measure height

2. Which single-gene human inherited disorders are explicitly listed in Lesson 82 as being identified by genetic mapping?

- A) Diabetes and hypertension
- B) Cystic fibrosis (and similar single-gene disorders)
- C) Common cold
- D) Baldness

3. How do geneticists perform human genetic mapping in practice?

- A) By exposing human cells to heavy radiation
- B) By analyzing DNA isolated from family pedigrees to identify polymorphic markers
- C) By karyotyping only
- D) By guessing

4. Which genetic markers are most commonly used on human genetic maps according to Lesson 82?

- A) Microsatellites and Single Nucleotide Polymorphisms (SNPs)
- B) Karyotypes
- C) Pedigrees
- D) Blood groups

Answer Key

1. B — Genetic mapping provides crucial clues about which chromosome carries a disease gene.
2. B — Genetic mapping successfully located single-gene disorders such as cystic fibrosis.
3. B — Human genetic mapping analyzes DNA isolated from family pedigrees to identify polymorphic markers.
4. A — Microsatellites and Single Nucleotide Polymorphisms (SNPs) represent the most commonly used markers.

Lesson 83: Multiple alleles

1. Multiple alleles are defined as how many alternative forms of a gene occupying the same locus on homologous chromosomes?

- A) One
- B) Three or more
- C) Exactly two
- D) Zero

2. What percentage of human genes are di-allelic (existing in two forms) according to Lesson 83?

- A) 10%
- B) 30%
- C) 50%
- D) 70%

3. What percentage of human genes exist in only one form (mono-allelic)?

- A) 30%
- B) 50%
- C) 70%
- D) 100%

4. Poly-allelic genes in humans are usually associated with which of the following?

- A) Ribosome size
- B) Tissue types (providing individual genetic identity, e.g., HLA)
- C) Hair color only
- D) Height only

Answer Key

1. B — Multiple alleles are defined as three or more alternative forms of a gene occupying the same locus.
2. B — The handout states that approximately 30% of human genes exist as di-allelic.
3. C — Roughly 70% of human genes are mono-allelic, existing in only one form.
4. B — Poly-allelic genes are associated with tissue types and provide an individual genetic signature.

Lesson 84: Multiple alleles - blood groups

1. What type of gene controls the ABO blood group system in humans?

- A) A mono-allelic gene
- B) A di-allelic gene
- C) A tri-allelic gene

D) A polygenic chromosome 18 gene

2. How many distinct genotypes can be generated by the three alleles of the ABO blood group system?

- A) 3
- B) 4
- C) 6
- D) 10

3. Which antigen is produced on the surface of red blood cells by allele i?

- A) Antigen A
- B) Antigen B
- C) Both Antigen A and B
- D) No antigen

4. Which ABO blood types have only one possible genotype each?

- A) Blood types A and B
- B) Blood types AB and O
- C) Blood types A and O
- D) Blood types B and AB

Answer Key

1. C — The ABO blood system is controlled by a tri-allelic gene consisting of Allele A, Allele B, and Allele i.
2. C — The three ABO alleles generate a total of 6 genotypes.
3. D — Allele I^A produces antigen A, Allele I^B produces antigen B, and allele i produces no antigen.
4. B — Blood types A and B can exist in homozygous or heterozygous forms, whereas blood types AB ($I^A I^B$) and O (ii) have only one genotype each.

Lesson 85: Blood groups and Transfusions

1. How does the human immune system interact with different blood group antigens during a blood transfusion?

- A) It ignores all blood group antigens
- B) It recognizes its own blood type as self and foreign blood types as non-self
- C) It destroys all red blood cells regardless of antigen type
- D) It converts non-self antigens into self antigens automatically

2. What type of antibodies are present in the plasma of an individual with Type A blood?

- A) Antibodies against Type A antigens (anti-A)
- B) Antibodies against Type B antigens (anti-B)
- C) Both anti-A and anti-B antibodies
- D) No antibodies against blood group antigens

3. What reaction occurs when anti-B antibodies in a recipient's plasma interact with donor red blood cells carrying Type B antigens?

- A) Hemolysis of white blood cells
- B) Agglutination (clumping) of red blood cells
- C) Conversion of Type B antigens to Type A
- D) Rapid duplication of blood platelets

4. What determines whether a foreign blood cell is targeted by host antibodies during a transfusion?

- A) The size of the donor red blood cells
- B) The presence of non-self surface antigens on the donor red blood cells
- C) The amount of hemoglobin inside host white cells
- D) The temperature of the donor blood bag

Answer Key

1. B — The immune system distinguishes self antigens from foreign non-self antigens on red blood cells.
2. B — Individuals with Type A blood produce antibodies against foreign Type B surface antigens.
3. B — Antibody binding to complementary foreign surface antigens causes red blood cell agglutination.
4. B — Host antibodies recognize foreign non-self surface antigens, triggering immune reactions against donor cells.

Lesson 86: Multiple alleles- rabbits

1. In rabbits, fur color is controlled by multiple alleles. Which allele produces the Agouti (dark gray) phenotype?

- A) Chinchilla allele
- B) Himalayan allele
- C) The dominant full-color allele (C)
- D) Albino allele

2. What phenotype is produced in a rabbit carrying the Himalayan allele in homozygous or dominant-recessive combination over white?

- A) Dark gray (Agouti)
- B) White fur with black paws, nose, ears, and tail
- C) Light gray (Chinchilla)
- D) Pure white

3. What is the correct hierarchy of dominance among the four rabbit fur color alleles?

- A) Albino > Himalayan > Chinchilla > Agouti
- B) Agouti (C) > Chinchilla (cch) > Himalayan (ch) > Albino (c)
- C) All are co-dominant
- D) Himalayan > Agouti > Albino > Chinchilla

4. A rabbit with the homozygous double-recessive genotype expresses which coat phenotype?

- A) Dark gray (Agouti)
- B) Light gray (Chinchilla)
- C) White with black points
- D) White (albino)

Answer Key

1. C — The dominant full-color allele specifies the dark gray (Agouti) phenotype.
2. B — The Himalayan allele produces white fur with black paws, nose, ears, and tail.
3. B — The dominance hierarchy for rabbit fur color is Agouti > Chinchilla > Himalayan > Albino.
4. D — The double recessive genotype results in a white (albino) coat.

Lesson 87: Alleles and genotypes

1. Which formula is used to calculate the total number of genotypes for a gene locus with n alleles?

- A) n^2
- B) $n(n+1)/2$
- C) $2n$
- D) $n!$

2. How is the number of homozygous genotypes determined for a gene with n alleles?

- A) Equal to n (the number of alleles)
- B) Equal to n^2
- C) Equal to $2n$
- D) Always equal to 1

3. What formula gives the number of heterozygous genotypes for a locus with n alleles?

- A) n
- B) $n(n-1)/2$
- C) $2n$
- D) n^2

4. For the 4 fur color alleles in rabbits, how many total genotypes can be generated?

- A) 4
- B) 6
- C) 10
- D) 16

Answer Key

1. B — The total number of genotypes for n alleles is calculated using $n(n+1)/2$.
2. A — The number of possible homozygous genotypes is equal to the number of alleles, n.
3. B — The number of heterozygous genotypes is given by $n(n-1)/2$.
4. C — For n = 4 alleles, $4(5)/2 = 10$ total genotypes are produced.

Lesson 88: Linkage and crossing over

1. How are linked genes defined in genetics?

- A) Genes located on different non-homologous chromosomes
- B) Genes located close together on the same chromosome
- C) Genes on sex chromosomes only
- D) Genes that never express

2. Genetically, linkage is defined as the failure of two genes to undergo which process?

- A) DNA replication
- B) Independent assortment
- C) Transcription
- D) Translation

3. During which cellular process do the phenomena of linkage and crossing over take place?

- A) Mitotic division

- B) Meiotic cell division
- C) Binary fission
- D) Budding

4. Linked genes tend to be inherited in what manner?

- A) Separately on different gametes
- B) Together as a unit (on the same physical chromosome)
- C) Randomly
- D) Only maternally

Answer Key

1. B — Linked genes are defined as genes located close together on the same chromosome.
2. B — Linkage is defined genetically as the failure of two genes to assort independently.
3. B — Linkage and crossing over are phenomena observed during meiotic cell division.
4. B — Because linked genes reside on the same physical chromosome, they tend to inherit together.

Lesson 89: Linked genes

1. What is the relationship between the physical distance separating two linked genes and their crossover frequency?

- A) Genes located closer together experience fewer crossover events than genes farther apart
- B) Closer genes cross over more
- C) Distance has no effect
- D) All genes cross over equally

2. On a genetic map, a crossover frequency of 1% corresponds to what distance?

- A) 0.1 micrometers
- B) 1 map unit (centiMorgan)
- C) 1 base pair
- D) 100 map units

3. If two genes show a crossover value of 12%, how far apart are they on a genetic map?

- A) 1 map unit
- B) 12 map units
- C) 120 map units
- D) 0.12 map units

4. How is crossover frequency used in chromosome mapping?

- A) It measures total nuclear volume
- B) It directly reflects relative map distance/units along a chromosome
- C) It counts ribosomes
- D) It measures cell size

Answer Key

1. A — Genes located closer together experience fewer crossover events than genes farther apart.
2. B — A 1% crossover value is defined as equivalent to 1 map unit.
3. B — A 12% crossover frequency indicates two genes are 12 map units apart.
4. B — Crossover frequency directly reflects relative map distance/units along a chromosome.

Lesson 90: Physical mapping

1. How is distance measured on a physical genome map compared to a genetic linkage map?

- A) In map units (centiMorgans)
- B) In base pairs (bp) or megabases (Mb) representing physical DNA distance
- C) In micrometers under light microscopy
- D) In number of histone octamers per loop

2. Which of the following is listed in Lesson 90 as a physical mapping technique?

- A) Monohybrid cross analysis
- B) In situ hybridization and restriction enzyme site mapping
- C) Pedigree analysis
- D) Feulgen hydrolysis testing

3. What serves as the fundamental basis for constructing a physical map of a chromosome?

- A) The frequency of recombination between markers
- B) The specific physical locations of landmarks along a DNA molecule
- C) The ratio of purines to pyrimidines
- D) The rate of cytosine methylation

4. What strategy characterizes top-down physical mapping?

- A) Random shotgun sequencing without prior mapping
- B) Ordering large-insert clones along a landmark map, selecting minimal overlaps, and subcloning for sequencing
- C) Mapping point mutations strictly through pedigree analysis
- D) Counting band numbers on metaphase chromosomes

Answer Key

- 1. B — Physical maps express distances directly in physical DNA units such as base pairs or megabases.
- 2. B — In situ hybridization, restriction enzyme site mapping, and cytogenetic mapping are physical mapping techniques.
- 3. B — Physical mapping relies on determining the physical coordinates of specific molecular landmarks along DNA.
- 4. B — Top-down physical mapping arranges large genomic clones into overlapping contigs before targeted subcloning and sequencing.

Lesson 91: Mapping diseased genes in Humans

1. What primary tools are used in human genetic mapping to locate disease genes in family pedigrees?

- A) Karyotyping only
- B) Pedigree analysis combined with DNA typing of polymorphic markers
- C) Blood tests only
- D) X-rays

2. What are the two common strategies used to identify diseased genes in human genetic mapping?

- A) Cloning and sequencing only
- B) Position-dependent and position-independent methods

- C) Karyotyping and pedigrees
- D) Drug trials

3. What does the "position-independent" method of mapping disease genes imply?

- A) Identifying a gene without prior location information (functional candidacy)
- B) Mapping without DNA
- C) Using only pedigrees
- D) Random selection

4. Why does having a higher density of DNA markers on a genetic map benefit disease gene identification?

- A) It looks nicer
- B) Dense marker coverage improves the probability of discovering a marker closely linked to the disease gene
- C) It reduces cost to zero
- D) It eliminates pedigrees

Answer Key

1. B — Human disease gene mapping relies on pedigree analysis combined with DNA typing of markers.
2. B — Two primary strategies for mapping disease genes are position-dependent and position-independent.
3. A — Position-independent mapping identifies disease genes without prior location information.
4. B — Dense marker coverage improves the probability of discovering a marker closely linked to the gene.

Lesson 92: Polygenic inheritance

1. How is polygenic inheritance defined in Lesson 92?

- A) A condition where a single gene controls multiple distinct traits
- B) A condition when two or more genes influence the expression of one trait
- C) A condition where DNA sequences are altered by UV radiation
- D) The complete loss of an autosomal chromosome

2. In the pepper color example, what phenotype results from the genotype R-/yy (red pigment present with chlorophyll)?

- A) Red
- B) Yellow
- C) Brown / orange
- D) Green

3. Crossing a pure-breeding brown pepper (RRyy) with a yellow pepper (rrYY) produces an F1 generation with which phenotype?

- A) All yellow
- B) All brown
- C) All red
- D) All green

4. What proportion of green peppers (rryy) appears in the F2 generation following self-fertilization of F1 red peppers?

- A) 9/16
- B) 3/16

- C) 1/16
- D) 1/4

Answer Key

1. B — Polygenic inheritance occurs when two or more genes influence the expression of a single trait.
2. C — Genotype R-/Y- produces red, R-/yy produces brown/orange, rr/Y- produces yellow, and rr/yy produces green.
3. C — Crossing RRyy and rrYY yields F1 hybrids with genotype RrYy, expressing the all-red phenotype.
4. C — Self-fertilizing F1 red peppers yields a 9:3:3:1 F2 phenotypic ratio, resulting in 1/16 green peppers (rryy).

Lesson 93: Examples of polygenic traits

1. Which of the following human traits is listed in Lesson 93 as an example of a polygenic/continuous trait?

- A) ABO blood group
- B) Eye color
- C) Presence of a Barr body
- D) Rh factor

2. In chicken comb shapes, what phenotype results from a genotype carrying dominant alleles at both loci (R-P-)?

- A) Rose comb
- B) Pea comb
- C) Single comb
- D) Walnut comb

3. What comb phenotype is expressed in chickens with the double homozygous recessive genotype (rrpp)?

- A) Walnut comb
- B) Rose comb
- C) Pea comb
- D) Single comb

4. Human hair color is controlled by alleles located on which four chromosomes?

- A) Chromosomes 1, 2, 3, and 4
- B) Chromosomes 3, 6, 10, and 18
- C) Chromosomes 5, 11, 13, and 21
- D) X and Y sex chromosomes only

Answer Key

1. B — Eye color, hair color, skin color, stature, shape of face, and fingerprint patterns are listed as polygenic traits.
2. D — Genotype R-P- produces Walnut combs, R-pp produces Rose combs, rrP- produces Pea combs, and rrpp produces Single combs.
3. D — Chickens carrying the double homozygous recessive genotype rrpp express Single comb shape.
4. B — Hair color is governed by alleles on chromosomes 3, 6, 10, and 18, with more dominant alleles yielding darker hair.

Lesson 94: Epistasis and modifier genes

1. How is positive epistasis characterized in Lesson 94?

- A) When two mutations together have a smaller effect than expected
- B) When a double mutation has a fitter phenotype than either single mutation
- C) When a mutation has the opposite effect in the presence of another mutation
- D) When two beneficial genes become deleterious together

2. What is negative epistasis?

- A) When a double mutation has a fitter phenotype than either single mutation
- B) When two mutations together have a smaller effect than expected from their individual effects
- C) When a mutation changes an amino acid to a stop codon
- D) When cytosine is methylated at carbon-5

3. Sign epistasis occurs when a mutation:

- A) Has no effect on protein function
- B) Has the opposite effect when in the presence of another mutation
- C) Reverts a mutant phenotype back to wild type
- D) Causes immediate embryonic lethality

4. Reciprocal epistasis is defined as a situation where:

- A) Two beneficial genes become lethal
- B) Two deleterious genes are beneficial when together
- C) A single gene affects multiple traits
- D) A gene is turned off by histone deacetylase

Answer Key

1. B — Positive epistasis between beneficial mutations generates improvements in function, producing a fitter phenotype in double mutants.
2. B — Negative epistasis occurs when two mutations together produce a smaller phenotypic effect than expected from their individual effects.
3. B — Sign epistasis is defined as occurring when one mutation exerts the opposite effect in the presence of another mutation.
4. B — Reciprocal epistasis is the case where two individually deleterious genes are beneficial when combined.

Lesson 95: Recessive Epistasis

1. How is recessive epistasis defined?

- A) An epistatic gene exerts its masking effect when present as a homozygous recessive
- B) An epistatic gene acts only when dominant
- C) It is the same as dominance
- D) It involves only one gene

2. In blue-eyed Mary plants, petal color is governed by recessive epistasis. What color are flowers when both enzymes are active?

- A) White
- B) Magenta
- C) Blue (anthocyanin)

D) Yellow

3. In the biochemical pathway of blue-eyed Mary plants, what product accumulates when Enzyme 1 is active?

- A) White pigment
- B) Precursor 1 (colorless) converts to precursor 2 (magenta)
- C) Blue pigment directly
- D) Nothing

4. In pigeons, which gene exhibits recessive epistasis over spread color and pattern genes?

- A) Ash-red
- B) Recessive red
- C) Blue
- D) Brown

Answer Key

1. A — In recessive epistasis, the epistatic gene masks other loci when present in the homozygous recessive state.
2. C — When both enzymes are active, precursor molecules are converted to blue anthocyanin pigment.
3. B — If Enzyme 1 is active, precursor 1 (colorless) converts to precursor 2 (magenta).
4. B — The recessive red gene in pigeons acts epistatically over both spread color and pattern genes.

Lesson 96: Dominant epistasis

1. How does dominant epistasis differ from recessive epistasis?

- A) The epistatic gene exerts its masking effect when present as a single dominant allele
- B) It requires two recessive alleles
- C) There is no difference
- D) It involves cytoplasmic genes

2. In summer squash fruit color, what effect does the dominant allele W have on fruit pigmentation?

- A) It produces yellow fruit color
- B) It produces green fruit color
- C) It inhibits color formation, yielding white fruit
- D) It has no effect

3. In summer squash, if a plant lacks the dominant inhibitor (ww) but carries dominant Y, what will be its fruit color phenotype?

- A) White
- B) Yellow
- C) Green
- D) Brown

4. In Shepherd's purse (Capsella bursa-pastoris), duplicate dominant epistasis controls fruit shape. What shape results from presence of a dominant allele?

- A) Narrow shape
- B) Heart (triangular) shape
- C) Oval shape

D) Circular shape

Answer Key

1. A — Dominant epistasis occurs when a single dominant allele at the epistatic locus masks another gene.
2. C — The dominant allele W inhibits color formation in summer squash, yielding white fruit.
3. B — In the absence of the dominant inhibitor W, dominant Y produces yellow fruit.
4. B — Presence of a dominant allele yields heart-shaped fruit, whereas double homozygous recessive produces narrow fruit.

Lesson 97: Example of Epistasis

1. In Labrador Retrievers, coat color is determined by two genes (B and E). What role does Gene B play?

- A) It controls pigment deposition
- B) It determines pigment color: dominant B produces black pigment, while recessive b produces brown pigment
- C) It has no role
- D) It controls tail length

2. What is the epistatic role of Gene E in Labrador Retriever coat color?

- A) Homozygous recessive ee prevents dark pigment deposition, resulting in golden/yellow fur regardless of Gene B
- B) Dominant E produces white fur
- C) Gene E has no effect on coat color
- D) Recessive e forces the coat to become black

3. What coat color phenotype is expressed by a Labrador Retriever with genotype bbE_?

- A) Black
- B) Brown (Chocolate)
- C) Golden (Yellow)
- D) Spotted

4. Which of the following genotypes will produce a Golden (Yellow) Labrador Retriever?

- A) BBEE
- B) bbEE
- C) BbEe
- D) Any genotype containing homozygous recessive ee (e.g., BBee)

Answer Key

1. B — Gene B determines pigment color: dominant B produces black pigment, while recessive b produces brown pigment.
2. A — Homozygous recessive ee blocks dark pigment deposition in fur, yielding a golden/yellow coat regardless of alleles at Gene B.
3. B — Genotypes bbE_ combine brown pigment alleles (bb) with functional pigment deposition (E_), producing brown fur.
4. D — Any genotype containing homozygous recessive ee results in a golden (yellow) retriever.

Lesson 98: Epigenetics

1. What does the term "epigenetics" literally mean?

- A) Below the genetics
- B) Above the genetics
- C) Outside of the nucleus
- D) Within the chromosome

2. How is an epigenetic trait defined in Lesson 98?

- A) A permanent change in nuclear base sequence caused by radiation
- B) A stable and heritable phenotype resulting from changes in chromosome without alterations in the DNA sequence
- C) A numerical abnormality resulting from meiotic non-disjunction
- D) A point substitution that changes a purine to a pyrimidine

3. Cytosine methylation involves adding a methyl group to which carbon position on cytosine residues?

- A) Carbon-1
- B) Carbon-3
- C) Carbon-5
- D) Carbon-6

4. Does DNA methylation alter the primary base sequence of DNA?

- A) Yes, it converts cytosine into thymine permanently
- B) No, DNA methylation represents an epigenetic change without altering the DNA sequence
- C) Yes, it deletes three consecutive base pairs
- D) Yes, it reverses the sequence orientation by 180 degrees

Answer Key

- 1. B — Epigenetics literally translates to 'above' genetics.
- 2. B — An epigenetic trait is defined as a stable, heritable phenotype arising from chromosome changes without altering the underlying DNA sequence.
- 3. C — Cytosine methylation specifically attaches a methyl group to carbon-5 of cytosine residues.
- 4. B — DNA methylation is an epigenetic modification that regulates gene expression without changing the DNA base sequence.

Lesson 99: Epigenetic Modifications

1. Which of the following is listed in Lesson 99 as one of the major steps/mechanisms of epigenetic modification?

- A) DNA replication
- B) Cytosine methylation / histone modification
- C) Protein denaturation
- D) Mitosis

2. At which carbon position on the cytosine ring does DNA methylation attach a methyl group?

- A) Carbon-1
- B) Carbon-3
- C) Carbon-5
- D) Carbon-6

3. What effect does histone acetylation (mediated by HAT) typically have on chromatin structure and gene expression?

- A) Condenses chromatin, silencing genes
- B) Relaxes chromatin into an open conformation, increasing gene expression
- C) No effect
- D) Destroys histones

4. Which enzymes are responsible for adding and removing acetyl groups on histone tails, respectively?

- A) HDAC then HAT
- B) HAT (Histone acetyltransferase) adds; HDAC (Histone deacetylase) removes
- C) DNA polymerase
- D) Ligase and helicase

Answer Key

1. B — Epigenetic modification mechanisms include cytosine methylation, histone modification, and RNA-based mechanisms.
2. C — DNA methylation specifically adds a methyl group to carbon-5 of cytosine residues.
3. B — Histone acetylation relaxes chromatin into an open conformation, increasing gene expression.
4. B — HAT (Histone acetyltransferase) attaches acetyl groups, whereas HDAC (Histone deacetylase) removes them.

Lesson 100: Common Epigenetics regulated phenomena

1. Which phenomenon involves down-regulating gene expression on one female X chromosome via DNA methylation?

- A) Genomic imprinting
- B) X chromosome inactivation
- C) Fragile X syndrome
- D) Mitosis

2. What is Genomic Imprinting?

- A) Epigenetic marking of a locus based on parental origin, causing monoallelic expression
- B) DNA mutation
- C) Chromosome loss
- D) Protein synthesis

3. Fragile X syndrome is linked to a CGG repeat expansion in the FMR1 gene. Hypermethylation and gene silencing occur beyond how many repeats?

- A) 50
- B) 100
- C) 200 CGG repeats
- D) 1000

4. Why do identical twins with identical genetic sequences develop phenotypic differences as they age?

- A) Different DNA sequences
- B) Environmental differences over time alter twin epigenomes, creating distinct individual patterns
- C) Different parents

D) Random mutation only

Answer Key

1. B — X chromosome inactivation silences one female X chromosome via epigenetic marks.
2. A — Genomic imprinting marks gene loci based on parental origin, expressing only the maternal or paternal allele.
3. C — Expansion beyond 200 CGG repeats correlates with hypermethylation at CpG dinucleotides and gene silencing.
4. B — Environmental differences over time alter twin epigenomes, creating distinct individual epigenetic patterns.

Lesson 101: Therapies for epigenetics

1. Unlike genetic sequence mutations, how are epigenetic modifications characterized regarding therapeutic potential?

- A) They are completely permanent and irreversible
- B) Epigenetic changes can be reversed
- C) They affect only mitochondrial DNA bases
- D) They permanently destroy histone core proteins

2. What is the mechanism of action of Vorinostat, an FDA-approved drug targeting epigenetics?

- A) DNA methyltransferase activator
- B) Histone deacetylase inhibitor
- C) Ribosomal RNA inhibitor
- D) DNA polymerase blocker

3. What effect does inhibiting histone deacetylase (increasing histone acetylation) have on gene expression?

- A) Decreases gene expression via chromatin condensation
- B) Increases gene expression by relaxing chromatin structure
- C) Halts DNA replication completely
- D) Promotes non-disjunction during meiosis

4. In ovarian cancer epigenetics, hypermethylation of tumor suppressor genes or hypomethylation of oncogenes can lead to:

- A) Enhanced DNA repair mechanisms
- B) Inactivation of tumor suppressors or activation of oncogenes leading to carcinogenesis
- C) Complete reversal of cellular aging
- D) Doubling of autosomal chromosome counts

Answer Key

1. B — In contrast to permanent genetic mutations, epigenetic modifications can be reversed through targeted therapies.
2. B — Vorinostat functions specifically as a histone deacetylase inhibitor.
3. B — Inhibiting deacetylase increases histone acetylation, which relaxes chromatin condensation and increases gene expression.
4. B — Hypermethylation of tumor suppressors or hypomethylation of oncogenes contributes to tumor progression and chemoresistance.

Lesson 102: Environmental factors and Epigenetics

- 1. According to Lesson 102, which maternal factor can directly influence the epigenome of a developing fetus?**
 - A) Eye color
 - B) Maternal nutrition and stress hormones
 - C) Blood type
 - D) Height
- 2. What role do lifestyle factors such as social interactions, physical activity, and diet play in epigenetics?**
 - A) No role
 - B) They are key factors that modify the epigenome
 - C) They change DNA sequence
 - D) They prevent all disease
- 3. How does the study of environmental epigenetics impact the "nature vs. nurture" debate?**
 - A) It proves only nature matters
 - B) It shows environmental influences directly modify gene expression via epigenetic marks
 - C) It ends all research
 - D) It proves nurture is irrelevant
- 4. Unlike permanent DNA sequence mutations, how are epigenetic modifications characterized regarding reversibility?**
 - A) They are permanent
 - B) They are dynamic and reversible through targeted pharmacological intervention
 - C) They cannot be studied
 - D) They always cause cancer

Answer Key

1. B — Maternal nutrition and stress hormones travel to the fetus and affect its developing epigenome.
2. B — Social interactions, physical activity, diet, and toxin exposure are key factors shaping the epigenome.
3. B — Environmental epigenetics shows that environmental influences directly modify gene expression.
4. B — Epigenetic modifications are dynamic and reversible through targeted pharmacological agents.

Lesson 103: Epigenetics- conclusions

- 1. What are the two prominent epigenetic mechanisms highlighted in the conclusions of Lesson 103?**
 - A) DNA methylation (gene silencing) and histone acetylation (gene activation)
 - B) Mitosis and meiosis
 - C) Replication and transcription
 - D) Mutation and selection
- 2. Which statement correctly describes how the epigenome changes over an individual's lifespan?**
 - A) It never changes
 - B) The epigenome is dynamic across an individual's lifetime, changing with age and environment
 - C) It changes only at birth

D) It is identical in all tissues always

3. What role do errors in epigenetic patterns play in human health according to Lesson 103?

- A) They have no effect
- B) Aberrant or errored epigenetic patterns contribute to human disease states, including cancer
- C) They improve health
- D) They only affect plants

4. What represents a major area of current research in epigenetic therapies?

- A) Synthesizing custom histones
- B) Developing pharmacological agents/drugs capable of modifying epigenetic marks
- C) Gene deletion
- D) Cloning

Answer Key

1. A — DNA methylation (gene silencing) and histone acetylation (gene activation) are two prominent mechanisms.
2. B — The epigenome is dynamic across an individual's lifetime, changing with age and environment.
3. B — Aberrant or errored epigenetic patterns contribute to human disease states, including cancer.
4. B — Research actively focuses on developing pharmacological agents/drugs capable of influencing epigenetic patterns.

Lesson 104: Penetrance and Expressivity

1. How is penetrance defined in genetics?

- A) The severity of symptoms observed in a single clinical case
- B) The proportion of individuals carrying a specific genotype who express the associated phenotype
- C) The number of genes involved
- D) The rate of mutation

2. What is expressivity defined as?

- A) The proportion of gametes carrying a mutant chromosome
- B) The extent or degree to which a particular genotype is expressed phenotypically
- C) The dominance of an allele
- D) The frequency of recombination

3. Reduced or incomplete penetrance (less than 100%) can result from which of the following factors?

- A) Genetic background factors (modifiers, epistasis) and environmental influences
- B) Nothing; penetrance is always complete
- C) Only age
- D) Only diet

4. In clinical medicine, the term "variable expressivity" is analogous to:

- A) The exact chromosome count
- B) Differing degrees or severity of phenotypic expression among individuals with the same genotype
- C) Complete penetrance
- D) Lethal mutations

Answer Key

1. B — Penetrance is defined as the proportion of individuals carrying a specific genotype who show the phenotype.
2. B — Expressivity describes the extent or degree to which a particular genotype is expressed.
3. A — Incomplete penetrance is influenced by genetic background factors (modifiers, epistasis) and environment.
4. B — Variable expressivity refers to differing degrees or severity of phenotypic expression.

Lesson 105: Types of Penetrance

1. When an allele exhibits complete penetrance, what proportion of individuals carrying the disease-causing mutation express clinical symptoms?

- A) All individuals (100%)
- B) Fewer than 10% of individuals
- C) Only homozygous recessive males
- D) Only individuals exposed to specific environmental toxins

2. How is incomplete (or reduced) penetrance defined in Lesson 105?

- A) Every individual carrying the allele shows 100% phenotypic severity
- B) Some individuals fail to express the trait even though they carry the allele
- C) The allele is expressed exclusively in response to soil pH
- D) The mutation permanently alters the base sequence of mitochondrial DNA

3. Why is low penetrance particularly challenging for geneticists to analyze?

- A) Because it is difficult to distinguish environmental factors from genetic factors
- B) Because it deletes both telomeres of the chromosome
- C) Because it affects only prokaryotic circular plasmids
- D) Because it converts euchromatin into heterochromatin in G0 phase

4. Multiple endocrine neoplasia 1 (MEN 1) is cited as an example of what type of penetrance phenomenon?

- A) Soil pH-dependent penetrance
- B) Age-related penetrance (7% at age 10, 100% at age 60)
- C) Complete non-penetrance across all ages
- D) Sex-limited penetrance in females only

Answer Key

1. A — An allele has complete penetrance when all individuals carrying the disease-causing mutation express clinical symptoms.
2. B — Reduced or incomplete penetrance occurs when some individuals carrying the allele fail to express the trait.
3. A — In low penetrance, it is difficult to distinguish environmental from genetic factors.
4. B — MEN 1 is an age-related penetrance disorder, rising from 7% at age 10 to 100% at age 60.

Lesson 106: Chromosomal Aberrations

1. How is a chromosomal aberration defined in Lesson 106?

- A) Any point mutation changing arginine to histidine

- B) Any change in the normal number or structure of chromosomes
- C) A change in flower color brought about by soil pH
- D) Cytosine methylation at carbon-5

2. Numerical chromosomal aberrations usually arise from what type of error?

- A) De novo errors during meiotic division
- B) Dietary protein deficiency in adulthood
- C) Histone deacetylase enzyme activation
- D) DNA polymerase slippage during mitosis

3. What mechanism explains why chromosomal aberrations cause clinical anomalies in affected individuals?

- A) By altering the amounts of gene products and disrupting gene balance
- B) By converting linear chromosomes into circular plasmids
- C) By destroying all cellular ribosomes
- D) By increasing the G-C percentage across the entire genome

4. Chromosomal aberrations occurring post-zygotically and giving rise to different cell lines within an individual result in:

- A) Complete penetrance
- B) Mosaicism
- C) Polygenic epistasis
- D) Reciprocal translocation

Answer Key

1. B — A chromosomal aberration is defined as any change in the normal number or structure of chromosomes.
2. A — Numerical chromosomal changes are usually due to de novo errors during meiosis.
3. A — Altered gene product amounts cause anomalies directly or disturb the balance of genes.
4. B — Aberrations in different cell lines occurring post-zygotically result in mosaicism.

Lesson 107: Karyotyping - Nomenclature

1. In human cytogenetic nomenclature, how is a normal male karyotype designated?

- A) 46, XX
- B) 46, XY
- C) 47, XXY
- D) 45, X

2. In the karyotype nomenclature "46, XY", what does the number 46 represent?

- A) The number of autosomes only
- B) The total count of chromosomes
- C) The number of genes
- D) The patient's age

3. What does the designation "XY" signify in karyotype nomenclature?

- A) The sex chromosome constitution
- B) The autosome number
- C) A disease

D) A band

4. How are chromosomal abnormalities or variants recorded in standardized cytogenetic nomenclature?

- A) They are ignored
- B) Specific numerical or structural abnormalities are appended after the sex chromosome designation (e.g., 47, XY, +21)
- C) Only verbally
- D) With colors

Answer Key

1. B — Standard human cytogenetic nomenclature designates a normal male chromosome constitution as 46, XY.
2. B — The leading integer in karyotype notation indicates the total count of chromosomes.
3. A — The letter pair following the chromosome count designates the sex chromosome constitution.
4. B — Specific numerical or structural abnormalities are appended immediately after the basic karyotype designation.

Lesson 108: Defining chromosome regions

1. What microscopic features are used to identify and classify individual chromosomes in a karyotype?

- A) Overall size, centromere position, and banding pattern
- B) Color only
- C) Weight
- D) Smell

2. How are chromosome arms designated in cytogenetic mapping?

- A) Short arm is 'p' (petite) and long arm is 'q'
- B) Short arm is 'q', long arm is 'p'
- C) Both arms are 'p'
- D) Arms are not designated

3. How are specific sub-regions and bands numbered along a chromosome arm?

- A) Numbered sequentially outward from the centromere
- B) Randomly
- C) From telomere inward only
- D) They are not numbered

4. What does the concept of "bands within bands" refer to in cytogenetics?

- A) Overlapping histone tails
- B) High-resolution cytogenetics resolving major chromosome bands into finer sub-bands and sub-sub-bands
- C) Staining errors
- D) Broken chromosomes

Answer Key

1. A — Chromosomes are identified and categorized by overall size, centromere position, and banding pattern.
2. A — Standard cytogenetic convention designates the short arm as 'p' (petite) and the long arm as 'q'.

3. A — Region and band numbers increase progressively moving outward from the centromere.
4. B — High-resolution cytogenetics resolves major chromosome bands into finer sub-bands.

Lesson 109: Common numerical abnormalities

1. Which numerical chromosomal abnormality corresponds to Down syndrome?

- A) Trisomy 13 (47, XX, +13)
- B) Trisomy 18
- C) Trisomy 21 (47, XX, +21 or 47, XY, +21)
- D) Monosomy X

2. Edwards syndrome and Patau syndrome represent which specific trisomies, respectively?

- A) Trisomy 18 and trisomy 13
- B) Trisomy 21 and trisomy 18
- C) Trisomy 13 and trisomy 21
- D) Monosomy and triploidy

3. What is the sex chromosome constitution of Klinefelter syndrome?

- A) 45, X
- B) 47, XXY
- C) 47, XXX
- D) 69, XXY

4. Triploidy is a numerical abnormality characterized by a total chromosome count of:

- A) 47 chromosomes
- B) 69 chromosomes (three complete haploid sets)
- C) 92 chromosomes
- D) 46 chromosomes

Answer Key

1. C — Down syndrome is caused by trisomy 21, designated as 47, XX, +21 (or 47, XY, +21).
2. A — Edwards syndrome is trisomy 18, whereas Patau syndrome is trisomy 13.
3. B — Klinefelter syndrome is characterized by an extra X chromosome in males, yielding 47, XXY.
4. B — Triploidy involves three complete haploid sets, resulting in 69 total chromosomes.

Lesson 110: Structural Abnormalities

1. Structural chromosome abnormalities result fundamentally from chromosome breakage followed by:

- A) Cell death
- B) Reunion of broken segments in abnormal configurations
- C) DNA replication
- D) Protein synthesis

2. How is a "balanced" structural chromosome rearrangement defined?

- A) Extra genetic material is present
- B) A normal, complete gene complement with rearranged order (no net gain or loss)
- C) Missing chromosomes
- D) Extra chromosomes

3. An "unbalanced" structural abnormality occurs when there is:

- A) An incorrect amount of genetic material (missing or duplicated segments)
- B) A normal gene complement
- C) No change at all
- D) Only inversions

4. Which of the following is listed in Lesson 110 as a type of structural chromosomal abnormality?

- A) Translocations, deletions, insertions, inversions, ring chromosomes, and isochromosomes
- B) Trisomy only
- C) Monosomy only
- D) Polyploidy only

Answer Key

1. B — Structural aberrations arise when chromosomes break and their segments reunite in abnormal patterns.
2. B — Balanced structural rearrangements contain a normal, complete gene complement with altered arrangement.
3. A — Unbalanced structural alterations result in missing or duplicated genetic material.
4. A — Translocations, deletions, insertions, inversions, ring chromosomes, and isochromosomes are listed types.

Lesson 111: Translocations

1. How is a translocation defined in cytogenetics?

- A) Loss of a single chromosome from a diploid set
- B) The transfer of genetic material from one chromosome to another
- C) Rotation of a chromosome segment by 180 degrees
- D) Doubling of the entire haploid genome

2. What are the two primary structural types of translocations described in Lesson 111?

- A) Point and frameshift translocations
- B) Reciprocal translocation and Robertsonian translocation
- C) Pericentric and paracentric translocations
- D) Monosomic and trisomic translocations

3. What defines a "balanced" translocation?

- A) A translocation where some genes are deleted or duplicated
- B) A translocation where no genetic information is extra or missing
- C) A translocation occurring exclusively on sex chromosomes
- D) A translocation involving centrosome duplication

4. What occurs during a reciprocal translocation?

- A) Two non-homologous chromosomes exchange genetic material between each other
- B) An entire chromosome arm is lost and degraded
- C) Two acrocentric centromeres fuse permanently
- D) A single base substitution converts an amino acid into a stop codon

Answer Key

1. B — A translocation is defined as the transfer of genetic material from one chromosome to another.
2. B — The handout classifies translocations into reciprocal translocations and Robertsonian translocations.
3. B — In a balanced translocation, the chromosome complement is complete with no extra or missing genes.
4. A — Reciprocal translocation involves a mutual exchange of chromosomal material between two different non-homologous chromosomes.

Lesson 112: Non-disjunction

1. What is defined as the failure of two homologous chromosomes or sister chromatids to separate during cell division?

- A) Crossing over
- B) Non-disjunction
- C) Polyploidy
- D) Translocation

2. What is the outcome if non-disjunction occurs during the first meiotic division?

- A) Only 50% of gametes are abnormal
- B) All sperms or ova from primary spermatocytes or oocytes will be abnormal
- C) All daughter cells become triploid
- D) Sister chromatids dissolve completely

3. What types of daughter cells are produced as a result of mitotic non-disjunction?

- A) Only haploid daughter cells
- B) Trisomic and monosomic daughter cells
- C) Polyploid and octaploid daughter cells
- D) Identical diploid daughter cells

4. Which of the following is listed as a cause of non-disjunction?

- A) Colchicine treatment during interphase
- B) Aging effect on primary oocytes, radiation, and delayed fertilization after ovulation
- C) High ambient environmental temperatures
- D) Histone acetylation in G1 phase

Answer Key

1. B — Non-disjunction is defined as the failure of homologous chromosomes or sister chromatids to separate during cell division.
2. B — If non-disjunction takes place during the 1st meiotic division, all sperms/ova derived from primary spermatocytes/oocytes will be abnormal.
3. B — Mitotic non-disjunction results in trisomic and monosomic daughter cells.
4. B — Causes of non-disjunction include the aging effect on primary oocytes, radiation exposure, and delayed fertilization after ovulation.

Lesson 113: Trisomy

1. How is trisomy defined in terms of chromosome composition?

- A) Having four copies of a particular chromosome
- B) Having three copies of a particular chromosome instead of the normal two
- C) Gaining an extra complete haploid set of chromosomes

D) Lacking a single sex chromosome

2. Which structural type of trisomy is characterized by an extra chromosome possessing duplicated arms?

- A) Primary trisomy
- B) Partial trisomy
- C) Secondary trisomy
- D) Tertiary trisomy

3. Which type of trisomy occurs when the extra chromosome consists of copies of arms from two other distinct chromosomes?

- A) Primary trisomy
- B) Secondary trisomy
- C) Tertiary trisomy
- D) Partial trisomy

4. What is the incidence rate of Trisomy 21 (Down syndrome) in live births, and what maternal factor is it associated with?

- A) 1 in 100 live births; associated with young maternal age
- B) 1 in 700 live births; associated with increased maternal age
- C) 1 in 5,000 live births; associated with maternal stress
- D) 1 in 10,000 live births; associated with low maternal nutrition

Answer Key

1. B — Trisomy is a form of polysomy and aneuploidy where three copies of a chromosome exist instead of the normal two.
2. C — Secondary trisomy occurs when the extra chromosome has duplicated arms.
3. C — Tertiary trisomy occurs when the extra chromosome is composed of copies of arms from two other chromosomes.
4. B — Trisomy 21 occurs in 1 in 700 live births and shows a strong association with increased maternal age.

Lesson 114: Monosomy

1. Monosomy is defined as the loss of:

- A) A complete haploid set of chromosomes
- B) A single chromosome from a diploid pair ($2n - 1$)
- C) Both homologs
- D) The entire genome

2. What is the general developmental outcome of autosomal monosomy in humans?

- A) Mild learning disability
- B) Incompatible with life (lethal)
- C) No effect
- D) Enhanced growth

3. Which human syndrome is a viable sex chromosome monosomy?

- A) Down syndrome
- B) Turner syndrome ($45, X / XO$)
- C) Klinefelter syndrome

D) Edwards syndrome

4. What are the two primary causes of monosomy outlined in Lesson 114?

- A) Meiotic/mitotic non-disjunction or chromosome loss during cell division (anaphase lag)
- B) Old age only
- C) Diet
- D) Radiation only

Answer Key

1. B — Monosomy represents the loss of a single chromosome from a diploid pair ($2n - 1$).
2. B — Monosomy involving any human autosome is incompatible with life and results in lethality.
3. B — Turner syndrome (XO / 45, X) is a viable human condition resulting from sex chromosome monosomy.
4. A — Monosomy arises through meiotic/mitotic non-disjunction or chromosome loss during division.

Lesson 115: Polyploidy

1. How is polyploidy defined in genetics?

- A) The loss of an autosome from a diploid set
- B) Having chromosome numbers that are multiples of the haploid number
- C) A point substitution that converts an amino acid into a stop codon
- D) A 180-degree rotation of a chromosome segment

2. What is the survival outcome of a polyploid fetus in humans?

- A) It develops into a normal fertile adult
- B) In humans, a polyploid fetus does not survive
- C) It develops into a viable adult with Klinefelter's syndrome
- D) It survives with mild physical anomalies

3. Which of the following is listed as a cause of polyploidy?

- A) Cytosine methylation
- B) Retention of a polar body, formation of diploid sperms, or di-spermy
- C) Histone deacetylase activation
- D) Equal meiotic crossing over

4. Which plant species is cited as an example of a hexaploid (6x) polyploid?

- A) Watermelon
- B) Cotton
- C) Wheat
- D) Strawberry

Answer Key

1. B — Polyploidy involves chromosome counts that are multiples of the haploid number (such as triploidy or tetraploidy).
2. B — Human polyploid fetuses do not survive.
3. B — Causes of polyploidy include retention of a polar body, formation of diploid sperms, and di-spermy (fertilization by two sperms).
4. C — Wheat (along with oat and kiwifruit) is listed as a hexaploid (6x) plant species.

Lesson 116: Allopolyploidy

1. What defines allopolyploidy compared to autopolyploidy?

- A) Allopolyploids have multiple chromosome sets derived from a single species
- B) Allopolyploids are polyploids with chromosomes derived from different species
- C) Allopolyploids possess only single-stranded RNA genomes
- D) Allopolyploids are generated exclusively through mitotic non-disjunction in somatic cells

2. How are offspring resulting from an inter-species allopolyploid cross generally described regarding fertility?

- A) Highly fertile
- B) Generally sterile
- C) Completely asexual
- D) Permanently haploid

3. Which chemical agent is used to promote polyploidy in plants?

- A) Colchicine
- B) Vorinostat
- C) Quinacrine
- D) Ethidium bromide

4. How does colchicine promote polyploidy at the cellular level?

- A) By binding to tubulin, disrupting microtubule formation, and blocking chromosome segregation
- B) By acetylating histone H1 tails
- C) By cleaving CEN DNA at CDE-III sub-domains
- D) By methylating cytosine residues at carbon-5

Answer Key

- 1. B — Allopolyploids carry chromosomes derived from different species through inter-species hybridization.
- 2. B — Offspring produced from inter-species crosses in allopolyploidy are generally sterile.
- 3. A — Colchicine is used to promote polyploidy and allopolyploidy in plants.
- 4. A — Colchicine binds to tubulin proteins, disrupting spindle microtubules and preventing chromosome segregation.

Lesson 117: Aneuploidy

1. How is aneuploidy defined in contrast to euploidy?

- A) Aneuploidy is a change in chromosome number by addition/deletion of a chromosome or part of a chromosome
- B) Aneuploidy is the gain of one or more complete sets of chromosomes
- C) Aneuploidy refers exclusively to base pair substitutions
- D) Aneuploidy is the complete loss of all nuclear DNA

2. Which of the following is classified as a variation in euploidy?

- A) Monosomy
- B) Trisomy
- C) Triploid
- D) Nullisomy

3. Which of the following is listed in Lesson 117 as an example of aneuploidy?

- A) Triploid
- B) Tetraploid
- C) Nullisomy
- D) Pentaploid

4. What defines euploidy according to the lesson comparison?

- A) Loss of a single sex chromosome
- B) Gain of one or more complete sets of chromosomes
- C) Single nucleotide point substitution
- D) Inversion of a gene segment by 180 degrees

Answer Key

1. A — Aneuploidy involves the addition or deletion of individual chromosomes or chromosome parts, altering chromosome number.
2. C — Triploid represents euploidy (gain of a complete set), whereas monosomy, trisomy, and nullisomy are forms of aneuploidy.
3. C — Nullisomy, monosomy, trisomy, and tetrasomy are listed as forms of aneuploidy.
4. B — Euploidy is defined as gaining one or more complete sets of chromosomes (such as triploidy or tetraploidy).

Lesson 118: Chromosomal deletions**1. Deletions of larger chromosomal portions are generally described as:**

- A) Harmless neutral variations
- B) Resulting in substantial loss of essential genes with severe phenotypic consequences
- C) Beneficial
- D) Unnoticeable

2. What percentage of chromosomal deletions are true de novo deletions vs. resulting from parental rearrangements?

- A) 50/50
- B) 85-90% true de novo deletions; 10-15% from parental rearrangements
- C) 100% inherited
- D) 0% de novo

3. How is a "terminal deletion" distinguished from an "interstitial deletion"?

- A) Terminal deletions lose a chromosome tip following one break, whereas interstitial deletions lose an internal segment after two breaks
- B) They are identical
- C) Terminal deletions are larger
- D) Interstitial deletions affect telomeres

4. Cri-du-chat syndrome is caused by a terminal deletion on which chromosome?

- A) Short arm of chromosome 5 (5p)
- B) Chromosome 21
- C) X chromosome
- D) Chromosome 1

Answer Key

1. B — Large chromosomal deletions result in substantial loss of essential genes and are phenotypically severe.
2. B — The handout specifies that 85-90% are true deletions, while 10-15% stem from parental rearrangements.
3. A — Terminal deletions lose a chromosome tip following one break, whereas interstitial deletions lose an internal segment.
4. A — Cri-du-chat syndrome is clinically associated with a terminal deletion of the short arm of chromosome 5.

Lesson 119: Chromosomal duplication

1. What role does gene duplication play during molecular evolution?

- A) It serves as a major mechanism through which new genetic material is generated
- B) It eliminates all non-coding intronic sequences
- C) It converts linear chromosomes into circular plasmids
- D) It prevents meiotic crossing over completely

2. Unequal crossing-over during meiosis between misaligned homologous chromosomes is termed:

- A) Ectopic homologous recombination
- B) Replication slippage
- C) Position-independent mapping
- D) Pericentric inversion

3. What error during DNA replication produces duplications of short genetic sequences when DNA polymerase copies DNA?

- A) Replication slippage
- B) Feulgen hydrolysis
- C) Colchicine binding
- D) Heterochromatin condensation

4. What can happen to tandem gene duplications over evolutionary time?

- A) They can mutate to take on altered roles or become inactive pseudogenes
- B) They always cause immediate embryonic lethality
- C) They convert into histone octamers
- D) They dissolve during cytokinesis

Answer Key

1. A — Chromosomal/gene duplication provides a primary mechanism for generating new genetic material in molecular evolution.
2. A — Duplications commonly arise from ectopic homologous recombination (unequal crossing-over) between misaligned homologous chromosomes.
3. A — Replication slippage is a DNA replication error that duplicates short nucleotide sequences.
4. A — Tandem duplicate copies can mutate to adopt new cellular functions or degenerate into pseudogenes.

Lesson 120: Chromosomal Inversions

1. What structural rearrangement occurs during a chromosomal inversion?

- A) A chromosome segment breaks in two places, rotates 180 degrees, and rejoins
- B) A segment is deleted
- C) A segment is duplicated
- D) Two chromosomes fuse

2. What is the key structural difference between a pericentric inversion and a paracentric inversion?

- A) Pericentric inversions span across the centromere, whereas paracentric inversions do not include the centromere
- B) They are identical
- C) Paracentric includes the centromere
- D) Neither involves breaks

3. Does a balanced chromosomal inversion involve a net loss or gain of genetic material?

- A) Yes, substantial loss
- B) No, inversions reorient genetic sequence without adding or subtracting total genomic content
- C) Yes, gain
- D) Always lethal loss

4. While many inversions have no direct phenotypic consequences for the carrier, they can cause what reproductive issue?

- A) Heightened risks of producing unbalanced gametes and offspring with abnormalities
- B) Nothing at all
- C) Increased fertility
- D) Twinning

Answer Key

1. A — Inversions occur when a chromosome segment breaks in two places, rotates 180 degrees, and rejoins.
2. A — Pericentric inversions span across the centromere, whereas paracentric inversions exclude it.
3. B — Inversions reorient genetic sequence without adding or subtracting total genomic content.
4. A — Carriers of balanced inversions face heightened risks of producing unbalanced gametes.

Lesson 121: Ring chromosome

1. How is a ring chromosome formed structurally?

- A) When breakage at chromosome ends allows the terminal arms to fuse together forming a ring
- B) By duplication
- C) By deletion of the centromere
- D) By translocation

2. How are ring chromosomes designated in human and Drosophila genetic nomenclature?

- A) Lowercase 'r' in human cytogenetics and uppercase 'R' in Drosophila
- B) 'R' in both
- C) 'r' in both
- D) No designation

3. What usually must happen to the ends of a chromosome to enable ring chromosome formation?

- A) Loss of terminal caps (telomeres) on both arms, producing reactive ends that fuse

- B) Nothing
- C) Centromere duplication
- D) Arm lengthening

4. What percentage of Turner syndrome cases are caused by a ring X chromosome?

- A) 5%
- B) 20%
- C) 50%
- D) 100%

Answer Key

1. A — Ring chromosomes occur when breakage at chromosome ends allows the terminal arms to fuse.
2. A — Cytogenetic notation uses lowercase 'r' for human ring chromosomes and uppercase 'R' in Drosophila.
3. A — Standard ring formation requires loss of terminal caps on both arms, producing reactive ends.
4. B — Approximately 20% of Turner syndrome cases are attributed to a ring X chromosome.

Lesson 122: Mutation

1. How does the definition of "mutation" vary across different scientific disciplines?

- A) In general genetics it is interpreted broadly as any sequence change; in clinical contexts it often implies a disease-causing change
- B) It is identical everywhere
- C) Only viruses mutate
- D) Mutations don't exist

2. Why do some geneticists prefer neutral terms like "sequence variant" or "allelic variant" instead of "mutation"?

- A) Neutral terminology avoids presuming disease causality
- B) They are shorter
- C) They sound better
- D) No reason

3. What are the potential phenotypic consequences of a gene mutation?

- A) Mutations can be harmful, beneficial, or neutral depending on location and functional impact
- B) Always lethal
- C) Always beneficial
- D) No consequences

4. How do mutations in somatic cells compare to mutations in germline cells regarding inheritance?

- A) Somatic mutations remain confined to body tissues of the host organism, whereas germline mutations are heritable
- B) Both are inherited equally
- C) Somatic mutations are inherited
- D) Germline mutations disappear

Answer Key

1. A — "Mutation" is interpreted broadly as any sequence change in general genetics, or as disease-causing in clinical usage.

2. A — Neutral terminology ("sequence/allelic variant") avoids presuming disease causality.
3. A — Depending on location and functional impact, mutations may exert harmful, beneficial, or neutral effects.
4. A — Somatic mutations remain confined to body tissues of the host organism, whereas germline mutations can be transmitted to offspring.

Lesson 123: Types of mutation

1. Which of the following is categorized as a type of point mutation involving base pair substitutions in Lesson 123?

- A) Translocation
- B) Missense mutation
- C) Polyploidy
- D) Monosomy

2. Under which major category of mutations are frameshift and triplet repeat mutations classified?

- A) Point mutations (base pair insertions/deletions)
- B) Variations in chromosome number
- C) Structural translocations
- D) Post-zygotic mosaicism

3. Variations in chromosome number are sub-classified into which two main categories?

- A) Silent and missense
- B) Aneuploidy and Euploidy
- C) Transitions and Transversions
- D) Forward and reverse

4. Which of the following structural chromosomal changes is listed under variations in chromosome structure in Lesson 123?

- A) Nonsense mutation
- B) Inversion
- C) Hypoploidy
- D) Triploidy

Answer Key

1. B — Silent, missense, and nonsense mutations are categorized as base pair substitutions under point mutations.
2. A — Frameshift and triplet repeat mutations are classified as base pair insertions/deletions under point mutations.
3. B — Variations in chromosome number are categorized into Aneuploidy and Euploidy.
4. B — Deletions, translocations, duplications, and inversions represent variations in chromosome structure.

Lesson 124: Somatic and Germline mutation

1. Where do somatic mutations arise, and how are they propagated to daughter cells within an organism?

- A) In non-germline body cells, propagated by mitosis

- B) In gametes
- C) They are not propagated
- D) By meiosis

2. Where do germline mutations occur, and what is their primary evolutionary significance?

- A) In gamete-forming lineages, enabling the altered allele to be inherited by offspring
- B) In skin cells
- C) They have no significance
- D) In dead cells

3. In multicellular organisms, when scientists use the unadorned term "mutation" without qualification, what does it conventionally denote?

- A) Inherited germline variants
- B) Somatic variants only
- C) Protein changes
- D) Nothing specific

4. Which of the following is listed in Lesson 124 as a method used to classify mutations?

- A) On the basis of molecular defect mechanism, phenotypic effect, and location
- B) By color
- C) By size of organism
- D) By age

Answer Key

- 1. A — Somatic mutations take place in non-germline body cells and divide mitotically among daughter cells.
- 2. A — Germline mutations occur in reproductive lineages, enabling the altered allele to be passed to offspring.
- 3. A — Standard genetic convention uses "mutation" to denote inherited germline variants.
- 4. A — Mutations are categorized systematically by molecular defect mechanism, phenotypic effect, and other criteria.

Lesson 125: Mutation - Forward, Reverse and Neutral

1. How is the "wild type" phenotype defined in Lesson 125?

- A) The rarest phenotype observed in a laboratory culture
- B) The most common phenotype in natural populations of an organism
- C) Any phenotype caused by a lethal chromosomal deletion
- D) A phenotype expressed only at elevated environmental temperatures

2. What is a "forward mutation"?

- A) A mutation that alters the wild type phenotype into a mutant phenotype
- B) A mutation that reverts a mutant phenotype back to wild type
- C) A mutation that changes DNA without altering any amino acid
- D) A mutation occurring exclusively in mitochondrial tRNA genes

3. A mutation that changes a mutant phenotype back into the original wild type phenotype is called a:

- A) Forward mutation
- B) Reverse mutation
- C) Conditional mutation

D) Dominant negative mutation

4. What defines a "neutral mutation" at the protein level?

- A) It converts an amino acid codon into a translation stop codon
- B) It alters the amino acid sequence of the protein without changing its function
- C) It causes complete loss of gene product
- D) It creates an entirely novel enzymatic activity

Answer Key

1. B — The wild type phenotype is defined as the most common phenotype in natural populations of the organism.
2. A — A forward mutation alters the wild type phenotype into a mutant phenotype.
3. B — A reverse mutation changes a mutant phenotype back into the wild type phenotype.
4. B — A neutral mutation alters the amino acid sequence but does not change protein function because the replaced amino acid is chemically similar or has little influence.

Lesson 126: Loss of function, Gain of function

1. How is a loss of function mutation defined, and what is its typical inheritance pattern?

- A) It produces a new trait; usually dominant
- B) It causes complete or partial loss of normal protein function; typically recessive
- C) It has no effect
- D) It only affects RNA

2. How is a gain of function mutation characterized according to Lesson 126?

- A) It is frequently recessive
- B) It produces a novel trait or expresses the gene product in inappropriate tissues/times; often dominant
- C) It deletes the gene
- D) It silences the gene

3. A temperature-sensitive mutation that causes cell death at high temperatures but has no deleterious effect at low temperatures is an example of:

- A) A lethal mutation
- B) A conditional mutation (expressed only under certain conditions)
- C) A silent mutation
- D) A nonsense mutation

4. Which statement regarding lethal mutations is correct according to the source handout?

- A) Death must occur at birth
- B) Lethal mutations significantly reduce longevity and cause death, which may take many forms/timing
- C) Lethal mutations are always recessive
- D) Lethal mutations don't exist

Answer Key

1. B — Loss of function mutations cause a complete or partial loss of normal protein function and are typically recessive.
2. B — Gain of function mutations produce a novel trait or express it in inappropriate tissues/times.

3. B — Conditional mutations, such as temperature-sensitive variants, are expressed only under specific conditions.
4. B — Lethal mutations significantly reduce longevity and cause death, which may take many forms.

Lesson 127: Spontaneous and Induced mutation

1. How are spontaneous mutations defined in Lesson 127?

- A) Mutations that result from natural changes in DNA
- B) Mutations caused exclusively by industrial chemicals
- C) Mutations induced by UV and X-ray radiation
- D) Mutations created artificially in recombinant plasmids

2. Induced mutations result from DNA changes caused by:

- A) Environmental factors, chemicals, and radiation
- B) Natural DNA replication proofreading only
- C) Normal histone acetylation
- D) Complete absence of mutagens

3. Which of the following is an origin of spontaneous mutations mentioned in the lesson?

- A) Replication errors and spontaneous DNA damage
- B) Exposure to medical X-ray radiation
- C) Treatment with alkylating chemical mutagens
- D) Colchicine application

4. What distinguishes an induced mutation from a spontaneous mutation?

- A) Induced mutations are caused by external environmental, chemical, or radiation agents
- B) Induced mutations never alter the nucleotide sequence
- C) Spontaneous mutations only occur in artificial cell cultures
- D) Spontaneous mutations affect only sex chromosomes

Answer Key

1. A — Spontaneous mutations arise naturally from endogenous cellular processes and natural changes in DNA.
2. A — Induced mutations stem from external mutagens including environmental agents, chemicals, and radiation.
3. A — Spontaneous mutations originate naturally through replication errors or spontaneous DNA damage.
4. A — Induced mutations are triggered by external environmental, chemical, or radiation exposure rather than natural internal processes.

Lesson 128: Suppressor mutation

1. How does a suppressor mutation differ fundamentally from a reverse mutation?

- A) Suppressor mutations are identical to reverse mutations
- B) A suppressor mutation occurs at a site distinct from the original mutation, whereas a reverse mutation restores the original site
- C) There is no difference
- D) Suppressor mutations delete genes

2. An organism carrying both an original background mutation and a suppressor mutation is a double mutant that:

- A) Shows no change
- B) Exhibits the phenotype (partially or fully restored toward wild-type)
- C) Always dies
- D) Becomes haploid

3. What defines intragenic suppression?

- A) A suppressor mutation occurring in a completely different gene
- B) Suppressor mutations that occur within the same gene as the original mutation
- C) Suppression between species
- D) Deletion of the gene

4. Intergenic (extragenic) suppression relieves the effect of a mutation by introducing a second mutation:

- A) In the same codon
- B) Situated on a different gene
- C) That deletes the chromosome
- D) In mitochondria only

Answer Key

1. B — A suppressor mutation occurs at a site distinct from the original mutation, whereas a reverse mutation corrects the original site.
2. B — An organism carrying a suppressor mutation is a double mutant that exhibits the phenotype restored toward normal.
3. B — Intragenic suppression results from suppressor mutations that occur within the same gene.
4. B — Intergenic suppression involves a second mutation situated on a different gene.

Lesson 129: Point mutations

1. Point mutations are defined as DNA alterations that:

- A) Delete entire chromosome arms
- B) Affect single sites on DNA through base substitutions or small additions/deletions
- C) Duplicate whole genomes
- D) Invert chromosomes

2. What type of base substitution occurs when a purine replaces another purine (e.g., Adenine to Guanine)?

- A) Transversion
- B) Transition
- C) Frameshift
- D) Deletion

3. A base substitution where a purine is replaced by a pyrimidine (or vice versa) is termed a:

- A) Transition
- B) Transversion
- C) Silent mutation
- D) Duplication

4. Which of the following is listed in Lesson 129 as a potential result of point mutations?

- A) Silent, missense, or nonsense mutations at the protein level
- B) Whole chromosome loss
- C) Polyploidy
- D) Ring chromosomes

Answer Key

1. B — Point mutations affect single sites on DNA through base substitutions or small additions/deletions.
2. B — A transition substitution occurs when a purine substitutes for a purine or a pyrimidine for a pyrimidine.
3. B — Transversion refers to a base substitution where a purine substitutes for a pyrimidine or vice versa.
4. A — Point mutations can yield silent, missense, or nonsense mutations at the protein level.

Lesson 130: Silent Mutation**1. How does a silent mutation affect an organism's phenotype?**

- A) It causes severe disease
- B) It has no effect on the protein or phenotype
- C) It truncates the protein
- D) It duplicates the gene

2. Which characteristic of the genetic code explains why many base substitutions result in silent mutations?

- A) Universality
- B) Degeneracy (multiple codons encode the same amino acid)
- C) Non-overlapping nature
- D) Triplet nature

3. In a codon, changes at which nucleotide position are most likely to result in a silent mutation?

- A) First base
- B) Second base
- C) Third base (wobble position)
- D) None

4. What happens at the protein level during a silent mutation?

- A) A stop codon is created prematurely
- B) The DNA base sequence changes while the amino acid sequence remains unchanged
- C) The protein is truncated
- D) The reading frame shifts

Answer Key

1. B — Silent mutations affect the DNA sequence but have no effect on the protein or phenotype.
2. B — Degeneracy of the genetic code allows multiple different codons to produce the same amino acid.
3. C — Third base changes in a codon often have no effect on the translated amino acid sequence.
4. B — Silent mutations alter the underlying DNA base sequence while leaving the protein unchanged.

Lesson 131: Missense mutation

1. How does a missense mutation affect protein synthesis?

- A) It substitutes one amino acid for another in the polypeptide chain
- B) It converts an amino acid codon into a premature stop codon
- C) It deletes three consecutive triplet base pairs
- D) It leaves the amino acid sequence completely unchanged

2. Why do some missense mutations have severe functional consequences while others have minimal effect?

- A) Because all missense mutations destroy the nuclear envelope
- B) It depends on the location in the protein structure and the chemical difference in the substituted amino acid
- C) Because missense mutations only occur in non-coding UTRs
- D) Because missense mutations always delete the centromere

3. In the example provided in Lesson 131, which amino acid substitution illustrates a missense mutation?

- A) Histidine substituted in place of Arginine
- B) Glutamate substituted in place of a stop codon
- C) Lysine substituted in place of a poly-A tail
- D) Cysteine substituted in place of a promoter

4. Does a missense mutation always alter the functional activity of the resulting protein?

- A) Yes, it always renders the protein completely non-functional
- B) No, it may change the function of the protein or it may not
- C) Yes, it always increases protein activity ten-fold
- D) No, because missense mutations never change the amino acid sequence

Answer Key

- 1. A — Missense mutations substitute one amino acid for another in the resulting protein product.
- 2. B — The phenotypic effect depends on where the substitution occurs in the protein structure and the magnitude of the amino acid change.
- 3. A — The lesson illustrates a missense mutation with Histidine substituted in for Arginine.
- 4. B — A missense mutation produces a change in the amino acid sequence, which may change protein function or may not.

Lesson 132: Nonsense mutation**1. A nonsense mutation is a point mutation that:**

- A) Replaces one amino acid with a chemically similar one
- B) Converts an amino acid-coding codon into a translation stop codon
- C) Has no effect
- D) Duplicates the gene

2. What is the primary structural effect of a nonsense mutation on the synthesized protein?

- A) It lengthens the protein
- B) Premature termination, shortening the protein (truncation)
- C) It adds amino acids
- D) No structural effect

3. Why might some nonsense mutations have relatively little functional impact on a protein?

- A) Because all truncations are harmless
- B) Truncations near the protein ends may have minimal impact if those regions are not functionally critical
- C) Because the protein is repaired
- D) Because of splicing

4. What is the most common functional outcome when a nonsense mutation occurs in a crucial coding region?

- A) Enhanced function
- B) A completely non-functional truncated protein
- C) No change
- D) Gene duplication

Answer Key

1. B — Nonsense mutations convert an amino acid-coding codon into a translation stop codon.
2. B — Converting a coding codon into a stop codon causes premature termination, shortening the protein.
3. B — Truncations near the protein ends may have minimal impact if those regions are not functionally essential.
4. B — Nonsense mutations frequently yield completely non-functional truncated proteins.

Lesson 133: Additions mutation**1. An addition mutation (insertion) involving non-triplet base counts typically results in what type of mutation?**

- A) Silent mutation
- B) Frameshift mutation
- C) Missense only
- D) No effect

2. How are addition mutations alternatively named in genetic literature according to Lesson 133?

- A) Deletions
- B) Insertions
- C) Inversions
- D) Duplications

3. What is the typical consequence of an addition/insertion frameshift mutation on protein function?

- A) Enhanced function
- B) Disruption of downstream amino acid sequences, often causing severe loss of function
- C) No consequence
- D) Increased protein length only

4. What happens to the translational reading frame when a single nucleotide is inserted into a coding sequence?

- A) It remains unchanged
- B) The reading frame shifts for all subsequent codons downstream
- C) It shifts only one codon

D) It reverses

Answer Key

1. B — Addition mutations alter the triplet reading alignment, resulting in a frameshift.
2. B — Lesson 133 specifies that addition mutations are also called insertions.
3. B — Addition mutations disrupt downstream amino acid sequences, often causing severe functional defects.
4. B — Inserting nucleotides shifts the reading frame for all subsequent codons downstream.

Lesson 134: Inversion mutation

1. According to Lesson 134, inversion mutations at the gene level affect:

- A) Entire chromosomes spanning millions of bases
- B) Only a small part of the gene (a localized segment)
- C) Whole genomes
- D) Mitochondria only

2. In the nucleotide sequence example provided in Lesson 134 (GGTCCTCTCAGCCA), how does the amino acid change after inversion?

- A) Reversing the sequence segment alters the second codon to translate Glycine instead of the original amino acid
- B) No amino acid change
- C) The protein doubles
- D) A stop codon appears at the start

3. How does a small intragenic inversion alter the local nucleotide sequence?

- A) By deleting a triplet
- B) By reversing the 180-degree orientation of a localized sequence segment
- C) By duplicating it
- D) It doesn't alter it

4. What distinguishes a small gene inversion from a large chromosomal inversion (Lesson 120 vs. Lesson 134)?

- A) Small gene inversions affect localized base sequences within a gene, whereas chromosomal inversions span large segments including the centromere region
- B) They are identical
- C) Small inversions are always lethal
- D) Large inversions affect single bases

Answer Key

1. B — Lesson 134 states that gene-level inversion mutations affect only a small part of the gene.
2. A — Reversing the sequence segment alters the second codon to translate Glycine instead of the original.
3. B — Inversion mutations reverse the 180-degree orientation of a localized sequence segment.
4. A — Small gene inversions affect localized base sequences within a gene, whereas chromosomal inversions affect large segments.

Lesson 135: Deletions Mutation

1. What is the primary effect of a single-nucleotide deletion on the downstream amino acid sequence?

- A) No effect
- B) All downstream amino acids from the mutation site change (frameshift)
- C) Only one amino acid changes
- D) The protein lengthens

2. Proteins produced following a frameshift deletion are generally:

- A) Hyper-functional
- B) Non-functional
- C) Longer
- D) Unaffected

3. In standard protein mutation nomenclature, how is a deletion of amino acid Lysine at position 2 designated?

- A) p.Lys2ins
- B) p.Lys2del
- C) p.Lys2
- D) c.2del

4. How is a deletion spanning from Cysteine-28 to Methionine-30 formatted according to Lesson 135 nomenclature?

- A) p.Cys28_Met30del
- B) p.Cys28del_Met30del
- C) c.28_30del
- D) p.del28-30

Answer Key

1. B — Deletion frameshifts cause all downstream amino acids from the mutation site to differ.
2. B — Proteins resulting from frameshift deletions are generally non-functional.
3. B — Single amino acid deletions are designated by "del" after the amino acid and position (e.g., p.Lys2del).
4. B — A range of deleted amino acids is indicated by separating the first and last residues (as formatted in the lesson).

Lesson 136: How mutations alter gene function

1. Which functional class of mutant allele results in a complete absence of gene product?

- A) Hypomorph
- B) Null allele (amorph / loss of function)
- C) Hypermorph
- D) Neomorph

2. What is a hypomorphic mutation?

- A) A mutation producing a completely novel activity
- B) A mutation causing decreased (partial loss of) gene product activity
- C) A mutation with increased activity
- D) A silent mutation

3. An allele whose product actively antagonizes or interferes with the normal wild-type protein product is called:

- A) Recessive
- B) Dominant negative (antimorph)
- C) Silent
- D) Neutral

4. A mutation that imparts an entirely novel biochemical activity to a gene product is classified as:

- A) Hypomorphic
- B) Neomorphic (gain of function)
- C) Amorphic
- D) Silent

Answer Key

1. B — A null allele (loss of function) produces no gene product whatsoever.
2. B — Hypomorphic mutations result in decreased gene product activity.
3. B — Dominant negative mutations produce a gene product that antagonizes the normal wild-type product.
4. B — Neomorphic mutations bestow a novel functional activity on the gene product.

Lesson 137: Mutation nomenclature

1. According to HGVS mutation nomenclature standards, at what most basic level should all variants be described?

- A) Protein level
- B) DNA level
- C) RNA level
- D) Chromosome level

2. What prefix letter indicates that a variant description uses a coding DNA reference sequence?

- A) g.
- B) c.
- C) m.
- D) r.

3. Which prefix letter is assigned to mitochondrial DNA reference sequences?

- A) c.
- B) m.
- C) r.
- D) g.

4. Official gene symbols used in standard mutation nomenclature must be approved by which authority?

- A) WHO
- B) HGNC (HUGO Gene Nomenclature Committee, www.genenames.org)
- C) FDA
- D) UNESCO

Answer Key

1. B — All variants should be described at the most basic level, namely the DNA level.
2. B — The prefix "c." designates a coding DNA reference sequence (e.g., c.76A>T).
3. B — The prefix "m." specifies a mitochondrial sequence (e.g., m.8993T>C).
4. B — Only official HGNC gene symbols (www.genenames.org) should be used when describing variants.

Lesson 138: Salient feature of Mutation nomenclature

1. In coding DNA mutation nomenclature, which specific nucleotide is designated as nucleotide 1?

- A) The first nucleotide of the chromosome
- B) The 'A' of the ATG translation initiation codon
- C) The promoter TATA box
- D) The stop codon

2. Is there a nucleotide designated as "0" (zero) in standard mutation nomenclature?

- A) Yes, it denotes the promoter
- B) No, there is no nucleotide 0
- C) Yes, it denotes the start codon
- D) Only in mitochondria

3. How are nucleotides located in the 5' untranslated region (upstream of the ATG start codon) numbered?

- A) With positive numbers
- B) Using negative numbers (-1, -2, ...) counting upstream from the ATG
- C) With asterisks
- D) They are not numbered

4. How are nucleotides situated in the 3' untranslated region (downstream of the translation stop codon) designated?

- A) With an asterisk (*1, *2, ...) counting downstream from the stop codon
- B) With negative numbers
- C) With zero
- D) They are not numbered

Answer Key

1. B — Nucleotide 1 is defined specifically as the 'A' of the ATG-translation initiation codon.
2. B — Salient nomenclature rules explicitly mandate that there is no nucleotide 0.
3. B — Nucleotides 5' of the ATG start codon are designated using negative numbers (-1, -2, ...).
4. A — Nucleotides 3' of the translation stop codon are designated with an asterisk (*1, *2, ...).

Lesson 139: Mutation at DNA, RNA and protein level

1. How are nucleotide letters written when describing mutations at the DNA level versus the RNA level?

- A) DNA-level mutations are written in capitals, whereas RNA-level descriptions use lower-case letters
- B) Both use capitals

- C) Both use lower-case
- D) DNA uses numbers only

2. A substitution at the RNA level changing Adenine to Uracil at position 76 is correctly formatted as:

- A) c.76A>U
- B) r.76a>u
- C) p.76A>U
- D) m.76A>U

3. How does a protein-level mutation description begin according to HGVS rules?

- A) Starting with a number
- B) With 'p.' followed by capital letters denoting the amino acid
- C) With 'c.'
- D) With the gene name

4. Which of the following represents a properly formatted protein substitution changing Lysine at position 2 to Asparagine?

- A) c.2A>T
- B) p.Lys2Asn
- C) r.2a>u
- D) g.Lys2Asn

Answer Key

1. A — DNA-level mutations are written in capitals, whereas RNA-level descriptions use lower-case letters.
2. B — RNA-level variants use the 'r.' prefix with lower-case nucleotide letters (e.g., r.76a>u).
3. B — Protein-level variants begin with 'p.' followed by capital letters denoting the amino acid.
4. B — Protein substitutions start with 'p.' followed by the three-letter amino acid codes (e.g., p.Lys2Asn).

Lesson 140: Mutation in intronic sequence

1. How is a mutation located at the beginning of an intron formatted in HGVS coding DNA nomenclature?

- A) Using the last nucleotide of the preceding exon, a plus sign, and the intronic offset (e.g., c.76+1)
- B) Using only the exon number
- C) With a minus sign
- D) It cannot be described

2. How is a mutation located near the end of an intron designated?

- A) The last nucleotide of the preceding exon with a plus
- B) The first nucleotide of the following exon with a minus sign and offset (e.g., c.77-1)
- C) With an asterisk
- D) With the intron number only

3. In the middle of an intron with an uneven number of nucleotides, how is the central nucleotide described?

- A) By the exon position
- B) As the last one described from the 5' end (or per the lesson's convention for the central nucleotide)
- C) It cannot be described

D) With zero

4. What symbol separates the preceding exon nucleotide position from the relative intronic position?

- A) Minus sign
- B) A plus sign (+) indicates a position counting downstream into the intron from the exon
- C) Underscore
- D) Colon

Answer Key

1. A — At the beginning of an intron, notation uses the last nucleotide of the preceding exon plus the intronic offset.
2. B — At the end of an intron, notation uses the first nucleotide of the following exon with a minus offset.
3. B — In an intron with an uneven base count, the central nucleotide is the last one described from the 5' side.
4. B — A plus sign (+) indicates a position counting downstream into the intron from the exon.

Lesson 141: Specific changes in sequence

1. In HGVS DNA mutation nomenclature, what symbol is explicitly used to indicate a base substitution?

- A) Underscore (_)
- B) Greater-than sign (>)
- C) Hyphen (-)
- D) Plus (+)

2. How are duplicating insertions described according to standard HGVS nomenclature rules?

- A) As insertions (ins)
- B) As duplications (dup)
- C) As inversions (inv)
- D) As substitutions

3. Which symbol is used to specify a range of affected residues between the first and last position?

- A) An underscore (_)
- B) A hyphen (-)
- C) A colon (:)
- D) A semicolon (;)

4. What abbreviation is used to denote an inversion of a sequence segment at the DNA level?

- A) del
- B) ins
- C) dup
- D) inv

Answer Key

1. B — The ">" symbol explicitly indicates a nucleotide substitution at the DNA level (e.g., c.76A>T).
2. B — Duplicating insertions are described specifically as duplications (dup) rather than insertions (ins).

3. A — An underscore (_) separates the first and last affected residues to indicate a sequence range (e.g., c.76_78delACT).
4. D — The abbreviation "inv" designates a sequence inversion (e.g., c.76_83inv).

Lesson 142: Mutations - Summary

1. In the sentence-analogy summary of mutations, how is a missense mutation represented relative to the original sentence?

- A) A word is deleted
- B) A single letter/amino acid is altered, changing "RED" to "RAD"
- C) The sentence is cut short
- D) Words are repeated

2. Which sentence analogy represents a nonsense mutation in the lesson summary?

- A) "THE BIG RAD DOG"
- B) A premature stop truncating the sentence (e.g., "THE BIG ...")
- C) "THE BGR EDD OGR"
- D) "THE BIG BIG BIG BIG BIG RED DOG RAN OUT"

3. How is a 3-bp in-frame deletion illustrated in the sentence-analogy summary?

- A) "THE BGR EDD OGR"
- B) Deleting 3 base pairs (one full word) removes "RED" without shifting the reading frame
- C) The sentence is cut in half
- D) Letters are swapped

4. The sentence "THE BIG BIG BIG BIG BIG RED DOG RAN OUT" serves as an analogy for which type of mutation?

- A) Missense
- B) Triplet repeat expansion mutation
- C) Nonsense
- D) Frameshift deletion

Answer Key

- 1. B — A missense mutation alters a single letter/amino acid, changing "RED" to "RAD".
- 2. B — A nonsense mutation introduces a premature stop, truncating the sentence.
- 3. B — Deleting 3 base pairs (one full word) removes "RED" without shifting the reading frame.
- 4. B — Repeated copies of the word "BIG" represent a triplet repeat expansion mutation.

Lesson 143: Genetic Disorders

1. What are the five main classes of genetic disorders according to Lesson 143?

- A) Multifactorial, single gene, chromosomal, mitochondrial, and somatic-mutation (cancer) disorders
- B) Only single gene disorders
- C) Only chromosomal disorders
- D) Infectious and genetic

2. How are "fully penetrant" genetic disorders characterized?

- A) Environmental factors exclusively determine expression

- B) Other background genes and environmental conditions have little influence; the genotype almost always produces the phenotype
- C) They never manifest
- D) They affect only one sex

3. In "low-penetrance" genetic disorders, what determines phenotypic expression?

- A) Genes play a small portion of risk along with environmental factors
- B) Genes play no role
- C) Only environment
- D) Only age

4. Cancer is listed under which category of genetic disorders in Lesson 143?

- A) Mitochondrial disorders
- B) Genetic disorders resulting from somatic mutations
- C) Single gene disorders
- D) Chromosomal disorders

Answer Key

1. A — Lesson 143 classifies genetic disorders into multifactorial, single gene, chromosomal, mitochondrial, and somatic mutation categories.
2. B — In fully penetrant disorders, other background genes and environmental conditions minimally modify expression.
3. A — Low-penetrance disorders involve genes contributing a small portion of risk along with environment.
4. B — Cancers are classified as genetic disorders resulting from somatic mutations.

Lesson 144: Inheritance patterns of genetic disorders

1. In autosomal dominant genetic disorders, what is the genotype of affected individuals?

- A) Heterozygote (one allele is mutated, while the other is normal)
- B) Homozygous recessive
- C) Hemizygous with no normal allele
- D) Nullizygous

2. What is the genetic requirement for expressing an autosomal recessive disorder?

- A) Both alleles of the gene must be mutated (homozygous)
- B) Only one allele needs to be mutated
- C) The mutation must reside on the Y chromosome
- D) Environmental factors must be 100% responsible

3. How are multifactorial genetic disorders characterized in Lesson 144?

- A) Both genes and the environment contribute to disease manifestation
- B) They are caused strictly by a single point mutation in a single gene
- C) They affect only mitochondrial DNA
- D) They are passed exclusively from father to son

4. How do chromosomal disorders affect an organism compared to single-gene disorders?

- A) They involve thousands of genes and affect multiple organ systems at multiple stages of gestation
- B) They affect only hair color and eye color
- C) They are always single-gene recessive traits

D) They have no impact on embryonic development

Answer Key

1. A — Affected individuals in autosomal dominant inheritance are heterozygotes carrying one mutated allele and one normal allele.
2. A — Autosomal recessive disorders manifest when an individual is homozygous for two mutated alleles.
3. A — Multifactorial disorders arise from the combined contributions of multiple genes and environmental factors.
4. A — Chromosomal disorders involve large numbers of genes, impacting multiple organ systems across gestational stages.

Lesson 145: Autosomal recessive inheritance / diseases

1. In autosomal recessive inheritance, an individual manifests the clinical disease phenotype when:

- A) They carry a single mutant allele
- B) They possess two mutant alleles (homozygous)
- C) They are hemizygous for the X chromosome
- D) They have a chromosomal deletion

2. Parents who are carriers for an autosomal recessive condition possess:

- A) Two copies of the altered allele
- B) One normal allele and one altered allele (heterozygous)
- C) No normal alleles at all
- D) Three copies of the gene

3. Which of the following phenotypic observations in a pedigree suggests an autosomal mode of inheritance?

- A) Only males are affected
- B) Equal phenotypic proportions between affected males and females
- C) The trait skips all generations
- D) Only the eldest child is affected

4. Which of the following is listed in the handout as a common autosomal recessive disorder?

- A) Marfan syndrome
- B) Cystic fibrosis
- C) Huntington disease
- D) Achondroplasia

Answer Key

1. B — Autosomal recessive diseases manifest only in individuals who possess two mutant alleles.
2. B — Heterozygous carrier parents possess one normal allele and one altered allele for the gene.
3. B — Equal phenotypic proportions between affected males and females indicate an autosomal inheritance pattern.
4. B — Cystic fibrosis is listed as a common autosomal recessive disease (alongside Bloom syndrome and others).

Lesson 146: Properties of Autosomal recessive inheritance

1. What term describes mating between individuals who share common ancestors, which increases the likelihood of autosomal recessive disease in offspring?

- A) Random mating
- B) Consanguinity
- C) Outbreeding
- D) Cloning

2. When two heterozygous carrier parents for an autosomal recessive disorder have children, what is the probability that each child will be affected?

- A) 50% (1 in 2)
- B) 25% (1 in 4)
- C) 100%
- D) 0%

3. Why does consanguinity increase the risk of offspring inheriting an autosomal recessive disease?

- A) Sharing common ancestors raises the likelihood that both parents inherited and carry the same mutant allele
- B) It causes new mutations in every generation
- C) It eliminates all dominant alleles
- D) It doubles the number of chromosomes

4. What is the expected proportion of unaffected carrier children born to two heterozygous carrier parents?

- A) 25% (1 in 4)
- B) 50% (2 in 4)
- C) 75% (3 in 4)
- D) 100%

Answer Key

1. B — Consanguinity refers to mating between parents with common ancestors, raising the risk of recessive disease.
2. B — Each sibling has a 25% (1 in 4) chance of inheriting two mutant alleles from heterozygous carrier parents.
3. A — Sharing common ancestors raises the likelihood that both parents inherited and carry the same rare mutant allele.
4. B — Crossing two carrier parents yields a 50% (2 in 4) probability of offspring being unaffected carriers.

Lesson 147: Autosomal dominant inheritance/ disorders

1. In autosomal dominant disorders, what is the relationship between the normal allele and the abnormal allele?

- A) The normal allele is dominant
- B) The abnormal allele exerts a dominant effect over the normal allele
- C) Both alleles are recessive
- D) The alleles are co-dominant

2. What proportion of children born to an affected individual (heterozygous for an autosomal dominant disorder) and an unaffected partner will inherit the disorder?

- A) 25%
- B) 50%
- C) 75%
- D) 100%

3. Which pedigree feature is a hallmark clue for identifying autosomal dominant inheritance?

- A) The trait skips generations
- B) The trait appears in every generation, with each affected person having an affected parent
- C) Only males are affected
- D) The trait appears only in twins

4. Which of the following conditions is listed as an autosomal dominant disorder in Lesson 147?

- A) Cystic fibrosis
- B) Marfan syndrome
- C) Hemophilia A
- D) Color blindness

Answer Key

1. B — In autosomal dominant conditions, the abnormal allele exerts a dominant effect over the normal allele.
2. B — An affected heterozygous parent has a 50% chance of transmitting the dominant disorder allele to each child.
3. B — Autosomal dominant traits characteristically appear in every generation, with each affected individual having an affected parent.
4. B — Marfan syndrome (along with Huntington's, Retinoblastoma, Achondroplasia, etc.) is listed as an autosomal dominant disorder.

Lesson 148: Variations in Autosomal dominant inheritance

1. What defines "incomplete or reduced penetrance" as a variation in autosomal dominant inheritance?

- A) Everyone with the genotype shows the trait
- B) Individuals with a disease-causing genotype do not always express the phenotype
- C) Only males express it
- D) It is the same as expressivity

2. Variable expressivity in autosomal dominant inheritance refers to differences in:

- A) Whether the trait appears at all
- B) Varying degrees of clinical severity among individuals with the same genotype
- C) Chromosome number
- D) Gender ratio

3. What term describes a single gene producing multiple distinct phenotypic effects?

- A) Penetrance
- B) Expressivity
- C) Pleiotropy
- D) Epistasis

4. In a typical autosomal dominant pedigree, what is the expected parental status of an affected person?

- A) Both parents normal
- B) Every affected individual typically has at least one affected parent
- C) Only the mother is affected
- D) Parents are always carriers

Answer Key

1. B — Incomplete penetrance occurs when individuals with a disease-causing genotype do not express the trait.
2. B — Variable expressivity denotes varying degrees of clinical severity among individuals with the same genotype.
3. C — Pleiotropy refers to a single gene exerting impacts on multiple distinct phenotypes.
4. B — In standard autosomal dominant inheritance, every affected individual typically has an affected parent.

Lesson 149: Dominance and Recessiveness

1. According to Lesson 149, dominance and recessiveness are fundamental properties of:

- A) Individual alleles in isolation
- B) Physical characters/traits rather than genes themselves
- C) Chromosomes as a whole
- D) Mitochondrial DNA only

2. How is a character defined as dominant in genetics?

- A) It is expressed only when homozygous
- B) It is expressed in a heterozygous individual
- C) It never appears in offspring
- D) It is always lethal

3. Why is Cystic fibrosis classified as a recessive trait?

- A) Because only homozygotes manifest the disease (two mutant alleles required)
- B) Because it is carried on the Y chromosome
- C) Because it affects only females
- D) Because it is caused by the environment

4. What term describes human males having only a single copy of gene loci present on the X and Y chromosomes?

- A) Homozygous
- B) Heterozygous
- C) Hemizygous
- D) Nullizygous

Answer Key

1. B — Dominance and recessiveness describe properties of physical characters/traits rather than the genes themselves.
2. B — A trait is defined as dominant if it manifests in a heterozygous individual.
3. A — Cystic fibrosis is recessive because two mutant alleles (homozygosity) are required for disease expression.

4. C — Males possess a single X and Y chromosome, making them hemizygous for sex-linked gene loci.

Lesson 150: X-linked recessive inheritance and disorders

1. Why are males significantly more frequently affected by X-linked recessive disorders than females?

- A) Males have two X chromosomes
- B) Being hemizygous (XY), males express X-linked recessive traits whenever their single X carries the mutant allele
- C) Females lack X chromosomes
- D) The Y chromosome protects females

2. In a pedigree showing X-linked recessive inheritance, how does transmission occur from an affected father?

- A) He passes the trait to all his sons
- B) Fathers pass their Y chromosome to sons, so affected fathers never transmit X-linked traits to sons (but pass the mutant X to all daughters)
- C) He passes the trait to 50% of children randomly
- D) He cannot have children

3. Which of the following is a general characteristic of X-linked recessive traits in family pedigrees?

- A) Traits never skip generations
- B) X-linked recessive traits characteristically skip generations and predominantly affect males
- C) Only females are ever affected
- D) Traits appear in every generation

4. Which of the following conditions is listed as an X-linked recessive disorder in Lesson 150?

- A) Achondroplasia
- B) Hemophilia A
- C) Marfan syndrome
- D) Huntington disease

Answer Key

- 1. B — Being hemizygous (XY), males express X-linked recessive traits whenever their single X chromosome carries the mutation.
- 2. B — Fathers pass their Y chromosome to sons, so affected fathers never transmit X-linked traits to sons.
- 3. B — X-linked recessive traits characteristically skip generations and predominantly affect males.
- 4. B — Hemophilia A (along with Hemophilia B, Duchenne Muscular Dystrophy, Color Blindness) is listed as an X-linked recessive disorder.

Lesson 151: X-linked dominant inheritance and disorders

1. In X-linked dominant inheritance, what is the pattern of transmission from an affected father?

- A) He passes the trait to all his sons
- B) An affected father passes his single dominant mutant X chromosome to all daughters (but none to sons)
- C) He passes the trait to no children
- D) He passes the trait only to his father

2. What is the expected inheritance outcome when an affected heterozygous female mates with an unaffected male?

- A) All children will be affected
- B) A heterozygous female ($(X^A X^a)$) has a 50% chance of transmitting the dominant mutant allele to any child, male or female
- C) No children will be affected
- D) Only sons will be affected

3. How does X-linked dominant inheritance differ from X-linked recessive inheritance regarding female expression?

- A) Females are never affected in X-linked dominant inheritance
- B) In X-linked dominant inheritance, a single mutant allele causes disease expression in females
- C) There is no difference
- D) X-linked dominant traits skip generations

4. Which of the following diseases is classified as an X-linked dominant disorder in Lesson 151?

- A) Rett syndrome
- B) Hemophilia A
- C) Cystic fibrosis
- D) Sickle cell anemia

Answer Key

1. B — An affected father passes his single dominant mutant X chromosome to all daughters.
2. B — A heterozygous female ($(X^A X^a)$) has a 50% chance of transmitting the dominant mutant allele to any child, male or female.
3. B — In X-linked dominant inheritance, a single mutant allele causes disease expression in heterozygous females.
4. A — Rett syndrome (along with Alport syndrome, Goltz syndrome, and Fragile-X) is an X-linked dominant disorder.

Lesson 152: Maternal inheritance of mitochondrial DNA

1. How is mitochondrial DNA (mtDNA) transmitted across human generations?

- A) Exclusively through sperm
- B) Exclusively through the ovum (maternal inheritance)
- C) Equally from both parents
- D) Only to sons

2. What is the expected pedigree outcome when a father carries a disease-causing mitochondrial DNA mutation?

- A) All his children will inherit it
- B) Because sperm do not contribute mitochondrial DNA to the zygote, affected fathers never transmit mtDNA diseases to offspring
- C) Only his daughters inherit it
- D) Only his sons inherit it

3. What term describes the presence of a mixture of normal and mutant mitochondrial DNA copies within a cell?

- A) Homoplasmy
- B) Heteroplasmy
- C) Aneuploidy
- D) Polyploidy

4. Which of the following conditions is listed as a mitochondrial disease in Lesson 152?

- A) Leber hereditary optic neuropathy (LHON)
- B) Cystic fibrosis
- C) Huntington disease
- D) Down syndrome

Answer Key

1. B — Mitochondrial DNA is inherited through the ovum, resulting in strict maternal inheritance.
2. B — Because sperm do not contribute mitochondrial DNA to the zygote, affected fathers never pass mtDNA mutations to children.
3. B — Heteroplasmy refers to co-existing wild-type and mutant mtDNA copies within a cell.
4. A — Leber hereditary optic neuropathy (LHON) is a classic example of a maternally inherited mitochondrial disorder.

Lesson 153: Multifactorial inheritance and disorders

1. How are multifactorial diseases defined in genetics?

- A) Disorders caused strictly by a single point mutation in a single gene
- B) Disorders arising from polygenic genetic susceptibility interacting with environmental factors
- C) Disorders caused only by the environment
- D) Disorders affecting only one organ

2. Unlike Mendelian single-gene disorders, how do risk and recurrence rates behave in multifactorial conditions?

- A) They follow strict 3:1 ratios
- B) Multifactorial conditions lack discrete Mendelian pedigree ratios (like 3:1 or 1:1)
- C) They affect 100% of offspring
- D) They never recur in families

3. Which of the following lists contains common human diseases classified as multifactorial in Lesson 153?

- A) Diabetes, Asthma, Heart diseases, and Hypertension
- B) Cystic fibrosis and Sickle cell anemia
- C) Down syndrome and Turner syndrome
- D) Hemophilia and Color blindness

4. In the spectrum of disease causality, multifactorial traits are characterized as:

- A) Completely unaffected by the environment
- B) Combining polygenic genetic factors with external environmental influences
- C) Caused by a single dominant allele
- D) Always lethal before birth

Answer Key

1. B — Multifactorial diseases arise from polygenic genetic susceptibility interacting with environmental factors.

2. B — Multifactorial conditions lack discrete Mendelian pedigree ratios (like 3:1 or 1:1).
3. A — Diabetes, Asthma, Heart diseases, and Hypertension are classic examples of multifactorial disorders.
4. B — Multifactorial traits combine polygenic genetic factors with external environmental influences.

Lesson 154: Frequency of genetic disorders

1. What is the estimated frequency of autosomal dominant disorders per 1,000 individuals according to Lesson 154?

- A) 0.5 – 1 per 1,000
- B) 2 – 3 per 1,000
- C) 3 – 9.5 per 1,000
- D) 15 – 20 per 1,000

2. According to the source handout, what is the frequency range of autosomal recessive disorders per 1,000 individuals?

- A) 0.1 – 0.5 per 1,000
- B) 2 – 2.5 per 1,000
- C) 3 – 9.5 per 1,000
- D) 10 – 15 per 1,000

3. Which category of genetic disorders has a reported frequency of 3 – 9.5 per 1,000 individuals?

- A) Autosomal recessive disorders
- B) Autosomal dominant disorders
- C) X-linked disorders
- D) Mitochondrial disorders

4. Comparing the frequencies provided in Lesson 154, how does the population frequency of autosomal dominant disorders relate to that of autosomal recessive disorders?

- A) Autosomal recessive disorders are more frequent
- B) Autosomal dominant disorders (3–9.5 per 1,000) have a higher overall population frequency than autosomal recessive disorders (2–2.5 per 1,000)
- C) Both have identical frequencies
- D) Frequencies cannot be compared

Answer Key

1. C — Lesson 154 states that the frequency of autosomal dominant disorders varies between 3 and 9.5 per 1,000 individuals.
2. B — The frequency of autosomal recessive disorders is specified as being between 2 and 2.5 per 1,000 individuals.
3. B — Autosomal dominant disorders occur at a frequency of 3 to 9.5 per 1,000 individuals.
4. B — Autosomal dominant disorders (3–9.5 per 1,000) have a higher overall population frequency than autosomal recessive disorders (2–2.5 per 1,000).

Lesson 155: Prenatal diagnosis

1. At how many weeks of gestation can amniotic fluid analysis be performed to detect genetic and chromosomal disorders?

- A) 4 – 6 weeks
- B) 8 – 10 weeks
- C) 14 – 16 weeks
- D) 30 – 32 weeks

2. Approximately how many different genetic diseases can be detected through amniotic fluid analysis?

- A) Fewer than 10
- B) About 50
- C) More than 200
- D) Exactly 1,000

3. Which prenatal diagnostic technique is specifically used to determine structural malformations of the fetus?

- A) Amniocentesis
- B) Ultrasonography of the fetus
- C) Maternal blood analysis
- D) Chorionic villus sampling

4. Which of the following is listed in Lesson 155 as a method used to observe an abnormal fetus?

- A) Karyotyping of parents
- B) Maternal blood analysis
- C) Pedigree drawing
- D) DNA fingerprinting

Answer Key

1. C — Analysis of amniotic fluid to determine genetic and chromosomal disorders is performed at 14–16 weeks of gestation.
2. C — More than 200 various genetic diseases can be easily detected via amniotic fluid analysis.
3. B — Ultrasonography of the fetus determines physical malformations of the head, internal organs, and limbs.
4. B — Maternal blood analysis is highlighted as a diagnostic approach to observe an abnormal fetus.

Lesson 156: Genetic Counseling

1. What is typically the first step taken by a genetic counselor during an evaluation?

- A) Performing amniocentesis
- B) Taking a complete family history of both prospective parents
- C) Ordering DNA sequencing
- D) Prescribing medication

2. How is gene therapy defined in Lesson 156?

- A) The microscopic counting of G-banded chromosomes
- B) A procedure involving the identification, manipulation, and reintroduction of genes to treat disease
- C) A type of prenatal diagnosis
- D) A method of pedigree analysis

3. What is the primary functional objective of gene therapy?

- A) To convert linear chromosomes into circular plasmids
- B) To replace defective genes and restore desired genetic activities
- C) To increase the number of chromosomes
- D) To prevent all mutations

4. Why is taking a complete background history of parents crucial in genetic counseling?

- A) To calculate the exact date of conception
- B) A complete family history allows counselors to evaluate inherited risks and genetic patterns
- C) To determine the parents' blood types only
- D) It is not important

Answer Key

1. B — A genetic counselor usually begins with a complete family history of both prospective parents.
2. B — Gene therapy is defined as a procedure involving the identification, manipulation, and reintroduction of genes into cells.
3. B — Gene therapy aims to replace defective genes and restore desired genetic activities.
4. B — A complete family history allows counselors to evaluate inherited risks and genetic patterns in the family.

Lesson 157: Complications in basic inheritance pattern

1. Which of the following is listed in Lesson 157 as a single-gene non-Mendelian complication?

- A) Codominance (also: incomplete dominance and overdominance)
- B) Independent assortment
- C) Segregation
- D) Dominance

2. How can multiple gene interactions complicate basic inheritance patterns?

- A) Through epistasis and polygenic interactions
- B) They cannot
- C) Only through mutation
- D) Only through environment

3. Which situation illustrates a pedigree complication where a recessive trait mimics a dominant pattern?

- A) X-linkage
- B) A common recessive trait appearing in every generation, falsely resembling dominant inheritance
- C) Mitochondrial inheritance
- D) Y-linkage

4. Sex-influenced traits and incomplete penetrance represent complications arising from interaction between:

- A) Genes and sex/environmental context
- B) Two Y chromosomes
- C) Nothing
- D) Cytoplasm only

Answer Key

1. A — Codominance, incomplete dominance, and overdominance are single-gene variations that complicate Mendelian patterns.
2. A — Multiple gene interactions produce complex inheritance through epistasis and polygenic effects.
3. B — A common recessive trait can appear in every generation, falsely resembling a dominant pattern.
4. A — Sex-influenced traits and incomplete penetrance illustrate complications stemming from gene-sex and gene-environment interactions.

Lesson 158: Drawing pedigree

1. In a standard pedigree diagram, how are generations designated?

- A) Using lowercase English letters
- B) Numbered with Roman numerals (I, II, III...), starting at the top
- C) With Arabic numerals from the bottom
- D) They are not designated

2. How are individual members within each generation numbered in a pedigree?

- A) Across the line from left to right
- B) Randomly
- C) Top to bottom
- D) They are not numbered

3. What is the primary purpose of drawing a family pedigree in genetics?

- A) To measure the exact length of chromosomes
- B) To trace and determine the mode of inheritance of traits/diseases
- C) To count family members
- D) For decoration

4. Where is generation I positioned on a pedigree chart?

- A) At the bottom of the diagram
- B) At the top (the top-most parental generation)
- C) In the middle
- D) On the left side

Answer Key

1. B — Generations are numbered with Roman numerals (I, II, III...), starting at the top.
2. A — Members within a single generation line are numbered sequentially from left to right.
3. B — Drawing pedigrees allows geneticists and counselors to trace and determine the mode of inheritance.
4. B — Generation I represents the top-most parental generation in a pedigree chart.

Lesson 159: Pedigree analysis

1. What are the two primary goals of pedigree analysis as outlined in Lesson 159?

- A) To determine cell size and shape
- B) To determine the mode of inheritance and calculate the probability of inheritance
- C) To count chromosomes under a microscope
- D) To measure DNA content

2. In pedigree diagrams, what geometric symbol is used to represent an unaffected male?

- A) A filled circle
- B) An open square
- C) A filled square
- D) An open circle

3. In a pedigree chart, what does an arrow pointing to a specific symbol indicate?

- A) Consanguineous mating
- B) The proband (propositus) — the individual through whom the family was identified
- C) A deceased individual
- D) An adopted individual

4. Which symbol represents consanguineous mating in a pedigree chart?

- A) A single horizontal line between partners
- B) A double horizontal line joining a male and female
- C) A vertical line
- D) A circle

Answer Key

1. B — Pedigree analysis aims to determine the mode of inheritance and calculate the probability of disease transmission.
2. B — Standard pedigree notation uses an open square for an unaffected male.
3. B — An arrow pointing to an individual in a pedigree identifies the proband (propositus).
4. B — Consanguineous mating is indicated by a double horizontal line joining a male and female partner.

Lesson 160: DNA polymorphism

1. How is a DNA polymorphism defined relative to a reference standard?

- A) A sequence difference present in at least 1% of the population compared to a reference standard
- B) A sequence difference compared to a reference standard
- C) A mutation causing disease in all carriers
- D) An identical sequence in all individuals

2. What proportion of DNA base pairs are identical among all humans?

- A) 50%
- B) 90%
- C) 99.9%
- D) 100%

3. What percentage of the human genomic sequence accounts for all genetically determined variability between individuals?

- A) 0.1%
- B) 1%
- C) 10%
- D) 50%

4. What size range can a DNA polymorphism span?

- A) Strictly 10 to 12 base pairs
- B) From a single nucleotide change to variations spanning large segments
- C) Exactly one chromosome

D) Only whole genes

Answer Key

1. B — DNA polymorphism is defined as a sequence difference compared to a reference standard.
2. C — In humans, 99.9% of genomic DNA base pairs are identical across individuals.
3. A — The remaining 0.1% of the genomic sequence accounts for unique differences and variability between individuals.
4. B — DNA polymorphisms range in scale from a single nucleotide change to variations spanning large segments.

Lesson 161: Types of Polymorphic DNA Sequences

1. Which of the following is classified as a single locus molecular marker in Lesson 161?

- A) DNA Fingerprinting
- B) RFLP
- C) RAPD
- D) AFLP

2. Which of the following molecular markers is categorized as a multi-locus marker?

- A) RFLP
- B) Microsatellites
- C) STS
- D) RAPD

3. Why is the study of DNA polymorphisms essential in genetics?

- A) To measure cellular water content
- B) For analyzing inheritance, mapping phenotypes, and understanding genetic diversity
- C) To count the number of cells
- D) To determine cell size

4. Which application utilizes polymorphic DNA markers in livestock and plant breeding?

- A) Feulgen hydrolysis
- B) Marker-assisted selection (MAS)
- C) Karyotyping
- D) Blood grouping

Answer Key

1. B — RFLP, Microsatellites, and STS are classified as single locus molecular markers.
2. D — RAPD, AFLP, and DNA Fingerprinting are categorized as multi-locus markers.
3. B — Studying polymorphisms is essential for analyzing inheritance, mapping phenotypes, and understanding diversity.
4. B — Polymorphic DNA markers enable marker-assisted selection (MAS) in breed development.

Lesson 162: Restriction Fragment Length Polymorphism

1. What is Restriction Fragment Length Polymorphism (RFLP)?

- A) A difference in homologous DNA sequences detected through differing restriction fragment lengths
- B) A type of mutation

- C) A sequencing method
- D) A PCR technique

2. In RFLP methodology, what is a labeled RFLP probe used for?

- A) To cut DNA at specific palindromic sites
- B) To hybridize to target digested DNA fragments on membranes after gel electrophoresis (Southern blot)
- C) To amplify DNA
- D) To sequence DNA

3. Which radioactive isotope is explicitly mentioned in Lesson 162 for labeling probes in Southern blotting?

- A) Carbon-14
- B) Phosphorus-32 (32P)
- C) Sulfur-35 (35S)
- D) Hydrogen-3 (3H)

4. Which human inherited diseases can be detected using RFLP mutation analysis according to Lesson 162?

- A) Huntington's disease, cystic fibrosis (among others listed)
- B) Common cold
- C) Flu
- D) None

Answer Key

1. A — RFLP detects homologous DNA sequence variations through differing restriction fragment lengths.
2. B — Labeled probes hybridize to target digested DNA fragments on membranes after gel electrophoresis.
3. B — Probes are labeled with Phosphorus-32 (32P) for autoradiographic detection on X-ray film.
4. A — RFLP mutation detection is explicitly listed for Huntington's disease, cystic fibrosis, and others.

Lesson 163: Variable Number Tandem Repeats

1. What is the typical size range of the repeating unit sequence in Variable Number Tandem Repeats (VNTRs)?

- A) 1 to 8 base pairs
- B) 7 (or 8) to a few tens (or >50) of base pairs
- C) 100 to 1000 base pairs
- D) Single nucleotides only

2. How are tandem repeats physically arranged on a chromosome?

- A) Clustered together and oriented in the same direction (contiguous blocks)
- B) Scattered randomly
- C) On different chromosomes
- D) Inverted alternately

3. Under basic genetic principles for VNTR data analysis, what does "Inheritance Matching" specify?

- A) A child inherits one matching VNTR allele from each parent
- B) Children inherit only from mothers

- C) No inheritance pattern
- D) Random inheritance

4. What two methods can be used to extract or amplify VNTR blocks prior to gel electrophoresis?

- A) Fluorescence microscopy
- B) Restriction digestion (RFLP) or PCR amplification
- C) Karyotyping
- D) Centrifugation only

Answer Key

1. B — VNTR repeat units range in length from 7 (or 8) to a few tens (or >50) of base pairs.
2. A — Tandem repeats are clustered contiguous blocks oriented in the same spatial direction.
3. A — Inheritance matching dictates that a child inherits one matching VNTR allele from each parent.
4. B — VNTR fragments are extracted via restriction digestion (RFLP) or amplified using PCR.

Lesson 164: Short Tandem Repeat Polymorphism

1. What is the size range of the repeating unit sequence in Short Tandem Repeats (STRs)?

- A) 1 to 8 base pairs
- B) 7 to 50 base pairs
- C) 100 to 200 base pairs
- D) 1000 base pairs

2. In STR analysis, how do the repeat region and the flanking regions compare between individuals?

- A) STR repeat counts vary between individuals, whereas flanking sequences remain constant
- B) Both vary equally
- C) Neither varies
- D) Flanking regions vary more

3. An individual carrying two STR alleles of different repeat lengths at a locus is designated as a:

- A) Homozygote
- B) Heterozygote (two distinct alleles)
- C) Hemizygote
- D) Null

4. STR markers are widely utilized in which major applications according to Lesson 164?

- A) DNA forensics and genetic linkage mapping
- B) Protein synthesis
- C) Karyotyping only
- D) Pedigree drawing

Answer Key

1. A — STRs consist of short repeating nucleotide motifs ranging from 1 to 8 base pairs.
2. A — STR repeat counts vary between individuals, whereas flanking sequences remain constant.
3. B — Differing repeat lengths on homologous chromosomes represent two distinct alleles (heterozygous).

4. A — STR markers are standard tools in DNA forensics and genetic linkage mapping.

Lesson 165: Random Amplified Polymorphic DNA & Amplified Fragment Length Polymorphism

1. What type of primer is used in Random Amplified Polymorphic DNA (RAPD) PCR amplification?

- A) Sequence-specific long primers
- B) Arbitrarily designed short PCR primers (10-12 bp) for random genomic amplification
- C) Single-stranded DNA probes
- D) RNA primers

2. Why does RAPD suffer from poor repeatability in laboratory testing?

- A) Because of unstable primer annealing during random amplification
- B) Because agarose gels destroy DNA fragments
- C) Because it uses single-stranded DNA probes
- D) Because of high cost

3. What initial step is performed in Amplified Fragment Length Polymorphism (AFLP) before adaptor ligation?

- A) Restriction endonuclease digestion of genomic DNA
- B) PCR amplification
- C) Gel electrophoresis
- D) Southern blotting

4. How does AFLP compare to RAPD regarding amplification stability and repeatability?

- A) AFLP has lower stability
- B) AFLP combines restriction digestion and specific PCR to yield stable, repeatable amplification
- C) They are identical
- D) RAPD is more stable

Answer Key

- 1. B — RAPD utilizes arbitrarily designed short PCR primers (10-12 bp) for random genomic amplification.
- 2. A — Unstable primer annealing during random amplification leads to poor repeatability.
- 3. A — AFLP methodology begins with restriction endonuclease digestion of genomic DNA, followed by adaptor ligation.
- 4. B — AFLP combines restriction digestion and specific PCR to yield stable amplification.

Lesson 166: Distribution of SNP

1. What is a Single Nucleotide Polymorphism (SNP)?

- A) A deletion of an entire chromosome arm
- B) A genetic variation resulting from the alteration of a single nucleotide base
- C) A duplication of a whole genome
- D) An inversion of a chromosome segment

2. Which detection method uses sequence-specific primers and PCR amplification for SNP analysis?

- A) RFLP

- B) PCR with sequence-specific primers
- C) Karyotyping
- D) Pedigree analysis

3. According to the comparison table in Lesson 166, what degree of polymorphism is exhibited by SNPs and STRs?

- A) Low
- B) Moderate
- C) High (both SNPs and STRs display a high degree of polymorphism)
- D) None

4. Which molecular marker technique relies on hybridization and requires radioactive probes, unlike PCR-based SNP assays?

- A) Microsatellites
- B) STS
- C) RFLP
- D) RAPD

Answer Key

1. B — A SNP is defined as a genetic variation resulting from the alteration of a single nucleotide.
2. B — SNP analysis utilizes PCR with sequence-specific primers.
3. C — Both SNPs and STRs display a high degree of polymorphism.
4. C — RFLP relies on hybridization and requires radioactivity, whereas SNP assays utilize PCR-based methods.

Lesson 167: SNP- Screening and importance

1. How frequently do Single Nucleotide Polymorphisms (SNPs) occur across the human genome?

- A) 1 in 10 bases
- B) Roughly 1 in 100 to 300 bases (up to 1 in 1,000 bases)
- C) 1 in 1 million bases
- D) 1 in 10 million bases

2. Where are SNPs distributed within genomic DNA?

- A) Exclusively in coding exons
- B) Exclusively in telomeres
- C) In both coding regions and, mostly, non-coding regions of the genome
- D) Only in mitochondrial DNA

3. What are the two main screening strategies used for SNP analysis according to Lesson 167?

- A) High-throughput sequencing only
- B) Analyzing many SNPs in a few individuals, or analyzing a few SNPs in many individuals
- C) Karyotyping and pedigree analysis
- D) Blood grouping and fingerprinting

4. How can human populations be categorized using individual genomic SNP patterns?

- A) Based on their eye color only

- B) Individual genomic SNP patterns/haplotypes allow people to be categorized into distinct population groups
- C) Based on height only
- D) They cannot be categorized

Answer Key

1. B — SNPs occur with high frequency, roughly 1 in 100 to 300 bases (to 1 in 1,000 bases).
2. C — SNPs are located in both coding regions and, mostly, non-coding regions of the genome.
3. B — The two SNP screening strategies are analyzing many SNPs in a few individuals, or a few SNPs in many individuals.
4. B — Individual genomic SNP patterns/haplotypes allow people to be categorized into distinct groups.

End of Quiz

BIO301 — 668 solved MCQs
Prepared by VU Learning HUB
vulearninghub.xyz