

MTH401 GRAND QUIZ

MTH401:Grand Quiz

Quiz Start Time: 08:01 AM, 06 July 2020

Question # 1 of 30 (Start time: 08:01:15 AM, 06 July 2020)

Total Marks: 1

The gravitational force exerted by the earth on a body of mass m is called ----- of the body.

Select the correct option

☐	Force
☐	weight

Question # 2 of 30 (Start time: 08:01:29 AM, 06 July 2020)

Total Marks: 1

The differential equation

$$\frac{du}{dx} + \frac{2}{x} dx = 3 - \frac{6}{x}$$

Select the correct option

[Reload Math Equations](#)

☐	$\frac{1}{x}$
☐	x^2
☐	$\frac{1}{x^2}$
☐	$-\frac{1}{x}$

Question # 3 of 30 (Start time: 08:01:43 AM, 06 July 2020)

Total Marks: 1

If the auxiliary equation $m^3 + m = 0$ has roots $m = 0, m = \pm i$ then the complementary function is

Select the correct option

[Reload Math Equations](#)

☐	$y_c = c_1 + c_2 \cos x + c_3 \sin x$
☐	$y_c = c_1 \cos x + c_2 \sin x$

Question # 4 of 30 (Start time: 08:01:48 AM, 06 July 2020)

Total Marks: 1

The differential equation

$$\frac{dy}{dx} - y = y^3$$

Select the correct option

[Reload Math Equations](#)

☐	Bernoulli's
☐	Homogeneous
☐	Cauchy
☐	Bessel

Question # 5 of 30 (Start time: 08:02:12 AM, 06 July 2020)

Total Marks: 1

A set of n functions

$$\{f_1(x), f_2(x), f_3(x), \dots, f_n(x)\}$$

is _____ if at least one of them can be expressed as a linear combination of the remaining.

Select the correct option

[Reload Math Equations](#)

☐	linear dependent
☐	linear independent

Question # 6 of 30 (Start time: 08:02:26 AM, 06 July 2020)

Total Marks: 1

What is annihilator operator of the function

$$g(x) = 4 \sin x$$

?

Select the correct option

[Reload Math Equations](#)

☐	$(D^2 - 1)$
☐	$(D^2 + 4)$
☐	$(D^2 - 4)$
☐	$(D^2 + 1)$

Question # 7 of 30 (Start time: 08:02:32 AM, 06 July 2020)

Total Marks: 1

Separable form $f(y)dy = g(x)dx$, of the differential equation: $y - x \frac{dy}{dx} = a \left(y^2 + \frac{dy}{dx} \right)$ is-----.

Select the correct option

[Reload Math Equations](#)

☐	$\frac{1}{y(1+ay)} dy = \frac{1}{x+a} dx$
☐	$\frac{1}{y(1-ay)} dy = \frac{1}{x+a} dx$
☐	$\frac{1}{y(1+ay)} dy = \frac{1}{x-a} dx$
☐	$\frac{1}{y(1-ay)} dy = \frac{1}{x-a} dx$

Question # 7 of 30 (Start time: 08:02:32 AM, 06 July 2020)

Total Marks: 1

Separable form $f(y)dy = g(x)dx$, of the differential equation: $y - x \frac{dy}{dx} = a \left(y^2 + \frac{dy}{dx} \right)$ is-----.

Select the correct option

[Reload Math Equations](#)

☒	$\frac{1}{y(1+ay)} dy = \frac{1}{x+a} dx$
☐	$\frac{1}{y(1-ay)} dy = \frac{1}{x+a} dx$
☐	$\frac{1}{y(1+ay)} dy = \frac{1}{x-a} dx$
☐	$\frac{1}{y(1-ay)} dy = \frac{1}{x-a} dx$

The differential equation $(\sin 2x - \tan y)dx - x \sec^2 y dy = 0$ is exact because - - - - -.

Select the correct option

[Reload Math Equations](#)

☒	$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} = -\sec^2 y$
☐	$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial y} = -\sec^2 y$
☐	$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} = \sec^2 y$
☐	$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial y} = \sec^2 y$

Which of the following is first order linear equation in unknown variable y?

Select the correct option

[Reload Math Equations](#)

☐	$x \frac{dy}{dx} + (\sin x) y = \cos x$
☐	$y \frac{dx}{dy} + (\sin y) x = \cos y$
☐	$y \frac{dx}{dy} + (\sin y) x = \cos x$
☐	$y \frac{dx}{dy} + (\sin x) x = \cos y$

Question # 10 of 30 (Start time: 08:03:06 AM, 06 July 2020)

Total Marks: 1

If initial amount of a radioactive isotope is 100g. What will be the amount at the end of 30 days such that $K=0.043$?

Select the correct option

☐	371.415
☐	380.560
☐	363.279
☐	360.351

Question # 11 of 30 (Start time: 08:03:13 AM, 06 July 2020)

Total Marks: 1

For $f(x, y) = x^2 - y^2$, $f(tx, ty) = \dots$

Select the correct option

[Reload Math Equations](#)

☐	$f(x, y)$
☐	$tf(x, y)$
☐	$t^2 f(x, y)$
☐	$t^3 f(x, y)$

Question # 12 of 30 (Start time: 08:03:19 AM, 06 July 2020)

Total Marks: 1

Which of following would be a constant solution of the separable differential equation: $\frac{dy}{dx} = e^{x+y}$?

Select the correct option

[Reload Math Equations](#)

- | | |
|-----------------------|----------------------------|
| ☐ | $y = 0$ |
| ☐ | $y = 1$ |
| ☐ | $y = a \in \mathbb{R}$ |
| ☐ | No constant solution exist |

Question # 13 of 30 (Start time: 08:03:24 AM, 06 July 2020)

Total Marks: 1

If we substitute

$$u = \frac{y}{x}$$

in differential equation

Select the correct option

[Reload Math Equations](#)

- | | |
|-----------------------|---------------------|
| ☐ | $ue^u du = dx$ |
| ☐ | $ue^{(-u)} du = dx$ |
| ☐ | $ue^u du = dy$ |
| ☐ | $ue^{(-u)} du = dy$ |

Question # 14 of 30 (Start time: 08:03:34 AM, 06 July 2020)

Total Marks: 1

Separable form $g(x)dx + f(y)dy = 0$, of the differential equation: $(2x \cos y) dx + x^2 (\sec y - \sin y) dy = 0$ is-----.

Select the correct option

[Reload Math Equations](#)

- ☐ $\left(\frac{2}{x}\right) dx + (\cos^2 y - \tan y) dy = 0$
- ☐ $\left(\frac{x}{2}\right) dx + (\sec^2 y - \cot y) dy = 0$
- ☐ $\left(\frac{x}{2}\right) dx + (\sec^2 y - \tan y) dy = 0$
- ☐ $\left(\frac{2}{x}\right) dx + (\sec^2 y - \tan y) dy = 0$

Question # 15 of 30 (Start time: 08:03:39 AM, 06 July 2020)

Total Marks: 1

A differential equation $M(x,y)dx + N(x,y)dy = 0$ is exact if there exists a multi - variable function $f(x,y)$ such that - - -

Select the correct option

[Reload Math Equations](#)

- ☐ $df(x,y) = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$
- ☐ $\int f(x,y) dx = \int \frac{\partial f}{\partial x} dx + \int \frac{\partial f}{\partial y} dy$
- ☐ $f(x,y) = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$
- ☐ $f(x,y) = \int \frac{\partial f}{\partial x} dx + \int \frac{\partial f}{\partial y} dy$

Question # 16 of 30 (Start time: 08:03:45 AM, 06 July 2020)

Total Marks: 1

 $\frac{d^3y}{dx^3} + y^4 = 0$ is a -----differential equation of order-----.

Select the correct option

[Reload Math Equations](#)

☐	linear, 4
☐	linear, 3
☐	non-linear, 4
☐	non-linear, 3

Which of the following is an equivalent form of the exact differential equation:
 $ydx + xdy = 0$?

Select the correct option

[Reload Math Equations](#)

☐	$d\left(\frac{x}{y}\right) = 0$
☐	$d\left(\frac{y}{x}\right) = 0$
☐	$d(xy) = 0$
☐	$d(x + y) = 0$

Question # 18 of 30 (Start time: 08:03:59 AM, 06 July 2020)

Total Marks: 1

In a Bernoulli equation

$$\frac{dy}{dx} - \frac{1}{xy} = xy^3$$

identify

Select the correct option

Reload Math Equations

☐	$\frac{1}{x}, x^3$
☐	y, xy^3
☐	$-y, xy^3$
☐	$-\frac{1}{x}, x^3$

Click to Save Answer & Move to Next Question

Question # 19 of 30 (Start time: 08:05:08 AM, 06 July 2020)

Total Marks: 1

The differential equation of orthogonal trajectory to the family of curves

$$x^2 + y^2 = C^2$$

is _____.

Select the correct option

Reload Math Equations

☐	$\frac{dy}{dx} = \frac{x}{y}$
☐	$\frac{dy}{dx} = \frac{y}{x}$
☐	$\frac{dy}{dx} = -\frac{x}{y}$
☐	$\frac{dy}{dx} = -\frac{y}{x}$

Click to Save Answer & Move to Next Question

Question # 20 of 30 (Start time: 08:05:17 AM, 06 July 2020)

Total Marks: 1

If the general solution of a separable differential equation is $\sin^{-1} y = \cos^{-1} x + c$, provided that $y(\frac{1}{\sqrt{2}}) = \frac{1}{\sqrt{2}}$, then $c = - - -$.

Select the correct option

[Reload Math Equations](#)

☐	$\frac{\pi}{2}$
☐	$\frac{\pi}{4}$
☐	$-\frac{\pi}{2}$
☐	0

[Click to Save Answer & Move to Next Question](#)

Maths 101: Grand Quiz

Quiz Start time: 08:07 AM, 06 July 2020

Question # 21 of 30 (Start time: 08:05:22 AM, 06 July 2020)

Total Marks: 1

The equation of free un-damped motion is _____.

Select the correct option

[Reload Math Equations](#)

☐	$\frac{d^2 x}{dt^2} + \frac{k}{m} x = 0$
☐	$\frac{d^2 x}{dt^2} - \frac{k}{m} x = 0$

Question # 22 of 30 (Start time: 08:05:35 AM, 06 July 2020)

Total Marks: 1

Which of the following is a boundary value problem(BVP)?

Select the correct option

[Reload Math Equations](#)

- | | |
|-----------------------|---|
| ☐ | $\frac{d^2y}{dx^2} + y = 0$ |
| ☐ | $\frac{d^2y}{dx^2} + y = 0, x \in [-2, 2]$ |
| ☐ | $\frac{d^2y}{dx^2} + y = 0, y(1) = -2$ and $y'(1) = 2$ |
| ☐ | $\frac{d^2y}{dx^2} + y = 0, y(0) = 0$ and $y\left(\frac{\pi}{2}\right) = 2$ |

A differential equation $M(x, y) dx + N(x, y) dy = 0$ is exact if and only if ---.

Select the correct option

[Reload Math Equations](#)

- | | |
|----------------------------------|---|
| ☒ | $\frac{\partial}{\partial x} M(x, y) = \frac{\partial}{\partial y} N(x, y)$ |
| ☐ | $\frac{\partial}{\partial y} M(x, y) = \frac{\partial}{\partial x} N(x, y)$ |
| ☐ | $\frac{d}{dx} M(x, y) = \frac{d}{dy} N(x, y)$ |
| ☐ | $\frac{d}{dy} M(x, y) = \frac{d}{dx} N(x, y)$ |

Saving

For 2nd order linear non-homogeneous differential equation

$$a_2(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

Select the correct option



☐	$y'(a) = y'_0, \quad y(b) = y_1$
☐	$y'(a) = y'_0, \quad y'(b) = y'_1$
☐	$y'(a) = y'_0, \quad y'(b) = y'_0$
☐	$y'(a) = y'_0, \quad y(b) = y_1$

Wronskian

$$W(1, e^x)$$

= _____.

Select the correct option



☐	0
☐	1
☐	-1
☐	e^x

For the non-exact differential equation $(x^2y + y) dx - xdy = 0$, if $\frac{\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x}}{N} = -x$, then the integrating factor is ---.

Select the correct option



- | | |
|-----------------------|--------------------|
| ☐ | $e^{\frac{1}{x}}$ |
| ☐ | $e^{-\frac{1}{x}}$ |
| ☐ | e^x |
| ☐ | e^{-x} |

MT H401: Grand Quiz

Quiz Start Time: 08:01 AM, 06 July 2020

Question # 27 of 30 (Start time: 08:06:23 AM, 06 July 2020)

Total Marks: 1

Which of following would be a general solution of the differential equation: $\frac{dy}{dx} = 4$?

Select the correct option



- | | |
|-----------------------|--------------|
| ☐ | $y = 4x + a$ |
| ☐ | $y = ax + 4$ |
| ☐ | $y = 4x + 4$ |
| ☐ | $y = ax + a$ |

Question # 28 of 30 (Start time: 08:06:29 AM, 06 July 2020)

Total Marks: 1

Which of the following substitution will transform the differential equation : $\frac{dy}{dx} = \frac{y}{x} + \sec\left(\frac{y}{x}\right)$, in to separable form?

Select the correct option

[Reload Math Equations](#)

☐	$y = v + x$
☐	$y = v - x$
☐	$y = vx$
☐	$x = vy$

Which of the following substitution will transform the differential equation : $\frac{dy}{dx} = \frac{y}{x} + \sec\left(\frac{y}{x}\right)$, in to separable form?

Select the correct option

[Reload Math Equations](#)

☐	$y = v + x$
☐	$y = v - x$
☐	$y = vx$
☐	$x = vy$

Question # 29 of 30 (Start time: 08:06:39 AM, 06 July 2020)

Total Marks: 1

Wronskian

$$W(x, 2x)$$

= _____.

Select the correct option

[Reload Math Equations](#)

☐	0
☐	1
☐	-1
☐	None of these

The differential equation $\frac{dx}{dy} + \frac{1}{y}x = 2 \sin y$ is first order linear in unknown ---

Select the correct option

[Reload Math Equations](#)

☐	variable x
☐	variable y
☐	multi - variables x and y
☐	$\frac{dy}{dx}$