

Mid-Term Quiz

Lecture # 1-22 😊

Grand Test

MTH645 Lec # 1-7

Question NO: 01 MCD's

1) if $X = \{1, 2, 3, 4, 5\}$ then $P(X) = ?$
tell about no of elements of Power set?

(a) 36 (b) 32 (c) 38 (d) 39

2) $\chi_A(x) = 1$ if

(a) $x \in A$ (b) $x \notin A$

3) $\chi_A(x) = 0$ if

(a) $x \in A$ (b) $x \notin A$

4) $A \cup B = \text{--- ?}$

(a) $\chi_A(x) \wedge \chi_B(x)$ (b) $\chi_A(x) \vee \chi_B(x)$

5) $A = \int \frac{\chi_A(x)}{x}$ is for ---?

(a) discrete case (b) continuous case

6) $\hat{A} = \{(x_1, 0), (x_2, 0.2)\}$

(a) Normal set

(b) Not a normal set

7) $A = \{(x_1, 1), (x_2, 0), (x_3, 1)\}$, $|A| = ?$

- (a) 2 (b) 3 (c) 4 (d) 5

8) $S(\vec{A}, \vec{B}) \in \text{---} ?$

- (a) (0, 1) (b) $[0, 1]$

9) $[a_1, b_1] \subseteq [a_2, b_2]$ iff --- ?

- (a) $a_1 \leq a_2$ and $b_1 \leq b_2$
(b) $a_2 \leq a_1$ and $b_1 \leq b_2$

10) $I_1 = [3, 5]$, $I_2 = [6, 7]$

- (a) $I_1 < I_2$ (b) $I_1 > I_2$

Question No: 02 :- Short Ques

Grand Test

Out

Mth645

lec 12-15

MCQ's:

- 1) $\prod_{i=1}^n A_i = \text{---?}$
 (a) A^2 (b) A^3 (c) A^n ✓
- 2) $(a, b) = (b, a)$ in cartesian product.
 (a) True (b) False ✓
- 3) Binary relation is subset of ---?
 (a) Fuzzy relation (b) cartesian product ✓
- 4) Many to one is also called ---?
 (a) surjection ✓ (b) injection
- 5) ~~Int~~jection also called ---?
 (a) Into, one to one ✓
 (b) Into, onto
- 6) R is said to be bijection if it is both ---?
 (a) one to many and one-to-one
 (b) Many to one and Into ✓

7) How many representation method of Relation.

(a) 3 (b) 4 (c) 2 (d) 5

8) $m_{ij} = 1$ if

(a) $(a_i, b_j) \in R$ (b) $(a_i, b_j) \notin R$

9) $R = \{(a, c), (b, d)\}$. $\text{Dom}(R) = ?$

(a) $\{a, b\}$ (b) $\{c, d\}$ (c) $\{a, c\}$

10) $M_R + M_S = ?$

(a) $M_{R \cup S}$ (b) $M_{R \cap S}$ (c) $M_{R \cdot S}$

Short Ques.

1) Define directed graph?

2) Define one to one correspondence.

3)

	x_1	x_2	x_3	x_4
x_1	0.2	0	0.5	0
x_2	0	0.3	0.7	0.8
x_3	0.1	0	0.4	0
x_4	0	0.6	0	0.1

$R =$

write $\text{dom}(R)$ and $\text{Rang}(R)$.

4) Define complement of fuzzy

relations.

Grand Test

MTH645 Lec # 16-18

Question No:01 MCQ's.

- 1) Triangular norms and co-norms were first introduced by theory of $\rightarrow$
 - (a) ☒ Probabilistic metric spaces.
 - (b) topological spaces.
- 2) How many conditions exists for T-norm?
 - (a) 1
 - (b) 3
 - (c) ☒ 4
 - (d) 6
- 3) $x \Delta_3 y = \text{---?}$
 - (a) xz
 - (b) ☒ xy
 - (c) $x+y$
- 4) For triangular norm the binary operation is $\rightarrow$
 - (a) ☒ delta
 - (b) Nebula
- 5) For co-norm the binary operation $\rightarrow$
 - (a) ☒ Nebula
 - (b) delta.
- 6) $0 \Delta x = \text{---?}$
 - (a) 1
 - (b) ☒ 0
 - (c) ∞
- 7) $x \Delta_5 y = \text{---?}$
 - (a) xy
 - (b) ☒ $x \wedge y$
 - (c) $x \vee y$

8) $x \Delta_3 y$ is not t-norm.

(a) true (b) False

9) $x \Delta_4 y$ is not t-norm.

(a) true (b) False

10) $x \leq y \Rightarrow -x \geq -y$

(a) True (b) False

Question No:02 Short Ques

1) Define t-norm?

2) Define co-norm?

3) write some examples of t-norm

4) Prove $0 \Delta x = 0$

5) Prove $x \Delta_2 (y \Delta_2 z) = (x \Delta_2 y) \Delta_2 z$

Long Ques. Ques # 03

1) Check min is t-norm or not?

2) Check $x \Delta_3 y$ is t-norm or not?

3) Check $x \Delta_4 y$ is t-norm or not?

Grand Test

Mth645

lec #21, 22

Question No: 01 MCQ's

1) if $\max(x, y) \neq 1$ then $x \Delta y =$
☒ (a) 0 (b) 1 (c) $\min(x, y)$

2) $\Delta_2 \geq \Delta_3$
 (a) True ☒ (b) False

3) $T(x, 0) =$ —
☒ (a) 0 (b) 1 (c) 2

4) $S(x, 1) =$ —
☒ (a) 1 (b) 0 (c) 2

5) $S(1, 0) =$ —
☒ (a) 1 (b) 0 (c) 2

6) $f(x) = x^x$ then $f^{-1}(x) =$ —
 (a) y ☒ (b) $x^{1/x}$ (c) $y^{1/y}$

7) $f(x) = x$ then $f^{-1}(x) =$ —
 (a) $\frac{1}{x}$ (b) x^3 ☒ (c) x

Which statement is correct

5 (a) Δ (union) with 0 is always 0

(b) Δ (union) with 0 is always positive.

(c) ∇ (union) with 0 cannot be negative.

6 $\Delta_1, \Delta_2, \Delta_3$ and Δ_4 all are _____ t-norm.

(a) Convex (b) Idempotent

(c) Archimedean. (d) None of these

MTH645

Lecture # 19, 20

Choose the correct answer.

1. Δ is idempotent if $x \Delta x =$ _____
 $\forall x \in [0, 1]$

a 1 ☒ (b) x ☐ 0 (d) None

2. A t-norm is Archimedean, if it
is _____ and $\forall a, b \in (0, 1)$
 $\exists$ a positive integer n s.t.

(a) $\text{convex}[a]^n = a \Delta a \Delta a \Delta a < b.$

(b) $\text{reave}[a]^n = a \Delta a \Delta a \Delta \dots < b.$

☒ (c) $\text{convex}[a]^n = a \Delta a \Delta a \Delta \dots \Delta a < b.$

(d) all of the above

3. $[a]^5$ is Archimedean t-norm

☒ (a) true (b) false

4. Min is t-norm as well
as archimedean t-norm

(a) true ☒ (b) false

MTH645

Lecture # 19, 20

Choose the correct answer

1. Δ is idempotent if $x \Delta x =$ _____
 $\forall x \in [0, 1]$

a 1 (b) x (c) 0 (d) None

2. A t-norm is Archimedean, if it
is _____ and $\forall a, b \in (0, 1)$
 $\exists$ a positive integer n s.t.

(a) $\text{convex}[a]^n = a \Delta a \Delta a \Delta a < b.$

(b) ~~convex~~ $[a]^n = a \Delta a \Delta a \Delta \dots < b.$

(c) $\text{convex}[a]^n = a \Delta a \Delta a \Delta \dots \Delta a < b.$

(d) all of the above

3. $[a]^5$ is Archimedean t-norm

(a) true (b) false

4. Min is t-norm as well
as archimedean t-norm

(a) true (b) false

4 $\min(1, x) = \text{---} ?$

- (a) 0
- (b) 1
- ☒ (c) x
- (d) None of these

5 Following is NOT the property of Δ -norm:

- (a) $x \Delta y = y \Delta x$
- ☒ (b) $0 \Delta x = x$
- (c) $(x \Delta y) \Delta z = x \Delta (y \Delta z)$
- (d) $1 \Delta x = x$

6 $\max(1, \emptyset) = \text{---} ??$

- (a) 0
- (b) $\emptyset$
- ☒ (c) 1
- (d) None.

$x \Delta_3 y = xy$ is not a Δ -norm

- (a) true
- ☒ (b) False

Choose the correct answer

1. If $I \Delta x = x$ satisfies for t -norm then it is true for,

(a) $I \Delta_1 x = x$

(b) $I \Delta_2 x = x$

(c) $I \Delta_3 x = x$

☒ (d) All of the above.

2. The binary operation for t -conorm will be:

(a) $\Delta : [0, 1]^2 \rightarrow [0, 1]$

(b) $\Delta : [0, 1] \rightarrow [0, 1]$

☒ (c) $\nabla : [0, 1]^2 \rightarrow [0, 1]$

(d) $\nabla : [0, 1] \rightarrow [0, 1]$

3. The binary operator nebula (∇) is used to represent $\neg$.

(a) t -norm

☒ (b) t -conorm

(c) both of these.

9

The distance between the points

$$\frac{x-a}{a_2-a_1} \text{ and } \frac{a_3-x}{a_3-a_2} \text{ in}$$

triangular fuzzy number is known

as _____

- (a) Diagonal (b) α -cut
(c) Chord (d) Tangent

10 According to the properties of α -cut, the interval of A_α become decrease/smaller when the value of α -cut _____.

- (a) decrease
(b) same as A_α
(c) increases
(d) (b) and (c) both.

4 There is only ~~one~~ maximum value at only one point in fuzzy number.

- (a) Triangular (b) Trapezoidal
(c) Cartesian (d) All of the above

5 The negative of triangular fuzzy number $\tilde{A} = (6, 7, 8)$ is

(a) $\tilde{A} = (-6, -7, -8)$

(b) $\tilde{A} = (-8, -7, -6)$

(c) $\tilde{A} = (-7, -6, -8)$

(d) None of these

6 If $\tilde{A} = (2, 4, 6)$ and $\tilde{B} = (3, 6, 7)$ the multiplication $\tilde{B} \times \tilde{A} =$

(a) $(6, 24, 42)$

(b) $(24, 15, 14)$

(c) None of these

7 $\tilde{A} - \tilde{B}$ and $\tilde{B} - \tilde{A}$ in triangular fuzzy numbers are always equal

(a) True

(b) False

MTH645

Lecture 8-11

Total Marks: 35

Choose the correct answer: $1 \times 10 = 10$

1. The convex condition

$$(\alpha' < \alpha) \Rightarrow (a_1^{(\alpha')} \leq a_1^{(\alpha)}, a_3^{(\alpha')} \leq a_3^{(\alpha)})$$

can also be written as:

(a) $\alpha' < \alpha \Rightarrow A_{\alpha'} \subset A_{\alpha}$

(b) $\alpha' < \alpha \Rightarrow A_{\alpha'} = A_{\alpha}$

(c) $\alpha' < \alpha \Rightarrow A_{\alpha'} \supset A_{\alpha}$

(d) $\alpha' < \alpha \Rightarrow A_{\alpha} \subset A_{\alpha'}$

2. According to the convex condition the α -cut is:

(a) discrete

(b) finite

(c) definite

(d) continuous

3. The triangular fuzzy number is always represented in _____ points

(a) 1
(c) 3

(b) 2

(d) 4

10 If $I_1 = [2, 3]$ and $I_2 = [0, 2]$

the intersection $I_1 \cap I_2$ is :

(a) $[1, 2]$

(b) $[0, 3]$

(c) Undefined

(d) None of these

Ans the following: $2 * 3 = 6$

11 Find the inclusion of I_1 in I_2
where $I_1 = [2, 6]$; $I_2 = [3, 7]$

12 Define support of fuzzy set and
give example

13 IF $\tilde{B} = \{(x_1, 0.5), (x_2, 0.7), (x_3, 0.6), (x_4, 1), (x_5, 0)\}$
then find $\bar{\tilde{B}}$??

Ans the following $3 * 3 = 9$

4 IF $C = \{7, 8, 9\}$, $D = \{2, 4, 7\}$
then find

$A \cup B \rightarrow \chi_{A \cup B}(x)$

where $x \in C$ and or $x \in D$

7 The fuzzy set $\tilde{A} : \mathbb{R} \rightarrow [0, 1]$ defined by $\tilde{A}(x) = e^{-x^2}$ is normal at

(a) $x = 1$

(b) $x = 0$

(c) $x = 1.5$

(d) $x = 0.5$

8 The degree of subset hood of $\tilde{B}$ in $\tilde{A}$ is defined as

(a) $S(\tilde{A}, \tilde{B}) = \frac{|\tilde{A} \cap \tilde{B}|}{|\tilde{A}|}$

(b) $S(\tilde{A}, \tilde{B}) = \frac{|\tilde{A} \cap \tilde{B}|}{|\tilde{B}|}$

(c) $S(\tilde{B}, \tilde{A}) = \frac{|\tilde{B} \cap \tilde{A}|}{|\tilde{B}|}$

(d) $S(\tilde{B}, \tilde{A}) = \frac{|\tilde{B} \cap \tilde{A}|}{|\tilde{A}|}$

9 Cardinality of fuzzy set C is where $C = \{(a, 0.5), (b, 0.6), (c, 0.7)\}$

(a) 2

(b) 1.8

(c) 1.6

(d) 2.5

4. If $A \subseteq B$, $B = \{a, b, c, d, e\}$

$A = \{c, d, e\}$ then

$$\chi_A(e) =$$

(a) 0

(b) 1

5. Fuzzy logic is an approach to computing based on:

(a) Degree of truth and false

(b) Degree of truth or false

(c) Degree of truth

(d) None of these.

6. If $A = \{(2, 0.4), (4, 0.6), (5, 1)\}$
the membership value of 4 is ____
(i.e. $\mu_A(4) =$ ____)

(a) 0.4

(b) 1

(c) 0.6

(d) 0

Ourself!! :)

MTH645

Fuzzy Logics

Lecture 1-7

Total Marks : 35

Choose the correct answer (10) = 1 * 10

1- If X and Y are two sets then the function defined as :

(a) It assigns elements of x to element of y

(b) It assigns unique elements of x to unique elements of y

(c) It assigns element of x to unique elements of y

2- If $A \subseteq X$ and $X = \{3, 4, 5, 6\}$
 $A = \{4, 5\}$. then $\chi_A(4) = \underline{\hspace{2cm}}$

(a) 1

(b) 0

3- The membership value of any fuzzy set lie in the interval

(a) $(0, 1)$ (b) $(0, 1)$

(c) $[0, 1)$ (d) $[0, 1]$

GRAND TEST

Math 645

LEC NO: 8 - 11

MCQS

1) The Convex Condition may be written as $(\alpha' < \alpha) \Rightarrow$ _____

- (a) ☒ $A_\alpha < A_{\alpha'}$ (b) $A_{\alpha'} < A_\alpha$

2) The Negative Triangular Fuzzy Number of $(6, 5, 9)$

- (a) $(-6, -5, -9)$ (b) ☒ $(9, -5, -6)$

3) $A_\alpha =$ _____

- (a) ☒ $[\alpha_1, \alpha_3]$ (b) $[\alpha_1, \alpha_2]$

4) The Inverse of Trapezoidal Number of $(4, 5, 6)$

- (a) ☒ $(\frac{1}{6}, \frac{1}{5}, \frac{1}{4})$ (b) $(\frac{1}{4}, \frac{1}{5}, \frac{1}{6})$

5) The Negative of Triangular Fuzzy Number is Denoted as:

- (a) ☒ $\tilde{A}$ (b) $-A$

6) $R^n =$ _____

- (a) Dimensional Euclidian vector
(b) ☒ Non Dimensional Euclidian vector space

LRAND TEST

Maths: (645)

Le§ 19.20

Q No:1 M.C.Q.S

1) Δ is idempotent $\text{iff } x \Delta x =$ _____

(a) Δx (b) x (c) xx

2) $C = \delta \Delta S$ where Δ is _____

(a) convex T-norm (b) Archimedean

3) Min is an Archimedean T-norm

(a) True (b) False

4) $\Delta_1, \Delta_2, \Delta_3$ and Δ_4 are all

Archimedean T-norm

(a) True (b) False

5) Δ is archimedean T-norm

So $\forall a > b$

(a) True (b) False

6) $\text{If } \Delta$ is Archimedean t-norm

Then $\forall a, b \in$ _____

(a) $(0, 1)$ (b) $[0, 1)$ (c) $[0, 1]$

RAND TEST

Maths: 645:

Question No: 1 M.C.Q.s

- i) $\prod_{i=1}^n A_i =$
☒ (a) A^n ☐ (b) A^i ☐ (c) A^m
- ii) If $\text{mid} = 1$ Then
☒ (a) $(a_i, b_j) \in R$ ☐ (b) $(a_i, b_j) \notin R$
- iii) If $\text{mid} = 0$ Then
☐ (a) $(a_i, b_j) \in R$ ☒ (b) $(a_i, b_j) \notin R$
- iv) Directed Graph is also called -
☒ (a) Digraph ☐ (b) Biograph
- v) Fuzzy Relation has degree of membership value lies in
☐ (a) $(0, 1)$ ☒ (b) $[0, 1]$ ☐ (c) $[0, 2]$
- vi) Sum $A+B =$
☒ (a) $\max[a_{ij}, b_{ij}]$ ☐ (b) $\min[a_{ij}, b_{ij}]$
- vii) If $A \leq B$ Then
☒ (a) $S_A \leq S_B$ ☐ (b) $S_A \geq S_B$
- viii) $MR_{US} =$
☒ (a) $MR \neq Ms$ ☐ (b) $MR = Ms$
- ix) $Ms.R = MR.Ms$
☒ (a) True ☐ (b) False