

Zoology

4th semester

MUHAMMAD HABIB



Smithsonian

Question # 8 of 10 (Start time: 07:06:48 PM, 23 December 2021)

Osmoregulatory structure that is found in turbellarians is called

Select the correct option

☐

Metanephridia

☐

Protanephridia

☐

Mesonephridia

☐

kidney

Question # 4 of 10 (Start time: 07:01:54 PM, 23 December 2021)

Jelly fish belongs to which class of Cnidarians?

Select the correct option

☐	Aconitina
☐	Anthozoa
☐	Hydrozoa
☐	Scyphozoa

ZOO201 - Animal Diversity: Invertebrates (QUIZ NO.1)

Question # 6 of 10 (Start time: 07:04:40 PM, 23 December 2021)

Each proglottid in tapeworm consists of a complete set of

Select the correct option

- | | |
|-----------------------|---------------------|
| ☐ | Respiratory system |
| ☐ | Excretory system |
| ☐ | Reproductive system |
| ☐ | Digestive system |

Question # 7 of 10 (Start time: 07:05:56 PM, 23 December 2021)

Class of Platyhelminthes that mostly possess the members that serve as fish parasites is

Select the correct option

☐ Turbellaria

☐ Trematoda

☐ Monogenea

☐ Cestodea

Ascaris lumbricoides commonly known as

Select the correct option

☒	The giant intestinal roundworm of humans
☐	The giant liver roundworm of humans
☐	The giant tapeworm of human
☐	The giant tapeworm of beef

Question # 2 of 10 (Start time: 06:59:26 PM, 23 December 2021)

Which class of cnidarians represents fresh water species?

Select the correct option

☐

scyphozoa

☐

anthozoa

☐

hydrozoa

☐

cubozoa

Question # 2 of 10 (Start time: 07:58:02 PM, 23 December 2021)

Members of phylum Kinorhyncha are exclusively

Select the correct option

☐	bottom dwellers
☐	parasitic
☒	marine
☐	freshwater

Question # 8 of 10 (Start time: 08:07:23 PM, 23 December 2021)

What is true about nemertean s?

Select the correct option

☐	They are monoecious
☐	None of these
☐	They are parthenogenic
☐	They are dioecious



ZOO201 - Animal Diversity: Invertebrates (QUIZ NO.1)

Question # 9 of 10 (Start time: 07:08:16 PM, 23 December 2021)

Common name of *Taeniarhynchus saginatus* is

Select the correct option

☐	Liver fluke
☐	Human worm
☐	Sheep tapeworm
☐	Beef tapeworm

Question #1 of 10 (Start time: 07:56:22 PM, 23 December 2021)

Class of the phylum Mollusca that serve as intermediate hosts for some medically important trematode parasites of human is

Select the correct option



caudofoveata



bivalvia



gastropoda



cephalopoda

Click to Save Answer & Move to

ZOO201 - Animal Diversity: Invertebrates (QUIZ NO.1)

Question # 6 of 10 (Start time: 08:05:10 PM, 23 December 2023)

The outer layer of Mollusca shell is called

Select the correct option

☐	mantle
☒	Periostracum
☐	mantle cavity
☐	Prismatic

Question # 5 of 10 (Start time: 08:03:17 PM, 23 December 2021)

Ephyra is the larval form of

Select the correct option

☐	Hydra
☐	Sea anemones
☐	Obelia
☐	Aurelia

Elephantiasis is caused by

Select the correct option

☐	Pork tapeworm
☐	Pin worm
☒	Filarial worm
☐	Elephant worm

 <https://www.mdpi.com> › pdf

The Emergence of the Bilateral Symmetry in Animals - MDPI

[More results](#)

What were the first bilateral animals?



630 million years ago

It seems likely that the first bilateral animal was **a kind of worm**. *Vernanimalcula guizhouena*, which dates from around 600 million years ago, may be the earliest bilateral animal found in the fossil record. 14-Jul-2009

NS <https://www.newscientist.com> › article

Timeline: The evolution of life | New Scientist

[More results](#)

Which animal has a bilaterally symmetrical body plan?



Nemertes Research - Wikipedia

Nemertes Research is an American research-advisory firm that specializes in analyzing and ... while others mention its name as a concrete...

[https://en.m.wikipedia.org › wiki](https://en.m.wikipedia.org/wiki/Nemertes_Research)



Nemertea - Wikipedia

Nemertea is a phylum of invertebrate animals also known as ribbon worms or proboscis worms.



More results from en.m.wikipedia.org

People also ask





WhatsApp • 10 messages from 6 ch... • Now



BS Zoology 1st Semster Vu: +92 303 4261378 D

BS ZOOLOGY VU: S Akhtar Bt101 ki assignme...

Safdar Bai Habib bhai midterm ki date sheet B'...

Zoology Helping Group (5 messages): +92 347...

+92 307 7598279 Thank you bro + 1

and asymmetry. Learning Objectives. Differentiate...

People also ask



What was the first animal to have bilateral symmetry?



The bilateral symmetry in Animals is probably developed from **a sea anemone (Cnidaria)** with radially symmetry (*Nematostella vectensis*) [7–11]. In addition, many plants exhibit different kinds of symmetry, e.g., rotational symmetry, but also bilateral symmetry. 04-Feb-2021



<https://www.mdpi.com> › pdf

The Emergence of the Bilateral

Question # 5 of 10 (Start time: 09:21:07 PM, 23 December 2021) Total Marks:

Nemertean are commonly called

Select the correct option

- | | |
|-----------------------|---------------------------------|
| ☐ | roundworms or flatworms |
| ☐ | flatworms or ribbon worms |
| ☐ | tapeworms |
| ☐ | proboscis worms or ribbon worms |

Click to Save Answer & Move to Next Question

Question # 10 of 10 (Start time: 09:27:02 PM, 23 December 2021) Total Marks:

The first bilaterally symmetrical animal belongs to which of the following phylum?

Select the correct option

☐	Annelida
☐	Coelenterate
☐	Mollusca
☐	Platyhelminthes

Click to Save Answer & Move to Next Question

Question # 7 of 10 (Start time: 09:23:54 PM, 23 December 2021) Total Marks: 1

Characteristics that define rotifers

Select the correct option

- | | |
|-----------------------|--|
| ☐ | Bilaterally symmetrical, coelomates, unsegmented organisms |
| ☐ | Bilaterally symmetrical, coelomates, unsegmented organisms |
| ☐ | Radially symmetrical, pseudocoelomates, unsegmented organisms |
| ☐ | Bilaterally symmetrical, pseudocoelomates, unsegmented organisms |

Click to Save Answer & Move to Next Question



Question # 6 of 10 (Start time: 09:22:15 PM, 23 December 2021) Total Marks:

In rotifers ovary and vitellarium often fuse to form a structure that is known as

▶ Select the correct option

☐	Seminal vesicle
☐	Seminal receptacle
☐	Germovitellarium
☐	Trophi

[Click to Save Answer & Move to Next Question](#)

Question # 3 of 10 (Start time: 09:18:53 PM, 23 December 2021) Total Marks:

Most of the nematodes are

Select the correct option

☐	monoecious and polymorphic
☐	dioecious and polymorphic
☐	dioecious and dimorphic
☐	monoecious and dimorphic

Click to Save Answer & Move to Next Question

Question # 2 of 10 (Start time: 09:18:11 PM, 23 December 2021) Total Marks:

The body of tapeworm consists of

Select the correct option

☐	Proglottids, scolex and strobila
☐	Scolex and strobila
☐	Scolex, neck and proglottids
☐	Scolex and neck

Click to Save Answer & Move to Next Question

Question # 9 of 10 (**Start time: 03:49:19 PM, 23 December 2021**)

The larva of phylum nemertean is known as

Select the correct option



pilidium larva



tadpole



planula larva



muller's larva

Question # 8 of 10 (Start time: 03:47:05 PM, 23 December 2021)

The group Aschelminthes has

Select the correct option



ten different phyla



seven different phyla



nine different phyla



eight different phyla

Question # 7 of 10 (Start time: 03:46:06 PM, 23 December 2021)

Which of following class of Platyhelminthes does not contain parasitic species?

Select the correct option

☒	Turbellaria
☐	Trematoda
☐	Cestoidea
☐	Monogenea

ZOO201 - Animal Diversity: Invertebrates (QUIZ NO.1)

Question # 10 of 10 (Start time: 03:50:39 PM, 23 December 2021)

Incomplete alimentary canal is found in which of the following class of Platyhelminthes

Select the correct option

☐	Turbellaria
☐	Nematode
☐	Cubozoa
☐	Trematode

The outer layer of Mollusca shell is called

Select the correct option

☐	Periostracum
☐	mantle
☐	mantle cavity
☐	Prismatic

Question # 4 of 10 (Start time: 03:42:43 PM, 23 December 2021)

Number of host involved in the life cycle of parasitic nematodes is/are

Select the correct option

☒	two
☐	four
☐	three
☐	one

Question #5 of 10 (Start time: 03:43:48 PM, 23 December 2021)

In *Taenia saginata* egg developed into six hooked larvae called

Select the correct option

☐	Cercariae
☒	Onchosphere
☐	Miracidia
☐	Metacercaria

Question # 2 of 10 (Start time: 03:26:15 PM, 23 December 2021)

Sexual maturity in a larval body form is called

Select the correct option

☐	Monecious
☐	Paedomorphosis
☐	Protandry
☐	Polygyny

Question # 5 of 10 (Start time: 03:29:00 PM, 23 December 2021)

Which class of cnidarians represents fresh water species?

Select the correct option

☐	scyphozoa
☐	cubozoa
☐	hydrozoa
☐	anthozoa

ZOO201 - Animal Diversity: Invertebrates (QUIZ NO.1)

Question # 10 of 10 (**Start time: 03:04:54 PM, 23 December 2021**)

Platyhelminthes are best described as

Select the correct option

☐	Flatworms, triploblastic, pseudocoelomate animals
☐	Flatworms, triploblastic, coelomates animals
☐	Flatworms, diploblastic , acoelomates animals
☐	Flatworms, triploblastic, acoelomates animals

Question # 1 of 10 (**start time: 03:17:49 PM, 23 December 2021**)

In tapeworms, glands that help in the formation of egg capsule are

Select the correct option

☐	Ovarian glands
☐	Salivary glands
☐	Vitelline glands
☐	Mehlis' glands

Question # 6 of 10 (**Start time: 03:30:07 PM, 23 December 2021**)

The body of tapeworm consists of

Select the correct option

☐	Proglottids, scolex and strobila
☐	Scolex and strobila
☒	Scolex, neck and proglottids
☐	Scolex and neck

Question # 8 of 10 (Start time: 03:01:08 PM, 23 December 2021)

The body surface of kinorhynch is devoid of cilia and is composed of 13 or 14 definite units called

Select the correct option

☐	placids
☐	scalids
☐	segments
☐	zonites

Click to

The rotifers are initially named as

Select the correct option

☐	Wheel animalcules
☐	Rods
☐	Collared organisms
☐	Crown organisms

Question # 6 of 10 (**Start time: 02:58:24 PM, 23 December 2021**)

_____is the Larval stage of Filarial worm.

Select the correct option

☐	Tadpole
☐	Filariform
☐	Rhabditiform
☐	Microfilariae

Question # 10 of 10 (**Start time: 02:48:46 PM, 23 December 2021**)

The intermediate host in the life cycle of *Taenia saginata* is

Select the correct option

☐	cattle
☐	fish
☐	sheep
☐	human

Sea anemones and corals belong to which class of phylum cnidarian?

Select the correct option

☐	cubozoa
☐	anthozoa
☐	hydrozoa
☐	scyphozoa

General characteristics that define phylum Mollusca are

Select the correct option

☐	Bilateral symmetry, Protostome , Triploblastic
☐	Bilateral symmetry, Protostome , Diploblastic
☐	Radial symmetry, Deuterostome, Diploblastic
☐	Radial symmetry, Protostome, Triploblastic

Question # 4 of 10 (**Start time: 02:55:40 PM, 23 December 2021**)

Nematodes commonly known as

Select the correct option

☐	Flat worms
☐	Crown worm
☐	Round worms
☐	Tape worms

Question # 2 of 10 (**Start time: 02:52:50 PM, 23 December 2021**)

In cnidarians partially digested food is phagocytized by which of the following cells

Select the correct option

☐	Epidermis cells
☐	Endodermis cells
☐	nutritive-muscular cells
☐	Nematocyst

Question # 1 of 10 (Start time: 02:51:21 PM, 23 December 2021)

Pharynx of rotifers contains a unique muscular structure called

Select the correct option



Mastax



Crown



Lori



Epiglottis

Gastrovascular cavity of cnidarians is mainly composed of

Select the correct option

☐	epidermis
☐	mesodermis
☐	gastrodermis
☐	endodermis

Question # 8 of 10 (Start time: 02:45:56 PM, 23 December 2021)

All animals are unsegmented worms except

Select the correct option

☐	Planaria	
☐	Fasciola hepatica	
☐	Polystoma	
☐	Taenia solium	

Question #1 of 10 (Start time: 02:34:00 PM, 23 December 2021)

Number of identified species of rotifers are

Select the correct option

☐	4000
☐	3000
☐	2000
☐	5000

Click to Save

Question # 4 of 10 (Start time: 02:39:47 PM, 23 Decembe

Sea anemone is

Select the correct option

☐	Triploblastic, bilaterally symmetrical animal
☐	Diploblastic, bilaterally symmetrical animal
☐	Triploblastic, radially symmetrical animal
☐	Diploblastic, radially symmetrical animal

Question # 3 of 10 (Start time: 02:38:12 PM, 23 December 2021)

Feeding polyp in Obelia is called

Select the correct option

☐	Gastrozoid
☐	Planula
☐	Oozoid
☐	Gonozoid

Question # 2 of 10 (**Start time: 02:36:52 PM, 23 December 2021**)

Which of following class of Platyhelminthes does not contain parasitic species?

Select the correct option

☐	Monogenea
☐	Trematoda
☐	Cestoidea
☐	Turbellaria

Question # 5 of 10 (Start time: 02:40:53 PM, 23 December 2021)

The larva of phylum nemertean is known as

Select the correct option

☐	muller's larva
☐	planula larva
☒	pilidium larva
☐	tadpole

Question # 6 of 10 (Start time: 02:43:23 PM, 23 December 2021)

Class of Platyhelminthes that mostly possess the members that serve as fish parasites is

Select the correct option

☐	Trematoda
☐	Monogenea
☐	Cestoidea
☐	Turbellaria

Question # 7 of 10 (Start time: 02:44:21 PM, 23 December 2021)

Glandular system in some species of aquatic nematodes consists of ventral glands that are called

Select the correct option

☐	Pituitary gland
☐	Master gland
☒	Renettes
☐	Antenna

Click to