MTH603 FINAL TERM SOLVED MCQs BY EMRAAN HAAKIM

(VISIT VURANK FOR MORE)

Exact solution of 2/3 is not exists.
TRUE
FALSE

The Jacobi's method is A method of solving a matrix equation on a matrix that has
____zeros along its main diagonal.

No

A 3 x 3 identity matrix have three and_____eigen values.
Same

Eigenvalues of a symmetric matrix are all_____.
Real
Comple
x Zero
Positive

The Jacobi iteration converges, if A is strictly diagonally dominant. TRUE
FALSE

Below are all the finite difference methods EXCEPT_____.

Jacobi's method
Newton's backward difference method
Stirlling formula
Forward difference method

If n x n matrices A and B are similar, then they have the same eigenvalues (with the same multiplicities).
TRUE
FALSE

Emraan Haakim.03349666619

If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix, the eigenvalues of A are the diagonal entries of A.

TRUE
FALSE

The characteristics polynomial of a 3x 3 Identity matrix is_____, if x is the Eigen values of the given 3 x 3 identity matrix. Where symbol ^ shows power.

(X-1)^3
(x+1)^3
X^3-1
X^3+1

Two matrices with the same characteristic polynomial need not be similar.

TRUE
FALSE

Bisection method is a

Bracketing method
Open method

Regula Falsi means

Method of Correct position
Method of unknown position
Method of false position
Method of known position

Eigenvalues of a symmetric matrix are all_____.
Select correct option:

Real
Zero
Positive
Negative

An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.
Select correct option:

TRUE
FALSE

Emran Haakim.03349666619

Exact solution of 2/3 is not exists.
Select correct option:

TRUE
FALSE

The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric _____definite matrices A.
Select correct option:

Positive
Negative

Differences methods find the_____solution of the system.
Select correct option:

Numerical
Analytical

The Power method can be used only to find the eigenvalue of A that is largest in absolute value—we call this Eigenvalue the dominant eigenvalue of A.
Select correct option:

TRUE
FALSE

The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its_____.
Select correct option:

Main diagonal
Last column
Last row
First row

Emraan Haakim.03349666619

If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix , the eigenvalues of A are the diagonal entries of A.
Select correct option:

TRUE
FALSE

A 3 x 3 identity matrix have three and different Eigen values.
Select correct option:

TRUE
FALSE

Newton Raphson method falls in the category of

Bracketing method
Open Method
Iterative Method
Indirect Method

Newton Raphson method is also known as
Tangent Method
Root method
Open Method
Iterative Method

Secant Method uses values for approximation

1
3
2
4

Secant Method is than bisection method for finding root
Slow
Faster

In Newton Raphson method

Root is bracketed
Root is not bracketed

Regula falsi method and bisection method are both

Emraan Haakim.03349666619

Convergent
Divergent

In bisection method the two points between which the root lies are

Similar to each other
Different
Not defined
Opposite

In which methods we do not need initial approximation to start
Indirect Method
Open Method
Direct Method
Iterative Method

Root may be

Complex
Real
Complex or real
None

In Regula falsi method we choose points that have signs

2 points opposite signs
3 points opposite signs
2 points similar signs
None of the given

In a bounded function values lie between
1 and -1
1 and 2
0 and 1
0 and -2

Newton Raphson method is a method which when it leads to division of number close to zero
Diverges
Converges

Which of the following method is modified form of Newton Raphson Method?
Regula falsi method
Bisection method
Secant method

Emraan Haakim.03349666619

Jacobi's Method

Which 1 of the following is generalization of Secant method?
Muller's Method
Jacobi's Method
Bisection Method
N-R Method

Secant Method needs starting points

2
3
4
1

Near a simple root Muller's Method converges than the secant method

Faster
Slower

If S is an identity matrix, then

$S^{-1} = S$
$S^t = S$
$S^{-1} = S^t$
All are true

If we retain r+1 terms in Newton's forward difference formula, we obtain a polynomial of degree ---- agreeing with y_x at $x_0, x_1, \ldots, x_r$

r+2
r+1
R
R-1

P in Newton's forward difference formula is defined as

$$p = \left(\frac{x - x_0}{h}\right)$$

$$p = \left(\frac{x + x_0}{h}\right)$$

Emraan Haakim.03349666619

$$p = (\frac{x + x_n}{h})$$

$$p = (\frac{x - x_n}{h})$$

Octal numbers has the base

10
2
8
16

Newton's divided difference interpolation formula is used when the values of the independent variable are

Equally spaced

Not equally spaced

Constant
None of the above

Given the following data

x	0	1	2	4
$f(x)$	1	1	2	5

Value of $f(2, 4)$ is

1.5

3
2

1

If $y(x)$ is approximated by a polynomial $p_n(x)$ of degree n then the error is given by

$\square(x) = y(x) + P_n(x)$

$\square(x) = y(x) - P_n(x)$

Emraan Haakim.03349666619

$\square(x) = y(x) \times P_n(x)$

$\square(x) = P_n(x) + y(x)$

Let I denotes the closed interval spanned by $x_0, x_1, x_2, x_3, x_4, x_5, x_6, \bar{x}_7, x$. Then $F(x)$ vanishes ------times in the interval I .

N-1
N+2
N
N+1

Differential operator in terms of forward difference operator is given by

$$D = \frac{1}{h}(\Delta + \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} + \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} + \ldots)$$

$$D = \frac{1}{h}(\Delta + \frac{\Delta^2}{2} + \frac{\Delta^3}{3} + \frac{\Delta^4}{4} + \frac{\Delta^5}{5} + \ldots)$$

$$D = \frac{1}{h}(\Delta - \frac{\Delta^2}{2} + \frac{\Delta^3}{3} - \frac{\Delta^4}{4} + \frac{\Delta^5}{5} \ldots)$$

$$D = \frac{1}{h}(\Delta - \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} - \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} \ldots)$$

Finding the first derivative of $f(x)$ at x =0.4 from the following table:

x	0.1	0.2	0.3	0.4
$f(x)$	1.10517	1.22140	1.34986	1.49182

Differential operator in terms of---------- will be used.

Forward difference operator
Backward difference operator
Central difference operator
All of the given choices

Emraan Hakim.03349666619

For the given table of values

x	0.1	0.2	0.3	0.4	0.5	0.6
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f'(0.1)$, using two-point equation will be calculated as.....

-0.5
0.5
0.75
-0.75

In Simpson's 1/3 rule, $f(x)$ is of the form

$ax+b$

- ax^2+bx+c
- ax^3+bx^2+cx+d
- $ax^4+bx^3+cx^2+dx+e$

$$I=\int_a^b f(x)dx$$

While integrating , h , width of the interval, is found by the formula-----.

$\frac{b-a}{n}$

$\frac{b+a}{n}$

$\frac{n}{a-b}$

$\frac{n}{}$

None of the given choices

To apply Simpson's 1/3 rule, valid number of intervals are.....

Emraan Haakim.03349666619

7
8
5
3

For the given table of values

x	0.1	0.2	0.3	0.4	0.5	0.6
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f''(0.2)$ **, using three-point equation will be calculated as ……**

17.5
12.5
7.5
-12.5

To apply Simpson's 1/3 rule, the number of intervals in the following must be

2
3
5
7

To apply Simpson's 3/8 rule, the number of intervals in the following must be

10
11
12
13

If the root of the given equation lies between a and b, then the first approximation to the root of the equation by bisection method is ……

$\frac{(a+b)}{2}$
$\frac{(a-b)}{2}$

Emraan Haakim.03349666619

$\frac{(b-a)}{2}$

None of the given choices

.....lies in the category of iterative method.

Bisection Method
Regula Falsi Method
Secant Method
All of the given choices

For the equation $x^3+3x-1=0$, the root of the equation lies in the interval.....

(1, 3)
(1, 2)
(0, 1)
(1, 2)

Rate of change of any quantity with respect to another can be modeled by

An ordinary differential equation
A partial differential equation

A polynomial equation

None of the given choices

If

$\frac{dy}{dx}=f(x,y)$

Then the integral of this equation is a curve in

None of the given choices

Xt-plane
Yt-plane
Xy-plane

In solving the differential equation

Emraan Haakim.03349666619

$y' = x + y \; ; y(0.1) = 1.1$

$h = 0.1$, **By Euler's method** $y(0.2)$ **is calculated as**

1.44
1.11
1.22
1.33

In second order Runge-Kutta method
k_1 is given by

$k_1 = hf(x_n, y_n)$
$k_1 = 2hf(x_n, y_n)$
$k_1 = 3hf(x_n, y_n)$
None of the given choices

In fourth order Runge-Kutta method, k_2 is given by

$k_2 = hf\left(x_n + \frac{h}{2}, y_n + \frac{k_1}{2}\right)$
$k_2 = hf\left(x_n + \frac{h}{3}, y_n + \frac{k_1}{3}\right)$
$k_2 = hf\left(x_n - \frac{h}{3}, y_n - \frac{k_1}{3}\right)$
$k_2 = hf\left(x_n - \frac{h}{2}, y_n - \frac{k_1}{2}\right)$

In fourth order Runge-Kutta method, k_4 **is given by**

$k_3 = hf(x_n + 2h, y_n + 2k_3)$
$k_3 = hf(x_n - h, y_n - k_3)$
$k_3 = hf(x_n + h, y_n + k_3)$
None of the given choices

Emraan Haakim 03349666619

Adam-Moulton P-C method is derived by employing

Newton's backward difference interpolation formula
Newton's forward difference interpolation formula
Newton's divided difference interpolation formula
None of the given choices

The need of numerical integration arises for evaluating the definite integral of a function that has no explicit_____or whose antiderivative is not easy to obtain

Derivatives
Antiderivative

If $|A| \neq 0$ then system will have a

Definite solution
Unique solution
Correct solution
No solution

If $|A| \neq 0$ then

There is a unique solution
There exists a complete solution
There exists no solution
None of the above options

Direct method consists of method
2
3
5
4

We consider Jacobi's method Gauss Seidel Method and relaxation method as
Direct method
Iterative method
Open method
All of the above

In Gauss Elimination method Solution of equation is obtained in
3 stages
2 stages
4 stages
5 stages

Emraan Haakim.03349666619

Gauss Elimination method fails if any one of the pivot values becomes
Greater
Small
Zero
None of the given

Changing the order of the equation is known as

Pivoting
Interpretation

Full pivoting is than partial pivoting
Easy
More complicated

The following is the variation of Gauss Elimination method

Jacobi's method
Gauss Jordan Elimination method

Courts reduction method is also known as Cholesky Reduction method
True
False

Jacobi's method is also known as method of Simultaneous displacement
True
False
Gauss Seidel method is also known as method of Successive displacement
False
True
In Jacobi's method approximation calculated is used for
Nothing
Calculating the next approximation
Replaced by previous one
All above

In Gauss Seidel method approximation calculated is replaced by previous one
True
False

Relaxation method is derived by
South well
Not defined

Power method is applicable for only
Real metrics

Emraan Haakim.03349666619

Symmetric
Unsymmetrical
Both symmetric and real

The process of eliminating value of y for intermediate value of x is know as interpolation
True
False

In Richardson's extrapolation method, we usually use two different step sizes ……and …… to yield a higher order method.

h, h/2
h, h/3
h, h/4
None

In Simpson's 3/8 rule, we divide the interval of integration into n sub-intervals. Where n is divisible by.....

3
4
5
None

1-Generally, Adams methods are superior if output at many points is needed.

- True
- False

2- Euler's method is only useful for a few steps and small step sizes; however Euler's method together with Richardson extrapolation may be used to increase the _____.

- **order and accuracy**
- divergence

3- The first lngrange polynomial with equally spaced nodes produced the formula for_____.

Emraan Haakim.03349666619

- Simpson's rule
- **Trapezoidal rule**
- Newton's method
- Richardson's method

4- The need of numerical integration arises for evaluating the indefinite integral of a function that has no explicit antiderivative or whose antiderivative is not easy to obtain.

- TRUE
- **FALSE**

5- The Trapezoidal Rule is an improvement over using rectangles because we have much less "missing" from our calculations. We used_____to model the curve in trapezoidal Rule

- **straight lines**
- curves
- parabolas
- constant

6- The Euler method is numerically unstable because of _____ convergence of error.
- Slow
- Fast
- Moderate
- No

7- Adams – Bashforth is a multistep method.

- **True**
- False

8- The need of numerical integration arises for evaluating the definite integral of a function that has no explicit _____or whose antiderivative is not easy to obtain.

- Antiderivative
- **Derivatives**

Ehraan Haakim.03349666619

9- In Runge – Kutta Method, we do not need to calculate higher order derivatives and find greater accuracy.

- **True**
- False

10- An indefinite integral may_____in the sense that the limit defining it may not exist.

- **Diverge**
- Converge

11- The Trapezoidal Rule is an improvement over using rectangles because we have much less "missing" from our calculations. We used_____to model the curve in trapezoidal Rule.

- **straight lines**
- curves
- parabolas
- constant

12- An improper integral is the limit of a definite integral as an endpoint of the interval of integration approaches either a specified real number or 8 or -8 or, in some cases, as both endpoints approach limits.

- True
- False

13- Euler's Method numerically computes the approximate derivative of a function.

- True
- False

14- If we wanted to find the value of a definite integral with an infinite limit, we can instead replace the infinite limit with a variable, and then take the limit as this variable goes to_____.

Emraan Haakim.03349666619

- Constant
- Finite
- Infinity
- zero

Question : While solving a system of linear equations, which of the following approach is economical for the computer memory?
Select correct option:

Direct
Iterative
Analytical
Graphical

Question : The basic idea of relaxation method is to reduce the largest residual to ……
Select correct option:

One
Two
Zero
None of the given choices

Question: The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its _____.
Select correct option:
main diagonal
last column
last row
first row

Question: If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix , the eigenvalues of A are the diagonal entries of A.
Select correct option:
TRUE
FALSE

Emraan Haakim.03349666619

Question : A 3 x 3 identity matrix have three and different eigen values.
Select correct option:
TRUE
FALSE

Question : Which of the following is a reason due to which the LU decomposition of the system of linear equations; x+y = 1, x+y =2 is not possible?
Select correct option:
- Associated coefficient matrix is singular
- All values of l's and u's can't be evaluated
- Determinant of coefficient matrix is zero
- All are equivalent

Question : Gauss - Jordan Method is similar to …….
Select correct option:
- Gauss–Seidel method
- Iteration's method
- Relaxation Method
- Gaussian elimination method

Question : While using Relaxation method, which of the following is the largest Residual for 1st iteration on the system; 2x+3y = 1, 3x +2y = - 4 ?
Select correct option:
- -4
- 3
- 2
- 1

Question : Gauss–Seidel method is also known as method of …….
Select correct option:
- Successive displacement
- Iterations
- False position
- None of the given choices

Emraan Hakim.03349666619

Question : Jacobi's Method is a/an……
Select correct option:

Iterative method
Direct method

Question : The characteristics polynomial of a 3x 3 identity matrix is_____, if x is the eigen values of the given 3 x 3 identity matrix. where symbol ^ shows power.
Select correct option:

(x-1)^3
(x+1)^3
x^3-1
x^3+1

Question : The Power method can be used only to find the eigenvalue of A that is largest in absolute value—we call this eigenvalue the dominant eigenvalue of A.
Select correct option:

TRUE
FALSE

Question: In.....method, a system is reduced to an equivalent diagonal form using elementary transformations.
Select correct option:

Jacobi's
Gauss-Seidel
Relaxation
Gaussian elimination

Question : The linear equation: 2x+0y-2=0 has--------- solution/solutions.
Select correct option:

unique
no solution
infinite many
finite many

Question : Under elimination methods, we consider, Gaussian elimination and..... methods.
Select correct option:

Emraan Haakim.03349666619

Gauss-Seidel
Jacobi
Gauss-Jordan elimination
None of the given choices

Question : Which of the following method is not an iterative method?
Select correct option:

Jacobi's method
Gauss-Seidel method
Relaxation methods
Gauss-Jordan elimination method

Question : An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.
Select correct option:
TRUE
FALSE

Question : Exact solution of 2/3 is not exists.
Select correct option:
TRUE

FALSE

Question : When the condition of diagonal dominance becomes true in Jacobi's Method. Then its means that the method is …….
Select correct option:

Stable
Unstable
Convergent
Divergent

Question : Gauss–Seidel method is similar to …….
Select correct option:

Emraan Haakim.03349666619

Iteration's method
Regula-Falsi method
Jacobi's method
None of the given choices

Question : Sparse matrices arise in computing the numerical solution of …….
Select correct option:

Ordinary differential equations
Partial differential equations
Linear differential equations
Non-linear differential equations

Question : While solving by Gauss-Seidel method, which of the following is the first Iterative solution for the system; x-2y =1, x+4y=4 ?
Select correct option:

(1, 0.75)
(0,0)
(1,0)
(0,1)

Question: While solving a system of linear equations by Gauss Jordon Method, after all the elementary row operations if there lefts also zeros on the main diagonal then which of the is true about the system?
Select correct option:

System may have unique solutions
System has no solution
System may have multiple numbers of finite solutions
System may have infinite many solutions

Question: Numerical methods for finding the solution of the system of equations are classified as direct and..... methods
Select correct option:

Emraan Haakim.03349666619

Indirect
Iterative
Jacobi
None of the given choices

Question : If the Relaxation method is applied on the system; 2x+3y = 1, 3x +2y = - 4, then largest residual in 1st iteration will reduce to----- --.
Select correct option:
zero
4
-1
-1

Question : While using Relaxation method, which of the following is the Residuals for 1st iteration on the system; 2x+3y = 1, 3x +2y =4 ?
Select correct option:
(2,3)
(3,-2)
(-2,3)
(1,4)

Question : If the order of coefficient matrix corresponding to system of linear equations is 3*3 then which of the following will be the orders of its decomposed matrices; 'L' and 'U'?
Select correct option:
Order of 'L' = 3*1, Order of 'U' = 1*3
Order of 'L' = 3*2, Order of 'U' = 2*3
Order of 'L' = 3*3, Order of 'U' = 3*3
Order of 'L' = 3*4, Order of 'U' = 4*3

Question : While solving the system; x–2y = 1, x+4y = 4 by Gauss-Seidel method, which of the following ordering is feasible to have good approximate solution?
Select correct option:
x+4y = 1, x-2y = 4
x+2y = 1, x- 4y =4
x+4y = 4, x–2y = 1
no need to reordering

Imran Hakim.03349666619

Question : Full pivoting, in fact, is more ..... than the partial pivoting.
Select correct option:

Easiest
Complicated

Question : Gauss–Seidel method is also known as method of ……..
Select correct option:
Successive displacement
Iterations
False position
None of the given choices

Question : For the equation $x^3 + 3x - 1 = 0$, the root of the equation lies in the interval.....

- (1, 3)
- (1, 2)
- (0, 1)
- (1, 2)

Question :..... lies in the category of iterative method.

- Bisection Method
- Regula Falsi Method
- Secant Method
- all of the given choices

Question : Power method is applicable if the eigen vectors corresponding to eigen values are linearly independent.
True
1. false

Emraan Haakim.03349666619

Question: A 3 x 3 identity matrix have three and different eigen values.

1. True

False

Question : If n x n matrices A and B are similar, then they have the different eigenvalues (with the same multiplicities).

1. True

False

Question : The Jacobi's method is a method of solving a matrix equation on a matrix that has_____zeros along its main diagonal.

No

1. At least one

Question : An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to_____.

Unity

1. zero

Question : If the root of the given equation lies between a and b, then the first approximation to the root of the equation by bisection method is ……

- $\frac{(a+b)}{2}$
- $\frac{(a-b)}{2}$
- $\frac{(b-a)}{2}$
- None of the given choices

Question : To apply Simpson's 3/8 rule, the number of intervals in the following must be

- 10

Emraan Haakm.03349666619

▶ 11
▶ 12
▶ 13

Question : The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric_____definite matrices A.
Select correct option:
positive
negative
Question : Differences methods find the_____solution of the system.
Select correct option:
numerical
Analytical
Question : To apply Simpson's 1/3 rule, the number of intervals in the following must be

▶ 2 (**Simpson**"s **1/3 rule must use** an even **number** of elements')
▶ 3
▶ 5
▶ 7

Question : The Power method can be used only to find the eigenvalue of A that is largest in absolute value we call this eigenvalue the dominant eigenvalue of A.
Select correct option:

TRUE
FALSE

Question : The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its_____.
Select correct option:

main diagonal
last column
last row
first row

Emraan Hakim.03349666619

Question : Bisection and false position methods are also known as bracketing method and are always
Divergent
Convergent

Question : The Inverse of a matrix can only be found if the matrix is
Singular
Every square non-singular matrix will have an inverse.
Scalar
Diagonal
Question : In interpolation is used to represent the δ

Forward difference
Central difference
Backward difference

Question : The base of the decimal system is _____
10
0
2
8
None of the above.

Question : Bisection method is ..... method
► Open Method
► **Bracketing Method**
Question : Exact solution of 2/3 is not exists.
TRUE
FALSE
Question : The Jacobi's method is a method of solving a matrix equation on a matrix that has____zeros along its main diagonal.
No
atleast one

Emran Haakim.03349666619

Question: A 3 x 3 identity matrix have three and _____eigen values.

same

different

Question : Eigenvalues of a symmetric matrix are all_____.

real

complex

zero

positive

Question : The Jacobi iteration converges, if A is strictly diagonally dominant.

TRUE

FALSE

Question : Below are all the finite difference methods EXCEPT _____.

jacobi's method

newton's backward difference method

Stirlling formula

Forward difference method

Question: If n x n matrices A and B are similar, then they have the same eigenvalues (with the same multiplicities).

TRUE

FALSE

Question : If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix , the eigenvalues of A are the diagonal entries of A.

TRUE

FALSE

Question: The characteristics polynomial of a 3x 3 identity matrix is _____, if x is the eigen values of the given 3 x 3 identity matrix. where symbol ^ shows power.

(x-1)^3

(x+1)^3

x^3-1

x^3+1

Question : Two matrices with the same characteristic polynomial need not be similar.

TRUE

FALSE

Imraan Haakim 03349666619

Question : The determinant of a diagonal matrix is the product of the diagonal elements.

True

1. False

Qusetion : The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric positive definite matrices A.

True

1. False

Question : The determinant of a_____matrix is the product of the diagonal elements.

Diagonal

1. Upper triangular
2. Lower triangular
3. Scalar

Question : For differences methods we require the set of values.

True

False

Question : If x is an eigen value corresponding to eigen value of V of a matrix A. If a is any constant, then x – a is an eigen value corresponding to eigen vector V is an of the matrix A - a I.

True

False

Question : Central difference method seems to be giving a better approximation, however it requires more computations.

True

False

Question : Iterative algorithms can be more rapid than direct methods.

True

1. False

Question : Central Difference method is the finite difference method.

True

Emraan Hakim.03349666619

1. False

Question : Back substitution procedure is used in **…….**
Select correct option:
Gaussian Elimination Method
Jacobi's method
Gauss-Seidel method
None of the given choices
Question : The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its main diagonal.

True
False1 .

Question: The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its_____.
•
main diagonal
last column
last row
first row
Question: . An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to_____.
Unity
Zero
Question: An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.

TRUE
FALSE
Question: . The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric positive definite matrices A.

True

Emraan Haakim.03349666619

By.Mohammad Imraan Haakim 03349666619
emranhaakim@gmail.comSoftvission Collage of IT Bahawalpur

False

Question : The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric_____definite matrices A.

Pos I t ive
Negative
Q u e s t i o n : . T h e d e t e r m i n a n t o f a d i a g o n a l m a t r i x i s t h e p r o d u c t o f t h e d i a g o n a l e l e m e n t s .

True
False1
Question : Power method is applicable if the eigen vectors corresponding to eigen values are linearly independent.

True
False
Question : Power method is applicable if the eigen values are _____.

real and distinct
real and equal
positive and distinct
negative and distinct

Question : Simpson's rule is a numerical method that approximates the value of a definite integral by using polynomials.

Quadratic
Linear
Cubic
Quartic

Q u e s t i o n : . I n S i m p s o n ' s R u l e , w e u s e p a r a b o l a s t o a p p r o x i m a t i n g e a c h p a r t o f t h e c u r v e . T h i s p r o v e s to be very efficient as compared to Trapezoidal rule.

True

Emraan Haakim.03349666619

By.Mohammad Imraan Haakim 03349666619
emranhaakim@gmail.comSoftvission Collage of IT Bahawalpur

False
Question : The predictor-corrector method an implicit method. (multi-step methods)

True
False
Question: Generally, Adams methods are superiorifoutputatmanypointsisneeded.

True
False

Question : The Trapezoidal rule is a numerical method that approximates the value of a._____.

Indefinite integral
Definite integral
Improper integral
Function
Question : The need of numerical integration arises for evaluating the definite integral of a function that has no explicit_____or whose antiderivative is not easy to obtain.

Anti deri vat ive
Derivatives.
Question: .Anindefiniteintegral may _____ in the sensethattheli mitdefining itmay notexist.

diverge
Converge
Question : An improper integral is the limit of a definite integral as an endpoint of the interval of integration approaches either a specified real number or ∞ or -∞ or, in some cases, as both endpoints approach limits.

TRUE
FALSE
Question : Euler's Method numerically computes the approximate derivative of a function.

Emraan Haakim 03349666619

TRUE
FALSE
Question :. Euler's Method numerically computes the approximate _____ of a function.

Antiderivative
Derivative
Error
Value
Question: If we wanted to find the value of a definite integral with an infinite limit, we can instead replace the infinite limit with a variable, and then take the limit as this variable goes to_____.
Chose the correct option :
Constant
Finite
Infinity
Zero

Question : Euler's Method numerically computes the approximate derivative of a function.
•
TRUE
•
FALSE

Question: .The Jacobi iteration _____, if A is strictly diagonally dominant.

converges
Diverges
Question :. Two matrices with the same characteristic polynomial need not be similar.

TRUE
fALSE

Emraan Haakim.03349666619

Question :. Differences methods find the _____ solution of the system.

Numerical

Analytica

Question : .By using determinants, we can easily check that the solution of the given system of linear equation exits and it is unique.

TRUE

FALSE

Question : The absolute value of a determinant (|detA|) is the product of the absolute values of the eigen values of matrix A

TRUE

FALSE

Question : Eigenvectors of a symmetric matrix are orthogonal, but only for distinct eigenvalues.

TRUE

FALSE

.

Question : Let A be an n ×n matrix. The number x is an eigenvalue of A if there exists a non-zerovector v such that_____.

Av = xv

Ax = xv not shore

Av + xv=0

Av = Ax1

Question : In Jacobi's Method, the rate of convergence is quite _____ compared with other methods.

slow

Fast

Question : .Numerical solution of 2/3 up to four decimal places is_____.

Emraan Haakim.03349666619

0.667
0.6666
0.6667
0.666671 .

Question : Symbol used for forward differences is

Δ Correct
δ
μ

Question:. There lationship between central difference operator and the shift operator is given by

$\delta = E - E^{-1}$

$\delta = E + E^{-1}$

$\delta = E^{1/2} + E^{1/2}$

$\delta = E^{1/2} - E^{1/2}$

Question : Muller's method requires -------- starting points

1

2

3

Question : By using determinants, we can easily check that the solution of the given system of linear equation_____and it is_____.

Select correct option:
exits, unique
exists, consistent
trivial, unique
nontrivial, inconsistent

Emraan Haakim.03349666619

Question : Two matrices with the_____characteristic polynomial need not be similar.

Select correct option:
same
different

Question : In..... method, the elements above and below the diagonal are simultaneously made zero.

Select correct option:
Jacobi's
Gauss-Seidel
Gauss–Jordon Elimination
Relaxation

Question : Which of the following is equivalent form of the system of equations in matrix form; AX=B ?

Select correct option:
XA = B
X = B(Inverse of A)
X =(Inverse of A)B
BX = A

Question : If the determinant of a matrix A is not equal to zero then the system of equations will have…….

Select correct option:
a unique solution
many solutions
infinite many solutions
None of the given choices

Question : Sparse matrix is a matrix with …….

Select correct option:
Some elements are zero
Many elements are zero
Some elements are one

Emraan Haakim.03349666619

Many elements are one

Question : An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.
Select correct option:

TRUE
FALSE

Question # 1 of 10 (Start time: 11:14:39 PM) Total Marks: 1
The Jacobi iteration_____, if A is strictly diagonally dominant.

Select correct option:
converges
diverges

Question # 2 of 10 (Start time: 11:16:04 PM) Total Marks: 1
The Jacobi's method is a method of solving a matrix equation on a matrix that has_____zeros along its main diagonal.

Select correct option:
No
atleast one

Question # 3 of 10 (Start time: 11:17:14 PM) Total Marks: 1
Power method is applicable if the eigen vectors corresponding to eigen values are linearly_____.

Select correct option:
independent
dependent

Question # 4 of 10 (Start time: 11:17:42 PM) Total Marks: 1
Power method is applicable if the eigen values are_____.

Select correct option:
real and distinct
real and equal
positive and distinct
negative and distinct

Emraan Haakim.03349666619

Question # 7 of 10 (Start time: 11:19:55 PM) Total Marks: 1
The determinant of a diagonal matrix is the product of the diagonal elements.

Select correct option:
TRUE
FALSE

Question # 8 of 10 (Start time: 11:21:14 PM) Total Marks: 1
For differences methods we require the set of values.

Select correct option:
TRUE
FALSE

Question # 10 of 10 (Start time: 11:23:55 PM) Total Marks: 1
Two matrices with the_____ characteristic polynomial need not be similar.

Select correct option:
Same
different

Question # 1 of 10 Total Marks: 1
While using Relaxation method, which of the following is the Residuals for 1st iteration on the system; 2x+3y = 1, 3x +2y =4 ?

Select correct option:
(2,3)
(3,-2)
(-2,3)
(1,4)

Question # 2 of 10 (Start time: 11:14:32 PM) Total Marks: 1
Sparse matrices arise in computing the numerical solution of …….

Select correct option:
Ordinary differential equations
Partial differential equations

Emraan Haakim.03349666619

Linear differential equations
Non-linear differential equations

Question # 3 of 10 (Start time: 11:15:18 PM) Total Marks: 1
In.....method, the elements above and below the diagonal are simultaneously made zero.

Select correct option:
Jacobi's
Gauss-Seidel
Gauss–Jordon Elimination
Relaxation

Question # 5 of 10 (Start time: 11:17:54 PM) Total Marks: 1
Which of the following is equivalent form of the system of equations in matrix form; AX=B ?

Select correct option:
XA = B
X = B(Inverse of A)
X =(Inverse of A)B
BX = A

Question # 7 of 10 (Start time: 11:20:24 PM) Total Marks: 1
If the determinant of a matrix A is not equal to zero then the system of equations will have…….

Select correct option:
A unique solution
many solutions
infinite many solutions
None of the given choices

Question # 8 of 10 (Start time: 11:21:37 PM) Total Marks: 1
Sparse matrix is a matrix with …….

Select correct option:
Some elements are zero

Ehtaam Haakim.03349666619

Many elements are zero
Some elements are one
Many elements are one

Question # 4 of 10 (Start time: 11:31:21 PM) Total Marks: 1
Back substitution procedure is used in …….

Select correct option:
Gaussian Elimination Method
Jacobi's method
Gauss-Seidel method
None of the given choices

Question # 5 of 10 (Start time: 11:32:12 PM) Total Marks: 1
The linear equation: 2x+0y-2=0 has--------- solution/solutions.

Select correct option:
unique
no solution
infinite many
finite many

Question # 8 of 10 (Start time: 11:35:30 PM) Total Marks: 1
For a system of linear equations, the corresponding coefficient matrix has the value of determinant; |A| = 0, then which of the following is true?

Select correct option:
The system has unique solution
The system has finite multiple solutions
The system has infinite may solutions
The system has no solution

Question # 9 of 10 (Start time: 11:36:21 PM) Total Marks: 1
For the system; 2x+3y = 1, 3x +2y = - 4, if the iterative solution is (0,0) and 'dxi = 2' is the increment in 'y' then which of the following will be taken as next iterative solution?

Select correct option:
(2,0)
(0,3)

Emran Hakim.03349666619

(0,2)
(1,-4)

Question # 2 of 10 (Start time: 11:42:14 PM)Total Marks: 1
Which of the following method is not an iterative?

Select correct option:
Gauss–Seidel method
Iteration's method
Relaxation Method
Gauss Jordan method

Question # 3 of 10 (Start time: 11:43:46 PM)Total Marks: 1
Sparse matrix is a matrix with …….

Select correct option:
Some elements are zero
Many elements are zero
Some elements are one
Many elements are one

Question # 4 of 10 (Start time: 11:44:33 PM)Total Marks: 1
While using Relaxation method, which of the following is the Residuals for 1st iteration on the system; 2x+3y = 1, 3x +2y =4

Select correct option:
(2,3)
(3,-2)
(-2,3)
(1,4)

Emraan Hakim.03349666619

Question # 6 of 10 (Start time: 11:47:15 PM)Total Marks: 1
Relaxation Method is a/an …….

Select correct option:

Direct method
Iterative method

Question # 9 of 10 (Start time: 11:50:33 PM)Total Marks: 1
Full pivoting, in fact, is more .....than the partial pivoting.

Select correct option:
Easiest
Complicated

Question # 10 of 10 (Start time: 11:51:55 PM)Total Marks: 1
Gauss–Seidel method is also known as method of …….

Select correct option:
Successive displacement
Iterations
False position
None of the given choices

Question # 2 of 10 (Start time: 11:31:28 PM) Total Marks: 1
Iterative algorithms can be more rapid than direct methods.

Select correct option:
FALSE
TRUE

Question # 3 of 10 (Start time: 11:32:02 PM) Total Marks: 1
Below are all the finite difference methods EXCEPT_____.

Select correct option:
jacobi's method
newton's backward difference method
Stirlling formula
Forward difference method

Question # 2 of 10 Total Marks: 1
Sparse matrices arise in computing the numerical solution of …….

Select correct option:
Ordinary differential equations
Partial differential equations
Linear differential equations

Imran Haakim.03349666619

Non-linear differential equations

Question # 9 of 10
If x is an eigen value corresponding to eigen value of V of a matrix A. If a is any constant, then x – a is an eigen value corresponding to eigen vector V is an of the matrix A - a I.

Select correct option:
TRUE
FALSE

Question # 10 of 10
An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.

Select correct option:
TRUE
FALSE

Question No: 1 (Marks: 1) - Please choose one
Symbol used for forward differences is

- ▶ ∇
- ▶ Δ
- ▶ □
- ▶ □

Question No: 2 (Marks: 1) - Please choose one
The relationship between central difference operator and the shift operator is given by

- ▶ □ $= E - E^{-1}$
- ▶ □ $= E + E^{-1}$
- ▶ □ $= E^{\frac{1}{2}} + E^{-\frac{1}{2}}$
- ▶ □ $= E^{\frac{1}{2}} - E^{-\frac{1}{2}}$

Question No: 3 (Marks: 1) - Please choose one
Muller's method requires-------- starting points

Emraan Haakim.03349666619

- 1
- 2
- **3**
- 4

Question No: 4 (Marks: 1) - Please choose one

If S is an identity matrix, then

- $S^{-1} = S$
- $S^t = S$
- $All\ are true$
- $S^{-1} = S^t$

Question No: 5 (Marks: 1) - Please choose one

If we retain r+1 terms in Newton's forward difference formula, we obtain a polynomial of degree----- agreeing with y_x at $x_0, x_1, \ldots, x_r$

- r+2
- r+1
- **r**
- r-1

Question No: 6 (Marks: 1) - Please choose one

P in Newton's forward difference formula is defined as

- $p = \left(\frac{x - x_0}{h}\right)$
- $p = \left(\frac{x + x_0}{h}\right)$
- $p = \left(\frac{x + x_n}{h}\right)$
- $p = \left(\frac{x - x_n}{h}\right)$

Question No: 7 (Marks: 1) - Please choose one

Octal number system has the base ----------

- 2
- **8**

Emraan Haakim.03349666619

- 10
- 16

Question No: 8 (Marks: 1) - Please choose one

Newton's divided difference interpolation formula is used when the values of the independent variable are

- Equally spaced
- **Not equally spaced**
- Constant
- None of the above

Question No: 9 (Marks: 1) - Please choose one

Given the following data

x	0	1	2	4
$f(x)$	1	1	2	5

Value of $f(2, 4)$ is

- 1.5
- **3**
- 2
- 1

Question No: 10 (Marks: 1) - Please choose one

If $y(x)$ is approximated by a polynomial $p_n(x)$ of degree n then the error is given by

- $\square(x) = y(x) + P_n(x)$
- $\square(x) = y(x) - P_n(x)$
- $\square(x) = P_n(x) - y(x)$
- $\square(x) = y(x) \times P_n(x)$

Question No: 11 (Marks: 1) - Please choose one

Let I denotes the closed interval spanned by $x_0, x_1, x_2, x_3, x_4, x_5, x_6, \bar{x}_7, x$. Then $F(x)$ vanishes ------times in the interval I.

Emraan Haakim.03349666619

- n-1
- **n+2**
- n
- n+1

Question No: 12 (Marks: 1) - Please choose one

Differential operator in terms of forward difference operator is given by

- $D = \frac{1}{h}(\Delta + \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} + \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} + ...)$
- $D = \frac{1}{h}(\Delta + \frac{\Delta^2}{2} + \frac{\Delta^3}{3} + \frac{\Delta^4}{4} + \frac{\Delta^5}{5} + ...)$
- $D = \frac{1}{h}(\Delta - \frac{\Delta^2}{2} + \frac{\Delta^3}{3} - \frac{\Delta^4}{4} + \frac{\Delta^5}{5} - ...)$
- $D = \frac{1}{h}(\Delta - \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} - \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} - ...)$

Question No: 13 (Marks: 1) - Please choose one

Finding the first derivative of $f(x)$ at x =0.4 from the following table:

x	0.1	0.2	0.3	0.4
$f(x)$	1.10517	1.22140	1.34986	1.49182

Differential operator in terms of ---------- will be used.

- Forward difference operator
- **Backward difference operator**
- Central difference operator
- None of the given choices

Question No: 14 (Marks: 1) - Please choose one

For the given table of values

x	0.1	0.2	0.3	0.4	0.5	0.6
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f'(0.1)$, using two-point equation will be calculated as.....

Emraan Haakim.03349666619

- -0.5
- **0.5**
- 0.75
- -0.75

Question No: 15 (Marks: 1) - Please choose one

In Simpson's 1/3 rule, $f(x)$ is of the form

- $ax+b$
- ax^2+bx+c
- ax^3+bx^2+cx+d
- $ax^4+bx^3+cx^2+dx+e$

Question No: 16 (Marks: 1) - Please choose one

$$I=\int_a^b f(x)dx$$

While integrating , h , width of the interval, is found by the formula----.

- $\frac{b-a}{n}$
- $\frac{b+a}{n}$
- $\frac{a-b}{n}$
- None of the given choices

Question No: 17 (Marks: 1) - Please choose one

To apply Simpson's 1/3 rule, valid number of intervals are.....

- 7
- **8**
- 5
- 3

Question No: 18 (Marks: 1) - Please choose one

For the given table of values

x	02	0.3	0.4	0.5	0.6	0.7
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

Emraan Haakim.03349666619

$f''(0.2)$, using three-point equation will be calculated as ……

- ▶ 17.5
- ▶ 12.5
- ▶ 7.5
- ▶ **-12.5**

Question No: 19 (Marks: 1) - Please choose one

To apply Simpson's 1/3 rule, the number of intervals in the following must be

- ▶ **2**
- ▶ 3
- ▶ 5
- ▶ 7

Question No: 20 (Marks: 1) - Please choose one

To apply Simpson's 3/8 rule, the number of intervals in the following must be

- ▶ 10
- ▶ 11
- ▶ **12**
- ▶ 13

Question No: 21 (Marks: 1) - Please choose one

If the root of the given equation lies between a and b, then the first approximation to the root of the equation by bisection method is ……

- ▶ $\frac{(a+b)}{2}$
- ▶ $\frac{(a-b)}{2}$
- ▶ $\frac{(b-a)}{2}$
- ▶ None of the given choices

Question No: 22 (Marks: 1) - Please choose one

.....lies in the category of iterative method.

- ▶ Bisection Method

Emraan Haakim.03349666619

► Regula Falsi Method
► Secant Method
► **All the given choices**

Question No: 23 (Marks: 1) - Please choose one

For the equation $x^3 + 3x - 1 = 0$, the root of the equation lies in the interval.....

► (1, 3)
► (1, 2)
► **(0, 1)**
► (1, 2)

Question No: 24 (Marks: 1) - Please choose one

Rate of change of any quantity with respect to another can be modeled by

► **An ordinary differential equation**
► A partial differential equation

► A polynomial equation

► None of the given choices

Question No: 25 (Marks: 1) - Please choose one

If
$\frac{dy}{dx} = f(x, y)$
Then the integral of this equation is a curve in

► None of the given choices

► xt-plane
► **yt-plane**
► **xy-plane**

Question No: 26 (Marks: 1) - Please choose one

In solving the differential equation
$y' = x + y \; ; y(0.1) = 1.1$
$h = 0.1$, By Euler's method $y(0.2)$ is calculated as

► 1.44
► 1.11
► **1.22**

Emraan Haakim.03349666619

► 1.33

Question No: 27 (Marks: 1) - Please choose one

In second order Runge-Kutta method

k_1 is given by

► $k_1 = hf(x_n, y_n)$
► $k_1 = 2hf(x_n, y_n)$
► $k_1 = 3hf(x_n, y_n)$
► None of the given choices

Question No: 28 (Marks: 1) - Please choose one

In fourth order Runge-Kutta method, k_2 is given by

► $k_2 = hf(x_n + \frac{h}{2}, y_n + \frac{k_1}{2})$
► $k_2 = hf(x_n + \frac{h}{3}, y_n + \frac{k_1}{3})$
► $k_2 = hf(x_n - \frac{h}{3}, y_n - \frac{k_1}{3})$
► $k_2 = hf(x_n - \frac{h}{2}, y_n - \frac{k_1}{2})$

Question No: 29 (Marks: 1) - Please choose one

In fourth order Runge-Kutta method, k_4 is given by

► $k_3 = hf(x_n + 2h, y_n + 2k_3)$
► $k_3 = hf(x_n - h, y_n - k_3)$
► $k_3 = hf(x_n + h, y_n + k_3)$
► None of the given choices

Question No: 30 (Marks: 1) - Please choose one

Adam-Moulton P-C method is derived by employing

► **Newton's backward difference interpolation formula**
► Newton's forward difference interpolation formula
► Newton's divided difference interpolation formula

Emraan Haakim.03349666619

▶ None of the given choices

Mth603 Solved MCQS for Final Term Exam

Solved by Mermaid with reference of book

Exact solution of 2/3 is not exists.
TRUE
FALSE

The Jacobi's method is
A method of solving a matrix equation on a matrix that has_____zeros along its main diagonal.

No
At least one

A 3 x 3 identity matrix have three and_____eigen values.
Same
Different

Eigenvalues of a symmetric matrix are all_____.
Real
Complex
Zero
Positive

The Jacobi iteration converges, if A is strictly diagonally dominant.
TRUE
FALSE

Below are all the finite difference methods EXCEPT_____.

Jacobi's method
Newton's backward difference method
Stirling formula
Forward difference method

If n x n matrices A and B are similar, then they have the same eigenvalues (with the same multiplicities).
TRUE
FALSE

Emraan Haakim.03349666619

If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix, the eigenvalues of A are the diagonal entries of A.

TRUE
FALSE

The characteristics polynomial of a 3x 3 Identity matrix is_____, if x is the Eigen values of the given 3 x 3 identity matrix. Where symbol ^ shows power.

(X-1)^3
(x+1)^3
X^3-1
X^3+1

Two matrices with the same characteristic polynomial need not be similar.

TRUE
FALSE

Bisection method is a

Bracketing method
Open method

Regula Falsi means

Method of Correct position
Method of unknown position
Method of false position
Method of known position

Eigenvalues of a symmetric matrix are all_____.
Select correct option:

Real
Zero
Positive
Negative

An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.
Select correct option:

TRUE
FALSE

Emraan Haakim.03349666619

Exact solution of 2/3 is not exists.
Select correct option:

TRUE
FALSE

The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric _____definite matrices A.
Select correct option:

Positive
Negative

Differences methods find the_____solution of the system.
Select correct option:

Numerical
Analytical

The Power method can be used only to find the eigenvalue of A that is largest in absolute value—we call this Eigenvalue the dominant eigenvalue of A.
Select correct option:

TRUE
FALSE

The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its_____.
Select correct option:

Main diagonal
Last column
Last row
First row

Emraan Haakim.03349666619

If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix , the eigenvalues of A are the diagonal entries of A.
Select correct option:

TRUE
FALSE

A 3 x 3 identity matrix have three and different Eigen values.
Select correct option:

TRUE
FALSE

Newton Raphson method falls in the category of

Bracketing method
Open Method
Iterative Method
Indirect Method

Newton Raphson method is also known as
Tangent Method
Root method
Open Method
Iterative Method

Secant Method uses values for approximation

1
3
2
4

Secant Method is than bisection method for finding root
Slow
Faster

In Newton Raphson method

Root is bracketed
Root is not bracketed

Emraan Haakim.03349666619

Regula falsi method and bisection method are both

Convergent
Divergent

In bisection method the two points between which the root lies are

Similar to each other
Different
Not defined
Opposite

In which methods we do not need initial approximation to start
Indirect Method
Open Method
Direct Method
Iterative Method

Root may be

Complex
Real
Complex or real
None

In Regula falsi method we choose points that have signs

2 points opposite signs
3 points opposite signs
2 points similar signs
None of the given

In a bounded function values lie between
1 and -1
1 and 2
0 and 1
0 and -2

Newton Raphson method is a method which when it leads to division of number close to zero
Diverges
Converges

Which of the following method is modified form of Newton Raphson Method?
Regula falsi method

Fmraan Haakim.03349666619

Bisection method
Secant method
Jacobi's Method

Which 1 of the following is generalization of Secant method?
Muller's Method
Jacobi's Method
Bisection Method
N-R Method

Secant Method needs starting points

2
3
4
1
Near a simple root Muller's Method converges than the secant method

Faster
Slower

If S is an identity matrix, then

$S^{-1} = S$
$S^t = S$
$S^{-1} = S^t$
All are true

If we retain r+1 terms in Newton's forward difference formula, we obtain a polynomial of degree ---- agreeing with y_x at $x_0, x_1, \ldots, x_r$

r+2
r+1
R
R-1
P in Newton's forward difference formula is defined as

$$p = \left(\frac{x - x_0}{h}\right)$$

Emraan Haakim.03349666619

$$p = (\frac{x + x_0}{h})$$

$$p = (\frac{x + x_n}{h})$$

$$p = (\frac{x - x_n}{h})$$

Octal numbers has the base

10
2
8
16

Newton's divided difference interpolation formula is used when the values of the independent variable are

Equally spaced

Not equally spaced

Constant
None of the above

Given the following data

x	0	1	2	4
$f(x)$	1	1	2	5

Value of $f(2,4)$ is

1.5

3
2

1

If $y(x)$ is approximated by a polynomial $p_n(x)$ of degree n then the error is given by

$$\square(x) = y(x) + \underline{P}_n(x)$$

Emraan Haakim.03349666619

$\square(x) = y(x) - P_n(x)$

$\square(x) = y(x) \times P_n(x)$

$\square(x) = P_n(x) + y(x)$

Let I denotes the closed interval spanned by $x_0, x_1, x_2, x_3, x_4, x_5, x_6, \bar{x}_7, x$. Then $F(x)$ vanishes ------times in the interval I.

N-1
N+2
N
N+1

Differential operator in terms of forward difference operator is given by

$$D = \frac{1}{h}(\Delta + \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} + \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} + \ldots)$$

$$D = \frac{1}{h}(\Delta + \frac{\Delta^2}{2} + \frac{\Delta^3}{3} + \frac{\Delta^4}{4} + \frac{\Delta^5}{5} + \ldots)$$

$$D = \frac{1}{h}(\Delta - \frac{\Delta^2}{2} + \frac{\Delta^3}{3} - \frac{\Delta^4}{4} + \frac{\Delta^5}{5} \ldots)$$

$$D = \frac{1}{h}(\Delta - \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} - \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} \ldots)$$

Finding the first derivative of $f(x)$ at x =0.4 from the following table:

x	0.1	0.2	0.3	0.4
$f(x)$	1.10517	1.22140	1.34986	1.49182

Differential operator in terms of---------- will be used.

Forward difference operator
Backward difference operator
Central difference operator
All of the given choices

Emraan Haaim.03349666619

For the given table of values

x	0.1	0.2	0.3	0.4	0.5	0.6
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f'(0.1)$, using two-point equation will be calculated as.....

-0.5
0.5
0.75
-0.75

In Simpson's 1/3 rule, $f(x)$ is of the form

$ax+b$

- ax^2+bx+c
- ax^3+bx^2+cx+d
- $ax^4+bx^3+cx^2+dx+e$

$$I=\int_a^b f(x)dx$$

While integrating , h , width of the interval, is found by the formula-----.

$\frac{b-a}{n}$

$\frac{b+a}{n}$

$\frac{a-b}{n}$

None of the given choices

Emraan Haakim.03349666619

To apply Simpson's 1/3 rule, valid number of intervals are.....

7
8
5
3

For the given table of values

x	0.1	0.2	0.3	0.4	0.5	0.6
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f''(0.2)$ **, using three-point equation will be calculated as ……**

17.5
12.5
7.5
-12.5

To apply Simpson's 1/3 rule, the number of intervals in the following must be

2
3
5
7

To apply Simpson's 3/8 rule, the number of intervals in the following must be

10
11
12
13

If the root of the given equation lies between a and b, then the first approximation to the root of the equation by bisection method is ……

$\frac{(a+b)}{2}$

Emraan Haakim.03496666619

$\frac{(a-b)}{2}$

$\frac{(b-a)}{2}$

None of the given choices

.....lies in the category of iterative method.

Bisection Method
Regula Falsi Method
Secant Method
All of the given choices

For the equation $x^3+3x-1=0$, the root of the equation lies in the interval.....

(1, 3)
(1, 2)
(0, 1)
(1, 2)

Rate of change of any quantity with respect to another can be modeled by

An ordinary differential equation
A partial differential equation

A polynomial equation

None of the given choices

If
$\frac{dy}{dx}=f(x,y)$

Then the integral of this equation is a curve in

None of the given choices

Xt-plane
Yt-plane
Xy-plane

Emraan Hadkim.03349666619

In solving the differential equation

$$y' = x + y \; ; \; y(0.1) = 1.1$$

$h = 0.1$, **By Euler's method** $y(0.2)$ **is calculated as**

1.44
1.11
1.22
1.33

In second order Runge-Kutta method
k_1 is given by

$k_1 = hf(x_n, y_n)$
$k_1 = 2hf(x_n, y_n)$
$k_1 = 3hf(x_n, y_n)$
None of the given choices

In fourth order Runge-Kutta method, k_2 is given by

$k_2 = hf\left(x_n + \frac{h}{2}, y_n + \frac{k_1}{2}\right)$
$k_2 = hf\left(x_n + \frac{h}{3}, y_n + \frac{k_1}{3}\right)$
$k_2 = hf\left(x_n - \frac{h}{3}, y_n - \frac{k_1}{3}\right)$
$k_2 = hf\left(x_n - \frac{h}{2}, y_n - \frac{k_1}{2}\right)$

In fourth order Runge-Kutta method, k_4 **is given by**

$k_3 = hf(x_n + 2h, y_n + 2k_3)$
$k_3 = hf(x_n - h, y_n - k_3)$

Emraanhaakim.03349666619

$k_3 = hf(x_n + h, y_n + k_3)$

None of the given choices

Adam-Moulton P-C method is derived by employing

Newton's backward difference interpolation formula
Newton's forward difference interpolation formula
Newton's divided difference interpolation formula
None of the given choices

The need of numerical integration arises for evaluating the definite integral of a function that has no explicit_____or whose antiderivative is not easy to obtain

Derivatives
Antiderivative

If $A \neq 0$ then system will have a

Definite solution
Unique solution
Correct solution
No solution

If $A \neq 0$ then

There is a unique solution
There exists a complete solution
There exists no solution
None of the above options

Direct method consists of method

2
3
5
4

We consider Jacobi's method Gauss Seidel Method and relaxation method as

Direct method
Iterative method
Open method
All of the above

In Gauss Elimination method Solution of equation is obtained in

3 stages
2 stages

Emraan Haakim.03349666619

4 stages
5 stages

Gauss Elimination method fails if any one of the pivot values becomes
Greater
Small
Zero
None of the given

Changing the order of the equation is known as

Pivoting
Interpretation

Full pivoting is than partial pivoting
Easy
More complicated

The following is the variation of Gauss Elimination method

Jacobi's method
Gauss Jordan Elimination method

Courts reduction method is also known as Cholesky Reduction method
True
False

Jacobi's method is also known as method of Simultaneous displacement
True
False
Gauss Seidel method is also known as method of Successive displacement
False
True
In Jacobi's method approximation calculated is used for
Nothing
Calculating the next approximation
Replaced by previous one
All above

In Gauss Seidel method approximation calculated is replaced by previous one
True
False

Relaxation method is derived by
South well
Not defined

Imraan Haakim.03349666619

Power method is applicable for only
Real metrics
Symmetric
Unsymmetrical
Both symmetric and real

The process of eliminating value of y for intermediate value of x is know as interpolation
True
False

Question No: 31 (Marks: 2)

If $F(h) = 256.2354$ and $F(\frac{h}{2}) = 257.1379$, then find $F_1(\frac{h}{2})$ using Richardson's extrapolation limit.

Question No: 32 (Marks: 2)

Evaluate the integral

$$\int_0^{\frac{\pi}{2}} (\cos x + 2)dx$$

Using Simpson's 3/8 rule

Take h= $\frac{\pi}{4}$

Question No: 33 (Marks: 2)

Write a general formula for Modified Euler's method of solving the given differential equation.

Question No: 34 (Marks: 3)

Evaluate the integral

$$\int_0^4 x^2 dx$$

Using Trapezoidal rule
Take h=1

Question No: 35 (Marks: 3)

Emraan Haakim.03349666619

Evaluate the integral

$$\int_{3}^{5} (\log x + 2)dx$$

Using Simpson's 3/8 rule
Take h=1

Question No: 36 (Marks: 3)

Write a formula for finding the value of k_3 in Fourth-order R-K method.

Question No: 37 (Marks: 5)

Find Newton's forward difference table from the following data.

x	0.0	0.1	0.2	0.3	0.4
$f(x)$	1	0.9048	0.8187	0.7408	0.6703

Question No: 38 (Marks: 5)

Evaluate the integral

$$\int_{0}^{3} (x^2 + x)dx$$

Using Simpson's 3/8 rule

Take h=1

Question No: 39 (Marks: 5)

Use Runge-Kutta Method of order four to find the values of k_1, k_2, k_3 and k_4 for the initial value problem

$y' = \frac{1}{2}(2x^3 + y), y(1) = 2$ taking $h = 0.1$

Emran Haakim.03346666619